Published in final edited form as:

Biochem Biophys Res Commun. 2011 January 21; 404(3): 762–766. doi:10.1016/j.bbrc.2010.12.039.

Electrical stimulation of cardiomyocytes activates mitochondrial matrix metalloproteinase causing electrical remodeling*

Thomas. P Vacek, Naira Metreveli, Neetu Tyagi, Jonathan C. Vacek, Sebastian Pagni*, and Suresh C. Tyagi

Department of Physiology and Biophysics, University of Louisville School of Medicine, Louisville, Kentucky

* Division of Thoracic and Cardiovascular Surgery, University of Louisville School of Medicine, Louisville, Kentucky

Abstract

Cardiac arrhythmias, instigated by mechanical and electrical remodeling, are associated with activation of extracellular matrix metalloproteinases (MMPs). However, the connection between intracellular MMPs activation and arrhythmogenesis is not well established. Previously, we determined localization of MMP in the mitochondria using confocal microscopy. We tested the hypothesis that electrical pacing induces the activation of mitochondrial MMP (mtMMP) and is associated with myocyte mechanical dysfunction. Myocytes were isolated and field stimulated at 1 and 4 Hz. Myocyte mechanics and calcium transient was studied using Ion-Optix system. Mitochondrial MMP-9 activation was evaluated using zymography. There was a 25% increase in 1Hz and 40% increase in 4Hz stimulation. We observed an increase in mtMMP activation with increase in electrical pacing compared to 0 Hz with a significant increase ($p < 0.05$, $n = 3$). Field stimulation at 4 Hz decreased cell re-lengthening. The levels of calcium transient were reduced with increase in contraction frequency. We conclude that electrical stimulation activates mtMMP-9 that is associated with myocyte mechanical dysfunction.

Keywords

Calcium transient; electrical pacing; arrhythmia; SCD; heart failure

Introduction

Heart failure (HF) involves abnormalities in systolic and/or diastolic function and increases the propensity for reentry arrhythmias (15). Persistent increased in cardiac sympathetic drive contributes to myocardial toxicity, leading to the decline in cardiac contractility.

Homocysteine (Hcy), a sulfur-containing amino acid, whose higher concentrations in blood known as hyperhomocysteinemia (HHcy) and have been linked independently to heart disease (5;18). Homocysteine (HCY) induces interstitial cardiac fibrosis leading to heart

* A part of this study was supported by NIH grants; HL-71010, HL-74185 and HL-88012.

Address for Correspondence: Suresh C. Tyagi, PhD, Department of Physiology and Biophysics, University of Louisville School of Medicine, Louisville, KY 40202, suresh.tyagi@louisville.edu, Phone: 502-852-3381, Fax: 502-852-6239.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

failure (3). During heart failure, increase in load cause increase in oxidative stress leading to activation of Matrix metalloproteinase, s (MMPs) (22).

MMPs are membrane-bound; zinc dependent endopeptidases involved in extracellular matrix proteins and are required for ECM remodeling. Considering the large family of MMPs, MMP-2 (gelatinase A) and MMP-9 (gelatinase B) are the key ones involved in the ECM remodeling of heart with both changing demands and response to injury (24). Both are known to break down non-fibrillar-forming type IV collagen and denatured interstitial collagens (1).

MMP-9 activation in HHCY decreases the collagen/elastin ratio and increases the fibrosis between endothelium and myocytes, and is arrhythmogenic (19;21;24). MMP-9 is playing an important role in a wide variety of physiological conditions and disease states, some of which include normal development, inflammation, wound healing and repair and vascular diseases (4). We and others have shown the presence of MMP-9 in the cardiac mitochondria (mtMMP) (9;16). Hcy was shown to activate mtMMP-9 in mitochondria with an enlargement of mitochondria in addition to disruption of cristae that resulted in contractile dysfunction (14;17). MMP-9 has also been shown to destroy SERCA pump, resulting in decreased calcium transient.

Calcium is a primary regulator of contraction in heart muscle in both initial depolarization and subsequent contraction (10). Calcium-dependent MMP-9 activation has been demonstrated in cells (13). Hoit et. al showed atrial remodeling by electronic pacing that resulted in systolic and diastolic dysfunction included chamber stiffness; there was an increase in MMP-9 and decrease in its tissue inhibitor (TIMP-4) with no change in calcium pumps (6). Russo et. al showed that MMP-2 expression was significantly increased in electrically-stimulated denervated skeletal muscle compared to control (20). MMP-2 is known to be involved in fibrotic proliferation and connective tissue remodeling (11).

A decrease in contractility is undoubtedly a major contributing factor in the pathophysiology of a significant portion of the patients with heart failure. A reasonable therapeutic goal in these patients would then be to increase cardiac contractility either acutely or long term (2). Recently, we showed that novel electrical stimulation activates mtMMP-9. Based on these results, we hypothesize that increased contraction frequency induced the activation of mitochondrial MMP (mtMMP) that has already been linked to myocyte mechanical dysfunction.

Methods

Mice and experimental protocol

C57BL/6J mice procured from The Jackson Laboratory (Bar Harbor, ME) were housed in a controlled environment on 12:12-h light-dark cycle. Mice were fed with normal mouse chow. Animal experimentation was performed according to the protocols approved by the Institutional Animal Care and Use Committee of the University of Louisville. All animal care and use procedures were carried out according to the National Institutes of Health (NIH) *Guide for the Care and Use of Laboratory Animals* (NIHPub. no. 86–23, revised 1985) and the regulations of the Animal Welfare Act.

Adult ventricular myocyte isolation

Single Ventricular myocytes from the adult mice heart were isolated according to the protocol as described elsewhere (15). Perfusion buffer was prepared from basic buffer, glucose and butanedione monoxime (BDM)(120.0 mM NaCl₂, 14.7 mM KCl, 0.6 mM KH₂PO₄, 0.6 mM Na₂HPO₄, 1.2 mM MgSO₄·H₂O, 10 mM Na-HEPES, 4.6 mM NaHCO₃,

30 mM Tauren, 10 mM BDM, 5.5 mM Glucose, pH 7.0). The perfusion system was prepared as follows: The temperature of the circulation water bath was set so that the temperature of the outflow liquid at the tip of the cannula (20-g needle with the nub filed flat and smooth) is 37°C. Flow rate was set at 4.0 ml/min. Prior to the cardiomyocyte isolation, about 100 ml of distilled water was run through the system. Then the system was perfused with perfusion buffer for at least 5 minutes.

A mouse of at least 20 grams was injected with heparin (1 000 U/kg; i.p.). Then the mouse was anesthetized with 2,2,2 tribromoethanol (240 mg/kg i.p.). The chest was wiped with 70 % ethanol. A skin incision was made revealing the xiphoid process. The rib cage was completely cut starting at the xiphoid process running up the chest cavity. To avoid heart damage, the diaphragm was cut as well. The heart was secured with forceps and all vessels were cut. The aorta was cut to leave the maximal length, which is important for rapid cannulation. The dissected heart was immediately placed in a Petri dish containing ice-cold calcium free perfusion buffer (pH 7.4). To expose the aorta, all the remnant excess tissue was removed and discarded. While holding the aorta with two fine forceps, a vertically-mounted cannula was inserted until the tip of the needle reached the aortic valve. The heart was secured on the needle with a small brass clip and was immediately perfused with Perfusion buffer at a flow rate of 4.0 ml/min. The aorta was tied to the needle with silk thread. The time from the heart dissection until the start of perfusion should not exceed 1 min.

The heart was perfused with perfusion buffer for 2 min, or until the outflow from apex was cleared from blood. Then the perfusion with digestion buffer consisting of 29 ml perfusion buffer and 1.0 ml of 10 mg/ml Liberase Blendzyme 4 (Roche Diagnostics Corp. Indianapolis, IN) was continued for 7–12 min. At the end of the perfusion, the tissue became soft, swollen and light pink.

After the perfusion, the heart was cut from the needle just below the atria using sterile fine scissors and is placed in a Petri dish with 10 ml room temperature Incubation buffer (135 mM NaCl, 4 mM KCl, 1mM MgCl₂, 10 mM HEPES, 0.33 mM NaH₂PO₄, 10 mM glucose, 10mM BDM, pH 7.4). The heart was cut in half and the tissue was gently teased into small pieces with fine forceps. The obtained suspension of cardiomyocytes was gently pipetted up and down with a plastic pipette (2 mm tip) several times. Then the cells were transferred to a 15 ml sterile polypropylene conical tube and 10, 20, 30, 30 and 30 µl of a 100 mM CaCl₂ solution is added at 5 min intervals. The final content of calcium was 1.2 mM. Isolated myocytes were maintained at room temperature in this buffer.

Cell shortening/re-lengthening

Mechanical properties of the ventricular myocytes were determined using a video-based edge-detection system (IonOptix, Milton, MA), as described elsewhere (26). The myocytes were field stimulate data frequency of 1 and 4 Hz using a pair of platinum wires placed on the opposite sides of the dish chamber and connected to a MyoPacer Field Stimulator (IonOptix). Cardiomyocyte contractility was controlled by electrical stimulation. Mechanical properties of isolated ventricular myocytes were assessed by video-based edge detection. An inverted microscope, a low light-level video camera and a computer-based motion analyzer are used to track the movement of cell edges.

The isolated myocytes were diluted approximately ten fold with contractility buffer (135 mM NaCl, 4 mM KCl, 1mM MgCl₂, 10 mM HEPES, 0.33 mM NaH₂PO₄, 10 mM glucose, 10 mM BDM, 1.2 mM CaCl₂) and placed on a Teflon glass coverslip dish mounted on the stage of an inverted microscope (Olympus, IX-70). The cells were field stimulated to contract by the MyoPacer field stimulator through a pair of platinum electrodes at a

frequency of 1.0 Hz, pulse duration of 4ms and amplitude of 10volts. The image of the myocyte is obtained with an Ion Optix Myo Cam camera side-mounted onto the microscope and displayed on a computer monitor using the Soft-Edge software. Typically 10–15 individual myocytes are recorded before replacing the myocytes with a fresh dilution of unstimulated myocytes. This prevents exhaustion of the myocytes due to prolonged stimulation. Fifteen to twenty myocytes were analyzed for each heart. The cells being studied were scanned every 8.3ms so that the amplitude and velocity of shortening and lengthening can be recorded with good fidelity. The displacements of cell edges at both ends of the myocyte were detected and converted to an analog voltage signal, which is then digitized and stored for offline analysis. Steady-state twitches (8–10) were analyzed for cell length changes using the Soft-Edge software and averaged for each myocyte. Cell shortening and re-lengthening were assessed by percentage of peak shortening (%PS), time to 90% peak shortening (TPS₉₀), time to 90% re-lengthening (TR₉₀), and maximal velocities of shortening and re-lengthening ($\pm dL/dt$).

Intracellular fluorescence measurement of calcium

Intracellular calcium was determined using a dual-excitation fluorescence photomultiplier system (IonOptix), as described elsewhere (26). A separate cohort of myocytes was loaded with calcium-specific radio-metric fura 2-AM dye (1 μ mol/l) for 30 min in the dark, and the fluorescence was recorded. Myocytes were placed in a dish chamber on the stage on an Olympus IX-70 inverted microscope and imaged through a Fluor x100 objective. Cells were exposed to light emitted by a 75-Watt lamp and were passed through either 360- or 380-nm filters (bandwidths will be placed at ± 15 nm) while being stimulated to contract at 1.0 and 4.0Hz. Fluorescence emission was detected between 480 and 520nm by a photomultiplier tube after exciting the cells at 360nm for 0.5s and then at 380nm for the duration of their cording protocol. A 360nm scan was repeated at end of the experiments, and the qualitative changes in intracellular calcium concentration were interpreted from the ratio. The calcium transients were measured as changes in fura fluorescence intensity (FFI). FFI was determined as the difference between the levels of calcium in systolic and diastolic settings (FFI = peak FFI – baseline FFI). The time course of calcium fluorescence signal decay (the duration at which calcium transient decays 67% from the peak level) was calculated to determine intracellular calcium clearing rate. The myocytes included in the cell shortening/re-lengthening and calcium measurement studies met the following criterion: 1) rod-shaped with clear striation pattern; 2) quiescent in the absence of electrical stimulation; 3) stable mechanical behavior at 0.25 or 2 Hz and 37°C for 15 min; and 4) absence of sarcolemmal blebs.

In-Gel-Gelatin Zymography

mtMMP-9 activity of mitochondrial fraction was measured using gel-gelatin zymography. Briefly, isolated myocyte, washed two times in PBS, and incubated in the protein extraction buffer (in mMol/L: 50 mM Tris pH7.4, 5 mM EDTA, 150 mM NaCl, and 1mM phenylmethylsulfonyl fluoride, 0.1% Triton X-100), for 1h on ice. The homogenate was centrifuged for 600g for 5 minutes. Pellet 1 that consists of nuclei and whole cells was discarded. The supernatant was centrifuged again at 14,000g for 45 minutes to acquire mitochondrial fraction. Protein concentration was assayed using the Bradford method. Samples were electrophoretically resolved by 10% SDS-PAGE containing 2 mg/ml gelatin as a substrate. At the end, gel was incubated in renaturation buffer (2.5% TritonX-100) for 30 min to remove SDS, rinsed in distilled water, and then incubated overnight at 37°C in water bath in activation buffer (50 mM Tris-HCl, pH 7.4, and 5 mM CaCl₂). Gel was stained using 0.5% Coomassie blue R-250 for 1h. MMP activity was detected as a white band on a dark blue background and quantity densitometrically using Un-Scan software (Silk Scientific, Orem, UT).

Statistics

The contractility measurements were performed on 12–15 myocytes from 6–8 hearts in each group. Values were mean $\pm$ SEM. Significance was carried out by paired Student's *t*-test. $P < 0.05$ was considered significant.

Results

The results suggested that mitochondrial MMP-9 was activated in the myocyte mitochondria with increased contraction frequency (Figure 1). It was demonstrated that mtMMP-9 was significantly elevated 25% at 1 Hz stimulation relative to 0 Hz ($P < 0.05$, $n = 3$) and elevated 40% at 4 Hz stimulation relative to 0 Hz ($P < 0.01$, $n = 3$).

There was a decrease in cell lengthening at higher contraction frequency that resulted in the following length shortening relative to control, i.e. 0.80 decrease in length shortening relative to 1 Hz ($p < 0.05$, $n = 3$) [Figure 2A–F].

The levels of calcium transient were reduced with increase in contraction frequency with the following levels, i.e. 0.80 decrease in calcium transient relative to 1 Hz ($P < 0.05$) [Figure 3A–E].

Although number of frequency was higher, the magnitude of myocyte lengthening was shorter in 4 Hz stimulation than 1 Hz (Figure 4).

Discussion

Calcium is a primary regulator of contraction within the heart; transients are important for both sinoatrial depolarization and myocardial contraction (11). An inward calcium transient is responsible for the depolarization (Phase 0) of sinoatrial node as well as the plateau phase (Phase 2) of myocardial depolarization and contraction (25). Moreover, the contraction of cardiomyocytes requires calcium binding of TnC to allow tropomyosin to expose myosin binding sites on the globular actin myofilament (11). Calcium deregulation and mishandling is known to be involved in several pathologies and arrhythmias. Calcium enters from voltage dependent L-type calcium channels (I_{Ca} current); these results in Calcium-induced calcium release (CICR) by activating calcium channels on sarcoplasmic reticulum (SR) called Ryr receptors. These Ryr receptors allow even greater release of calcium from SR that allows for contraction to occur. Calcium exits cardiomyocytes via Ca/Na exchanger that requires the Na⁺/K⁺ pump to establish a gradient of extracellular sodium; moreover, intracellular calcium is mainly sequestered into SR via SR ATPase protein, SERCA.

Under physiological electrical stimulation, more calcium enters the cell from a greater number of action potentials (11). Moreover, the interval for calcium uptake is decreased; therefore, more calcium remains in the cell with subsequent contractions. A shorter interval may prevent NCX from extruding calcium quick enough; hence, intracellular calcium may accumulate to greater levels and increase force of contraction as more calcium binds to more TnC and unveils greater number of actin binding sites. SERCA uptake and release of calcium proceed at a greater rate to accommodate the higher number of action potential. Electrical stimulation of astroglia and fibroblast cells also resulted in induction of heat shock protein (hsp70) gene whose role is to maintain protein integrity and aid in protein folding (20).

Our lab has worked largely with homocysteine, a sulfur-containing amino acid that has been shown to be an independent risk factor for heart disease (18;22). Hcy-induced calcium-dependent PKC can phosphorylate ERK 1/2 to activate intracellular MMP-9 that proceeds

with vascular remodeling in vitro (13). Hence, calcium is an important inducer of MMP activity. It is known that calpain-1 disrupts inner mitochondrial membrane electronic gradient that results in an accumulation of reactive oxygen species within mitochondria (14). These ROS can activate mitochondrial MMP as well as diffusing from mitochondria to cytosol to activate intracellular MMP (14). Activation of MMP-9 via ROS can occur either directly or via ERK 1/2 pathways that have been shown to activate MMP-9 (14).

Our lab investigated Hcy-induced activation of mtMMP via Hcy agonizing NMDA-R receptor and increasing calcium within the cell; we also investigated its effect on mitochondria via ultra structural analysis and mitochondrial swelling assay; results showed that there was an enlargement of mitochondria as well as fragmentation of cristae that resulted in contractile dysfunction (12;15). This is logical considering how disruption of inner mitochondrial membrane would destroy the electric potential necessary to generate ATP. Contractile dysfunction was measured via the following: decrease in percent cell shortening, maximal rate of contraction and maximal rate of relaxation. Moreover, decrease of calcium transient amplitude was faster in wild type compared with HHcy. This was confirmed using pharmacological blocker of NMDA-R i.e MK-801, in presence of Hcy; contractile function returned (12;23). In fact, the contractile dysfunction in previous experiments resembles the data that we have collected upon stimulation of myocytes. This is strong evidence with the data that we have collected that calcium mishandling within cell can lead to mtMMP activation in the mitochondria. This is associated with an increase in contraction frequency and decrease in myocyte contractility

This can have serious implications in arrhythmogenesis. Abnormal electric pacing via ectopic electrical stimulation within the heart could possibly lead to fibrotic remodeling and long-term contractile dysfunction. Moreover, electrical stimulation or pacing as a therapeutic measure should be evaluated closely in higher pacing ranges.

In rat choroidallantois membranes, calcium was shown to be necessary for the development of a complex of MMPs that proceed with connective tissue degradation (25). Remodeling induced by electronic pacing has been demonstrated in atria of dogs; rapid pacing produced the following results: systolic, diastolic dysfunction, increased chamber stiffness, atrial enlargement, increased activity of intracellular MMP-9 by 50% with a concomitant decrease in TIMP-4 by 50%, and an unchanged number of calcium pumps (10). One interesting experiment found that there was an increase in MMP-2 expression in denervated muscle, and that electrical stimulation (ES) produced an increasing trend of MMP-2 expression compared to denervated muscle alone (20). PCR analysis resulted in an increased MMP-2 expression that was only significantly higher in the denervated muscle +ES group compared to control (20). The role of MMP-2 in denervated or electrically stimulated muscle is not well understood as of yet whether it is detrimental or beneficial.

It is known that there is a deposit and proliferation of connective tissue after muscle degeneration that leads to the impairment of mechanical and electrical properties (20). There has been some suggestion that it may be involved in allowing new nerve sprouting; however, others suggested that it may be related to connective tissue proliferation that would hinder mechanical function. The latter is more likely considering connective tissue density in denervated muscle in this experiment was much greater versus normal muscle. Moreover, increased expression of MMP-2 and MMP-9 are well-established to be involved in inflammatory connective tissue disease (7). It was shown that MMP-9 activation in synovial fluid from rheumatoid arthritis patients when extracted and incubated was delayed in absence of calcium. Hence, calcium is an important regulator. In cancer cells, an increase in extracellular calcium resulted in a dose-dependent of MMP2 expression. In fact, it was just recently discovered that MMP-7 deletion improved survival and myocardial conduction

pattern after a myocardial infarction; moreover, it was shown that connexin-43 is a substrate for MMP-7 (8). Finally, a study showed that MMP-9 deletion reduced neutrophil proliferation after ischemia/reperfusion injury in vivo.

The decrease in myocyte lengthening upon stimulation is due to inability of the cardiac muscle to fully relax at higher contraction rates (4Hz). Lesser calcium is released from SR before being sequestered and released again; we suggest based on previous studies that MMP can inhibit the SERCA pump thereby preventing normal release of calcium. This could be a potential negative feedback mechanism since MMP-9 activation requires calcium increase, and MMP-9 activation can use SERCA as a substrate to increase calcium within cell. This is consistent with lesser calcium transient with greater electrical stimulation rate (4Hz).

Research highlights

- MMP-9 was activated in the myocyte mitochondria with increased contraction frequency
- Myocyte cell lengthening was decrease at higher contraction frequency
- Calcium transient were reduced with increase in contraction frequency

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

References

1. Ahtikoski AM, Tuominen H, Korpelainen JT, Takala TE, Oikarinen A. Collagen synthesis and degradation in polyneuropathy and myopathies. *Muscle Nerve*. 2004; 30:602–608. [PubMed: 15389647]
2. Cornelussen RN, Splett V, Klepfer RN, Stegemann B, Kornet L, Prinzen FW. Electrical modalities beyond pacing for the treatment of heart failure. *Heart Fail Rev*. 2010
3. Gao X, Xu X, Pang J, Zhang C, Ding JM, Peng X, Liu Y, Cao JM. NMDA receptor activation induces mitochondrial dysfunction, oxidative stress and apoptosis in cultured neonatal rat cardiomyocytes. *Physiol Res*. 2007; 56:559–569. [PubMed: 16925458]
4. Gill SE, Parks WC. Metalloproteinases and their inhibitors: regulators of wound healing. *Int J Biochem Cell Biol*. 2008; 40:1334–1347. [PubMed: 18083622]
5. Herrmann M, Taban-Shomal O, Hubner U, Bohm M, Herrmann W. A review of homocysteine and heart failure. *Eur J Heart Fail*. 2006; 8:571–576. [PubMed: 16504575]
6. Hoit BD, Takeishi Y, Cox MJ, Gabel M, Kirkpatrick D, Walsh RA, Tyagi SC. Remodeling of the left atrium in pacing-induced atrial cardiomyopathy. *Mol Cell Biochem*. 2002; 238:145–150. [PubMed: 12349902]
7. Lichtinghagen R, Seifert T, Kracke A, Marckmann S, Wurster U, Heidenreich F. Expression of matrix metalloproteinase-9 and its inhibitors in mononuclear blood cells of patients with multiple sclerosis. *J Neuroimmunol*. 1999; 99:19–26. [PubMed: 10496173]
8. Lindsey ML, Escobar GP, Mukherjee R, Goshorn DK, Sheats NJ, Bruce JA, Mains IM, Hendrick JK, Hewett KW, Gourdie RG, Matrisian LM, Spinale FG. Matrix metalloproteinase-7 affects connexin-43 levels, electrical conduction, and survival after myocardial infarction. *Circulation*. 2006; 113:2919–2928. [PubMed: 16769909]
9. Ma YS, Chen YC, Lu CY, Liu CY, Wei YH. Upregulation of matrix metalloproteinase 1 and disruption of mitochondrial network in skin fibroblasts of patients with MERRF syndrome. *Ann N Y Acad Sci*. 2005; 1042:55–63. [PubMed: 15965045]
10. Maier LS, Bers DM. Calcium, calmodulin, and calcium-calmodulin kinase II: heartbeat to heartbeat and beyond. *J Mol Cell Cardiol*. 2002; 34:919–939. [PubMed: 12234763]

11. Makowski GS, Ramsby ML. Autoactivation profiles of calcium-dependent matrix metalloproteinase-2 and -9 in inflammatory synovial fluid: effect of pyrophosphate and bisphosphonates. *Clin Chim Acta*. 2005; 358:182–191. [PubMed: 15921672]
12. Moshal KS, Kumar M, Tyagi N, Mishra PK, Metreveli N, Rodriguez WE, Tyagi SC. Restoration of contractility in hyperhomocysteinemia by cardiac-specific deletion of NMDA-R1. *Am J Physiol Heart Circ Physiol*. 2009; 296:H887–H892. [PubMed: 19181966]
13. Moshal KS, Sen U, Tyagi N, Henderson B, Steed M, Ovechkin AV, Tyagi SC. Regulation of homocysteine-induced MMP-9 by ERK1/2 pathway. *Am J Physiol Cell Physiol*. 2006; 290:C883–C891. [PubMed: 16251475]
14. Moshal KS, Singh M, Sen U, Rosenberger DS, Henderson B, Tyagi N, Zhang H, Tyagi SC. Homocysteine-mediated activation and mitochondrial translocation of calpain regulates MMP-9 in MVEC. *Am J Physiol Heart Circ Physiol*. 2006; 291:H2825–H2835. [PubMed: 16877562]
15. Moshal KS, Tipparaju SM, Vacek TP, Kumar M, Singh M, Frank IE, Patibandla PK, Tyagi N, Rai J, Metreveli N, Rodriguez WE, Tseng MT, Tyagi SC. Mitochondrial matrix metalloproteinase activation decreases myocyte contractility in hyperhomocysteinemia. *Am J Physiol Heart Circ Physiol*. 2008; 295:H890–H897. [PubMed: 18567713]
16. Moshal KS, Tipparaju SM, Vacek TP, Kumar M, Singh M, Frank IE, Patibandla PK, Tyagi N, Rai J, Metreveli N, Rodriguez WE, Tseng MT, Tyagi SC. Mitochondrial matrix metalloproteinase activation decreases myocyte contractility in hyperhomocysteinemia. *Am J Physiol Heart Circ Physiol*. 2008; 295:H890–H897. [PubMed: 18567713]
17. Nelson KK, Melendez JA. Mitochondrial redox control of matrix metalloproteinases. *Free Radic Biol Med*. 2004; 37:768–784. [PubMed: 15304253]
18. Ni M, Zhang XH, Jiang SL, Zhang Y. Homocysteinemia as an independent risk factor in the Chinese population at a high risk of coronary artery disease. *Am J Cardiol*. 2007; 100:455–458. [PubMed: 17659928]
19. Rucklidge GJ, Milne G, McGaw BA, Milne E, Robins SP. Turnover rates of different collagen types measured by isotope ratio mass spectrometry. *Biochim Biophys Acta*. 1992; 1156:57–61. [PubMed: 1472539]
20. Russo TL, Peviani SM, Durigan JL, Salvini TF. Electrical stimulation increases matrix metalloproteinase-2 gene expression but does not change its activity in denervated rat muscle. *Muscle Nerve*. 2008; 37:593–600. [PubMed: 18293392]
21. Sood HS, Cox MJ, Tyagi SC. Generation of nitrotyrosine precedes activation of metalloproteinase in myocardium of hyperhomocysteinemic rats. *Antioxid Redox Signal*. 2002; 4:799–804. [PubMed: 12470508]
22. Tyagi N, Mishra PK, Tyagi SC. Homocysteine, hydrogen sulfide (H₂S) and NMDA-receptor in heart failure. *Indian J Biochem Biophys*. 2009; 46:441–446. [PubMed: 20361707]
23. Tyagi N, Vacek JC, Givvimani S, Sen U, Tyagi SC. Cardiac specific deletion of N-methyl-D-aspartate receptor 1 ameliorates mtMMP-9 mediated autophagy/mitophagy in hyperhomocysteinemia. *J Recept Signal Transduct Res*. 2010; 30:78–87. [PubMed: 20170426]
24. Tyagi SC, Meyer L, Schmaltz RA, Reddy HK, Voelker DJ. Proteinases and restenosis in the human coronary artery: extracellular matrix production exceeds the expression of proteolytic activity. *Atherosclerosis*. 1995; 116:43–57. [PubMed: 7488332]
25. Verkerk AO, Wilders R, Zegers JG, van Borren MM, Ravestloot JH, Verheijck EE. Ca(2+)-activated Cl(−) current in rabbit sinoatrial node cells. *J Physiol*. 2002; 540:105–117. [PubMed: 11927673]
26. Ye G, Metreveli NS, Donthi RV, Xia S, Xu M, Carlson EC, Epstein PN. Catalase protects cardiomyocyte function in models of type 1 and type 2 diabetes. *Diabetes*. 2004; 53:1336–1343. [PubMed: 15111504]

Figure 1.

Zymographic analysis of myocyte mitochondrial (mt) MMP-9 activity. Myocytes were stimulated at 0, 1, and 4 hertz (Hz); mitochondria were isolated using differential centrifugation and analyzed on 1% gelatin-gel zymography (gel panel). The bar graphs represent the accumulative densitometric scan data from n=3 in each group. *, p<0.05 compared to 0 Hz.

Figure 2. Single myocyte lengthening stimulated at 1Hz versus 4Hz. (A) lengthening at 1 Hz; (B) at 4 Hz; (C) maximum rate of relaxation; (D) maximum rate of contraction; (E) cell shortening; (F) time to release calcium. Each bar represents mean $\pm$ SEM from n=3. *, p<0.05 compared to 1 Hz.

Figure 3.

Single myocyte calcium (340nm/380nm) transient was analyzed at 1 Hz (A) versus 4 Hz (B). (C) Cell shortening (%); (D) rate of calcium release; (E) time to release calcium. Each bar represents mean $\pm$ SEM from n=3. p<0.05 compared to 1 Hz.

Figure 4.
Representative tracing of myocyte lengthening at 1 and 4 Hz.