CORRESPONDENCE

Vision Screening in Preschool Children: Do the Data Support Universal Screening?

by Prof. Dr. med. Wolf A. Lagrèze in volume 28–29/2010

Children With Problems at School

The Professional Association of German Ophthalmologists (Berufsverband der Augenärzte Deutschlands, BVA) and the Ophthalmologische Gesellschaft (DOG, the medical scientific association of ophthalmology in Germany) protested against IQWiG's rejection of vision screening in preschool children, in some cases insulting and blaming the institute's staff. As Professor Lagrèze showed in his article, this was unjustified. Vision screening was intended to identify in particular amblyopia with strabismus, in order to prevent bilateral sight impairment in case additional eye disorders were to develop in old age.

However, this should be completed if possible by the 3rd year of life because the chance of a cure notably decreases after that age. Transferring preventive eye investigations in early childhood from pediatricians to ophthalmologists would make sense. Screening measures at any later stage therefore hardly make any sense.

In my long years' experience with many thousands of affected children, something that really puts a burden on children after the 6th year of life, when they have entered school, are latent binocular vision impairments, which affect children's ability to perform and thus endanger their academic and later professional careers.

Severe asthenopic symptoms with headache/migraine, impaired gross and fine motor skills, difficulties in reading/spelling, and AD(H)D will then lead to severe performance impairments.

Children with problems at school should therefore be investigated for associated heterophorias (latent strabismus in which one eye tends to deviate either medially or laterally) with the Pola-test according to Haase's particularly effective measurement and correction method, in order to prevent maldevelopments.

Ophthalmologists, pediatricians, psychologists, and optometrists, as well as educators have joined forces to be contacted under the umbrella of the International Association for Binocular Fullcorrection (IVBV).

DOI: 10.3238/arztebl.2011.0039a

REFERENCES

1. Lagrèze WA: Vision screening in preschool children: Do the data support universal screening? Dtsch Arztebl Int 2010; 107(28–29): 495–9.

Dr. med. Fritz Gorzny
Düppelstr. 47
56179 Vallendar, Germany
dr.fritz_gorzny@web.de

Early Start of Screening

Lagrèze says that studies up to now did not sufficiently prove the benefits of preschool screening. We agree if only one screening is done during or after the maturation of the visual system. However, therapeutic success depends on the age of the child at the onset of treatment. The available studies were published before the age of the internet and therefore unfortunately hardly find any attention today.

Urgently our aim is not therapy but rather prevention on the basis of known risk factors. These are age dependent and well defined for the second half of the first year of life, as shown by British (1) and Swedish (2) studies.

In a pilot project, "prevention of amblyopia and strabismus in infants and toddlers," we are trying to implement this early detection system. Since 2001, in the obstetric clinic of Kulmbach Medical Center we have measured refraction by skiascopy in neonates under cycloplegia with tropicamide and recommend further investigations on the basis of our findings. This early start of screening is necessary because of the rapid development of the visual system within the first 12 months of life.

In two cross sectional studies in school starters in 2005 (without screening) and in 2008 (with screening) we observed a reduced rate of amblyopia from 6.2% (2005) to 2.6% (2008). In the first longitudinal study (3) we found 5 children with strabismus among 27 children with >3.5 diopters of hyperopia that was not corrected by the 3rd year of life. In a group of 22 children who had started wearing spectacles during the first year of life no child developed strabismus. According to Atkinson et al. these results suggest that it is necessary to correct refraction errors early in the first year of life if we want to prevent amblyopia and strabismus.

DOI: 10.3238/arztebl.2011.0039b

REFERENCES

1. Atkinson J, Braddick O, Robier B, et al.: Eye 1996; 10: 189–98.
2. Kvarnström G, Jakobsson P, Lennerstrand G: Visual screening of Swedish children: An ophthalmological evaluation. Acta ophthalmol Scand 2001; 79: 240–4.
3. Petzold G et al.: Augenärztliche Prävention im 1. Lebensjahr. Monatsschr Kinderheilk 2007; 155: 61–7.
4. Lagrèze WA: Vision screening in preschool children: Do the data support universal screening? Dtsch Arztebl Int 2010; 107(28–29): 495–9.

Prof. Dr. med. Wolfgang Haase
Dr. med. Helmut Bock

Dr. med. Gernot Petzold
für die Arbeitsgruppe
Hans Hacker Str. 1
95326 Kulmbach, Germany
gernot.petzold@t-online.de

In Reply:

Gorzny correctly says that uncorrected hyperopia ($\geq +3.5$ dpt) increases the risk for amblyopia. However, only vague indications support the idea that such hyperopia results in general symptoms such as headaches, writing/spelling problems, and AD(H)D. Improvements of such symptoms are often reported once a child starts wearing spectacles, but these may be due to a placebo effect, chance, or regression to the mean. This criticism also applies to the mentioned “associated heterophoria,” which fans of the measuring and correction method according to Hans-Joachim Haase (MKH) treat with prism spectacles.

In Haase’s method, a very slight misalignment of one eye from the fixed object (fixation disparity) is seen as indicative for the affected person’s problems to overcome larger misalignments completely.

However, the premise that such a slight misalignment could be detected on the basis of the perception of certain test figures has not been confirmed by objective, high-resolution measurements of eye alignment (1).

The cross sectional studies by Petzold, Bock, and Haase have confirmed the findings of earlier studies (for example, Atkinson et al, 1996), which found a reduction in the prevalence of amblyopia as a result of screening. The mention of the very early timing of the first refraction measurement is valuable and interesting. What is surprising, however, is the high prevalence of amblyopia in the non-screened cohort.

It is to be hoped that the results of the comparison of these two cross sectional studies in children with or without screening will be published soon. With regard to the strength of evidence of a clinical study, we’d like to point out that in these days of evidence based medicine the requirements for study designs have clearly changed and older studies should therefore be interpreted with caution. Kvarnström et al (2), for example, wrote in the abstract of their much cited study that screening reduced the prevalence of amblyopia in

Sweden from 2.0% to 0.2%. However, these values are not included in the actual publication itself, and neither was an unscreened control group. Instead the authors cite their own work of 1998, in which they cite a Danish study (3) that investigated adults older than 60 for macular degeneration and also amblyopia. These adults had not been examined in childhood. This underlines the fact that only few studies meet the required quality criteria and lesser, but still available, evidence is at risk of being ignored—sometimes rightly so, sometimes wrongly so. Ultimately, the resource allocating institutions (GBA [Gemeinsamer Bundesausschuss — Federal Joint Committee]/IQWiG/statutory health insurers) will prioritize a screening program only if existing indications of positive effects and benefits of childhood screening for the general public are confirmed by randomized controlled studies.

DOI: 10.3238/arztebl.2011.0040

REFERENCES

1. Gerling J, de Paz H, Schroth V, Bach M, Kommerell G: Ist die Feststellung einer Fixationsdisparation mit der Mess- und Korrektionsmethode nach H.-J. Haase (MKH) verlässlich? *Klin Monatsbl Augenheilkd* 2000; 216: 401–11.
2. Kvarnström G, Jakobsson P, Lennerstrand G: Screening for visual and ocular disorders in children, evaluation of the system in Sweden. *Acta Paediatr* 1998; 87: 1173–9.
3. Vinding T, Gregersen E, Jensen A, Rindziunski E: Prevalence of amblyopia in old people without previous screening and treatment. *Acta Ophthalmol* 1991; 69: 796–8.
4. Lagrèze WA: Vision screening in preschool children: Do the data support universal screening? *Dtsch Arztebl Int* 2010; 107(28–29): 495–9.

Prof. Dr. med. Wolf A. Lagrèze

Universitäts-Augenklinik
Killianstr. 5
79106 Freiburg, Germany
mail@lagreze.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.