Published in final edited form as:

Nat Genet. 2009 February ; 41(2): 160–162. doi:10.1038/ng.292.

15q13.3 microdeletions increase risk of idiopathic generalized epilepsy

Ingo Helbig¹, Heather C Mefford², Andrew J Sharp³, Michel Guipponi³, Marco Fichera⁴, Andre Franke¹, Hiltrud Muhle¹, Carolien de Kovel¹, Carl Baker², Sarah von Spiczak¹, Katherine L Kron¹, Ines Steinich¹, Ailing A Kleefuß-Lie¹, Costin Leu¹, Verena Gaus¹, Bettina Schmitz¹, Karl M Klein¹, Philipp S Reif¹, Felix Rosenow¹, Yvonne Weber¹, Holger Lerche¹, Fritz Zimprich¹, Lydia Urak¹, Karoline Fuchs¹, Martha Feucht¹, Pierre Genton⁵, Pierre Thomas⁶, Frank Visscher¹, Gerrit-Jan de Haan¹, Rikke S Møller¹, Helle Hjalgrim¹, Daniela Luciano⁴, Michael Wittig¹, Michael Nothnagel¹, Christian E Elger¹, Peter Nürnberg¹, Corrado Romano⁴, Alain Malafosse³, Bobby P C Koeleman¹, Dick Lindhout¹, Ulrich Stephani¹, Stefan Schreiber¹, Evan E Eichler^{2,7}, and Thomas Sander¹

¹ EPICURE Consortium; a list of participating centers is provided at the end of the paper ² Department of Genome Sciences, University of Washington, Seattle, Washington 98195, USA ³ Department of Genetic Medicine and Development, University of Geneva Medical School and University Hospitals of Geneva, 1211 Geneva, Switzerland ⁴ IRCCS Associazione Oasi Maria Santissima, 94018 Troina, Italy ⁵ Centre Saint Paul-Hôpital Henri Gastaut, 13258 Marseilles 09, France ⁶ Unité Fonctionnelle EEG-Epileptologie, Service de Neurologie, Hôpital Pasteur, 06002 Nice Cedex 01, France ⁷ Howard Hughes Medical Institute, University of Washington, Seattle, Washington 98195, USA

Abstract

We identified 15q13.3 microdeletions encompassing the *CHRNA7* gene in 12 of 1,223 individuals with idiopathic generalized epilepsy (IGE), which were not detected in 3,699 controls (joint $P = 5.32 \times 10^{-8}$). Most deletion carriers showed common IGE syndromes without other features previously associated with 15q13.3 microdeletions, such as intellectual disability, autism or schizophrenia. Our results indicate that 15q13.3 microdeletions constitute the most prevalent risk factor for common epilepsies identified to date.

Idiopathic generalized epilepsies (IGE) are common seizure disorders accounting for up to one-third of all epilepsies¹. The vast majority of individuals with IGE have a complex genetic etiology², for which the underlying genetic alterations remain largely unknown. Recently, a 15q13.3 microdeletion syndrome has been identified in 0.2–0.3% of individuals

Correspondence should be addressed to T.S. (sandert@uni-koeln.de).

Note: Supplementary information is available on the Nature Genetics website.

AUTHOR CONTRIBUTIONS

T.S. and E.E.E. initiated and designed the study; I.H., H.M., S.v.S., I.S., A.A.K.-L., V.G., B.S., K.M.K., P.S.R., F.R., Y.W., H.L., F.Z., L.U., K.F., M. Feucht, F.V., G.-J.d.H., R.S.M., H.H., D. Luciano, C.R., D. Lindhout, C.E.E., U.S. and T.S. recruited and phenotyped the EPICURE sample; H.C.M., A.J.S., M.G., M. Fichera, C.B., P.G., P.T., A.M. and E.E.E. recruited and phenotyped the mixed IGE sample; A.F., M.W., M.N. and S.S. recruited and phenotyped the PopGen control sample; I.H., A.F., C.L., K.L.K., I.S., M.W., M.N., P.N. and T.S. performed the CNV analysis on SNP arrays; H.C.M., A.J.S., M. Fichera, C.B. and D. Luciano performed the qPCR screening; H.C.M., M. Fichera, C.B. and D. Luciano performed the screening using Illumina Genotyping BeadChips; H.C.M., A.J.S. and C.B. performed the confirmation using NimbleGen arrays; C.d.K., B.P.C.K. and D. Lindhout performed the confirmation using Illumina CNV BeadChips; I.H., H.C.M., A.J.S., M.G., M. Fichera, A.F., C.d.K., K.L.K., C.R., B.P.C.K., D. Lindhout, E.E.E. and T.S. coordinated the work and prepared the manuscript.

with mental retardation and epilepsy³, schizophrenia^{4,5}, autism and other neuropsychiatric features⁶. The critical region of the 1.5-Mb deletion on 15q13.3 contains at least seven genes, including the *CHRNA7* gene coding for the $\alpha 7$ subunit of the nicotinic acetylcholine receptor, which is considered a plausible candidate gene for the epilepsy phenotype.

Susceptibility loci for common idiopathic epilepsies, comprising benign epilepsy of childhood with centrotemporal spikes⁷ and common IGE syndromes^{8,9}, have also been mapped to the 15q13–q14 region. To test whether the 15q13.3 deletion increases risk of common epilepsies, we screened for structural variants within the 15q13.3 region in two independent samples of individuals with IGE and ancestrally matched controls. The first sample comprised 647 unrelated IGE cases of Western European ancestry (EPICURE sample) and 1,202 German controls (PopGen) genotyped using the Affymetrix Genome-Wide Human SNP array 6.0. We identified the 15q13.3 microdeletion in 7 of 647 IGE cases (Supplementary Fig. 1 online) with different IGE syndromes (Table 1). All but one of these showed segmental breakpoints BP4 and BP5, as observed in the 15q13.3 microdeletion syndrome⁴. The other showed a 3.8-Mb deletion defined by BP3 and BP5. The 15q13.3 deletion was not detected in any of 1,202 controls examined.

We next examined the frequency of 15q13.3 microdeletions in a second independent sample of 576 IGE cases from Switzerland ($n = 205$), North America ($n = 133$) and Northern Europe ($n = 238$) as well as 2,497 controls from North America of predominantly European ancestry genotyped with various methods (Supplementary Methods online). In this sample, we found 15q13.3 microdeletions in 5 of 576 cases and 0 of 2,497 controls. Altogether, we identified 15q13.3 deletions in 12 of 1,223 IGE cases and in 0 of 3,699 controls ($P = 5.32 \times 10^{-8}$, Fisher's exact test). We used custom array CGH (10 of 12 cases; Fig. 1) or SNP and CNV arrays (2 of 12 cases) to verify the deletions in all IGE probands and their available first-degree family members. We identified duplications involving *CHRNA7* in 12 of 1,223 cases and 23 of 3,699 controls ($P = 0.27$, Pearson's χ^2 test; Supplementary Methods and Supplementary Fig. 2 online). Thus, our results suggest that the 15q13.3 deletion only, and not the reciprocal duplication, represents a major risk factor for IGE.

In our study, parental DNA was available for 5 of 12 probands with the deletion (Supplementary Fig. 3 online). In one individual (L1371), the deletion was apparently de novo, as we found that both clinically unaffected parents did not carry the deletion using custom array CGH. For the other four individuals, we observed parental transmission from one father and three mothers. One mother (E562M) suffered from panic disorder, a phenotype previously associated with 15q13.3 deletions⁶. The other three transmitting parents were apparently clinically unaffected, although we cannot exclude that subclinical manifestations such as age-dependent paroxysmal EEG discharges or undetected mild and remitting IGE phenotypes might have been missed. In one family (E562), both siblings carried the deletion and were affected by IGE. In family 254, the proband's brother (254B) also carried the deletion and was affected by IGE. In the family of IGE proband 1674, the 15q13.3 microdeletion was present in a brother (2376) with severe intellectual disability but without a history of seizures.

The phenotypes of most of the individuals reported here with the 15q13.3 microdeletion differ notably from those reported originally for the 15q13.3 microdeletion syndrome, in which individuals showed marked intellectual disability, seizures, growth retardation and dysmorphic features³. Although we observed severe intellectual disability in 1 of 12 and mild intellectual disability in 2 of 12 probands, 9 of 12 of the probands in our sample had characteristic features of IGE without dysmorphic features or intellectual disability (Table 1). Seizure types in these individuals included typical absence seizures, myoclonic seizures

and primary generalized tonic-clonic seizures, all of them occurring at the typical age of onset. Electroencephalographic (EEG) records were available for all individuals and showed normal background activity with paroxysmal generalized spike-wave discharges, which represent the EEG hallmark of IGE.

Two studies have reported association of the 15q13.3 microdeletion with schizophrenia and related psychoses^{4,5}. Consistent with our study, those studies observed intellectual disability in only a small fraction of individuals carrying the deletion. In the current study, none of the individuals with IGE carrying a 15q13.3 deletion had a history of a psychotic episode. Thus, our findings extend the phenotypic spectrum related to the 15q13.3 deletion to common IGE syndromes without the previously reported neuropsychiatric features. Taken together, the current studies reveal extensive variability in the phenotypic manifestation associated with the 15q13.3 deletion, ranging from apparently healthy individuals to severely affected individuals with a broad spectrum of neuropsychiatric disorders^{3–6}. These findings imply that shared mechanisms are involved in the pathogenesis of a spectrum of seemingly unrelated neuropsychiatric disorders, and argue for a new framework for understanding complex genetic diseases.

The critical region affected by the BP4-BP5 deletions harbors at least seven genes (*ARHGAP11B*, *MTMR15*, *MTMR10*, *TRPM1*, *KLF13*, *OTUD7A* and *CHRNA7*) that might contribute to the seizure phenotype, including *CHRNA7* as the prime candidate gene. Compelling evidence suggests that the impairment of neuronal ion channel function may be pivotal to the pathology of IGE¹⁰. Cholinergic pathways have several important functions in the brain, and nicotinic acetylcholine receptors containing the $\alpha 7$ subunit are widely expressed throughout the central nervous system¹¹. These receptors are localized both pre- and postsynaptically and are thought to modulate excitatory and inhibitory pathways. In particular, *CHRNA7* is highly expressed in the reticular thalamus¹², indicating a role in modulating thalamo-cortical pathways, which are central to the generation of primarily generalized seizures seen in IGE¹³. *CHRNA2*, *CHRNA4* and *CHRNA2*, which code for the $\alpha 2$, $\alpha 4$ and $\beta 2$ subunits of the nicotinic acetylcholine receptor, respectively, have a causative role in autosomal dominant nocturnal frontal lobe epilepsy (ADNFLE)¹¹. Together, these lines of evidence strongly support an involvement of *CHRNA7* in epileptogenesis.

The genetic architecture of common seizure disorders is thought to show a biological continuum ranging from rare monogenic forms to common epilepsies with complex inheritance. Positional cloning of epilepsy genes has been successful in large families with monogenic inheritance, and multiple rare gene variants, including mutations in *CACNA1H* and *EFHC1*, have been found in a small proportion of individuals with IGE¹⁰. However, common susceptibility alleles have not been identified so far. Here we describe a structural variant that is virtually absent in the general population (<0.02%)^{4,5} and roughly 50 times more frequent in individuals with IGE in the present study (1%). Furthermore, the frequency of 15q13.3 deletions in IGE seems to be higher than that reported in intellectual disability or schizophrenia (Supplementary Table 1 online). Given the strong epileptogenic effect, the 15q13.3 microdeletion represents the most prevalent major risk factor for IGE identified to date.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank all the subjects and their families for participating in this study. We are also grateful to H. Bode for referral of subjects and A. Ackerhans and K. Moldenhauer for database management. This study used samples from the NINDS Human Genetics Resource Center DNA and Cell Line Repository (<http://ccr.coriell.org/ninds>), as well as clinical data. NINDS Repository sample numbers corresponding to the samples used are available upon request. This study was supported by grants from the German Research Foundation (SA434/4-1, T.S., P.N.), the German Federal Ministry of Education and Research (National Genome Research Network, NGFN-2: NeuroNet, NGFNplus: EMINet), the European Community (FP6 Integrated Project EPICURE, LSHM-CT-2006-037315; grant agreement 219250, A.J.S.), the PopGen biobank, the University of Kiel (I.H.), the Danish National Research Foundation (R.S.M.), in part by grants from the NIH (HD043569, E.E.E.), the National Epilepsy Funds (NEF, grant no. 04-08, B.P.C.K., D. Lindhout), the Netherlands Organization for Scientific Research (NOW, grant no. 917.66.315, B.P.C.K., C.d.K.) and the German Research Foundation/German Federal Ministry of Education and Research (DFG/BMBF) excellence cluster "Inflammation at Interfaces" (A.F., M.W., S.S.). E.E.E. is an investigator of the Howard Hughes Medical Institute.

References

1. Jallon P, Latour P. *Epilepsia* 2005;46(Suppl 9):10–14. [PubMed: 16302871]
2. Ottman R. *Epilepsia* 2005;46(Suppl 10):7–14. [PubMed: 16359464]
3. Sharp AJ, et al. *Nat Genet* 2008;40:322–328. [PubMed: 18278044]
4. International Schizophrenia Consortium. *Nature* 2008;455:237–241. [PubMed: 18668038]
5. Stefansson H, et al. *Nature* 2008;455:232–236. [PubMed: 18668039]
6. Miller DT, et al. *J Med Genet* advance online publication. September 19;2008 10.1136/jmg.2008.059907
7. Neubauer BA, et al. *Neurology* 1998;51:1608–1612. [PubMed: 9855510]
8. Elmslie FV, et al. *Hum Mol Genet* 1997;6:1329–1334. [PubMed: 9259280]
9. Sander T, et al. *Hum Mol Genet* 2000;9:1465–1472. [PubMed: 10888596]
10. Helbig I, Scheffer IE, Mulley JC, Berkovic SF. *Lancet Neurol* 2008;7:231–245. [PubMed: 18275925]
11. Steinlein OK, Bertrand D. *Biochem Pharmacol* 2008;76:1175–1183. [PubMed: 18691557]
12. Breese CR, et al. *J Comp Neurol* 1997;387:385–398. [PubMed: 9335422]
13. Blumenfeld H. *Epilepsia* 2005;46(Suppl 9):21–33. [PubMed: 16302873]

Participating centers in the EPICURE Integrated Project are as follows

Department of Neuropediatrics, University Medical Center Schleswig-Holstein (Kiel Campus), Schwanenweg 20, 24105 Kiel, Germany (I.H., H.M., S.v.S., K.L.K., I.S., U.S.). Institute for Clinical Molecular Biology, University Medical Center Schleswig-Holstein (Kiel Campus), Arnold-Heller-Strasse 3, 24105 Kiel, Germany (A.F., M.W., S.S.). Institute of Medical Informatics and Statistics, University Medical Center Schleswig-Holstein (Kiel Campus), Brunswiker Strasse 10, 24105 Kiel, Germany (M.N.). Cologne Center for Genomics, University of Cologne, Zùlpicher Strasse 47, 50674 Cologne, Germany (T.S., C.L., P.N.). Max-Delbrück-Center for Molecular Medicine, Robert-Rössle-Strasse 10, 13125 Berlin, Germany (C.L., V.G., T.S.). Department of Epileptology, University of Bonn, Sigmund Freud Strasse 1, 53105 Bonn, Germany (A.A.K.-L., C.E.E.). Department of Neurology, Charité University Medicine, Campus Virchow Clinic, Humboldt University of Berlin, Augustenburger Platz 1, 13353 Berlin, Germany (V.G., B.S., T.S.). Interdisciplinary Epilepsy-Center, Department of Neurology, Philipps University Marburg, 35033 Marburg, Germany (K.M.K., P.S.R., F.R.). Department of Neurology, University of Ulm, Helmholtzstrasse 8/1, 89081 Ulm, Germany (Y.W., H.L.). Department of Clinical Neurology (F.Z.) and Department of Pediatrics and Neonatology (L.U., M.F.), Medical University of Vienna, Währinger Gürtel 18–20, A-1090 Vienna, Austria. Center of Brain Research, Department of Biochemistry and Molecular Biology, Medical University of Vienna, Spitalgasse 4, A-1090 Vienna, Austria (K.F.). Department of Neurology, Danish

Epilepsy Centre, Dianalund Kolonivej 1, 4293 Dianalund, Denmark (R.S.M., H.H.).
Wilhelm Johannsen Centre for Functional Genome Research, University of Copenhagen,
Blegdamsvej 3, 2200 Copenhagen N, Denmark (R.S.M.). Netherlands Section Complex
Genetics, Department of Medical Genetics, University Medical Center Utrecht, Str. 2.112
Universiteitsweg 100, 3584 CG Utrecht, The Netherlands (B.P.C.K., C.d.K., D. Lindhout).
Department of Neurology, Oosterschelde Hospital, P.O. Box 106, 4460 BB Goes, The
Netherlands (F.V.). SEIN Epilepsy Institute in The Netherlands, P.O. Box 540, 2130AM
Hoofddorp, The Netherlands (G.-J.d.H., D. Lindhout).

Figure 1. Confirmation of 15q13.3 microdeletions using custom array CGH. High-resolution oligonucleotide array mapping of the 15q12–q13.3 region in 10 of 12 IGE probands with 15q13.3 microdeletions. Probes with $\log_2$ ratios above or below a threshold of 1.5 s.d. are colored green (duplications) or red (deletions). Hashed lines indicate the breakpoint regions BP3–BP5.

Table 1

Phenotypic features of individuals with 15q13.3 microdeletions

Individual	Descent	Diagnosis	Seizure types	Age of onset	EEG	Cognition
EPIPURE sample						
A067	German	JME	Myoclonus GTCS	16 y 19 y	GSW	Normal
1674	German	CAE	Absence	5 y	GSW	ID
254A	German	IAE	Absence GTCS	9 y 13 y	GSW	Normal
60A	German	IAE	Absence GTCS	12 y 14 y	GSW	Normal
EZ1194	Austrian	JME	Absence Myoclonus GTCS	6 y 10 y 20 y	GSW	Normal
40281601	German	CAE	Absence GTCS	Uncertain 3 y	GSW	Normal
D04u0213	Dutch	JME	Absence Myoclonus GTCS	4 y 13 y	Irreg. GSW PPR	Normal
Mixed IGE sample						
E421	Northern African	JME	Myoclonus GTCS	<26 y 26 y	Irreg. GSW	Normal
E435	French	JME	Myoclonus GTCS	12 y 12 y	GSW	Mild deficits
L1371	German	CAE	Absence	4 y	Irreg. GSW	Normal
D07u0771	Dutch	IAE	Absence	14 y	GSW	Normal
E562	French	JME	Myoclonus GTCS	12 y 13 y	GSW	Mild ID (IQ = 73)
Affected first-degree family members						
2376 (brother of 1674)	German		None			Severe ID
254B (sister of 254A)	German	JME	Absence Myoclonus	15 y 15 y	GSW	Normal
E562M (mother of E562)	French		None			Panic disorder
E562B (brother of E562)	French	EGTCS	GTCS	Uncertain	Unknown	Normal

CAE, childhood absence epilepsy; JAE, juvenile absence epilepsy; JME, juvenile myoclonic epilepsy; EGTCS, idiopathic epilepsy with GTCS; GTCS, generalized tonic-clonic seizures; GSW, generalized spike-wave discharges (2.5–5.0 Hz); Irreg. GSW, irregular generalized spike-wave discharges; PPR, photo-paroxysmal response; ID, intellectual disability.