Maternal investment of female mallards is influenced by male carotenoid-based coloration

M. Giraudeau^{1,*}, C. Duval^{2,†}, G. Á. Czirják^{2,3,‡}, V. Bretagnolle⁴,
C. Eraud⁵, K. J. McGraw¹ and P. Heeb²

¹*School of Life Sciences, Arizona State University, Tempe, AZ 85287-4501, USA*

²*Laboratoire Évolution et Diversité Biologique (EDB), UMR 5174 Centre National de la Recherche Scientifique (CNRS)–Université Paul Sabatier, 118 Route de Narbonne, F-31062 Toulouse, France*

³*Department of Infectious Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine, Mănăştur Street 3-5, 400372 Cluj-Napoca, Romania*

⁴*Centre d'Etudes Biologiques de Chizé, CNRS UPR 1934, 79360 Beauvoir-sur-Niort, France*

⁵*Office National de la Chasse et de la Faune Sauvage, Direction des Etudes et de la Recherche, CNERA Avifaune migratrice, Carrefour de la Canauderie, 79360 Villiers-en-Bois, France*

The differential allocation hypothesis predicts that females modify their investment in a breeding attempt according to its reproductive value. One prediction of this hypothesis is that females will increase reproductive investment when mated to high-quality males. In birds, it was shown that females can modulate pre-hatch reproductive investment by manipulating egg and clutch sizes and/or the concentrations of egg internal compounds according to paternal attractiveness. However, the differential allocation of immune factors has seldom been considered, particularly with an experimental approach. The carotenoid-based ornaments can function as reliable signals of quality, indicating better immunity or ability to resist parasites. Thus, numerous studies show that females use the expression of carotenoid-based colour when choosing mates; but the influence of this paternal coloration on maternal investment decisions has seldom been considered and has only been experimentally studied with artificial manipulation of male coloration. Here, we used dietary carotenoid provisioning to manipulate male mallard (*Anas platyrhynchos*) bill coloration, a sexually selected trait, and followed female investment. We show that an increase of male bill coloration positively influenced egg mass and albumen lysozyme concentration. By contrast, yolk carotenoid concentration was not affected by paternal ornamentation. Maternal decisions highlighted in this study may influence chick survival and compel males to maintain carotenoid-based coloration from the mate-choice period until egg-laying has been finished.

Keywords: maternal investment; yolk carotenoid concentration; egg mass; albumen lysozyme concentration; paternal ornamentation; carotenoid-based coloration

1. INTRODUCTION

Life-history theory predicts that individual females might modulate their investment in a particular breeding attempt according to its reproductive value because a trade-off may exist between current and future reproduction [1]. One trait that females may use to adjust current effort is the quality of its mate. One prediction of this 'differential allocation' hypothesis (DAH, [2,3]) is that females will increase reproductive investment when mated to high-quality males. In a classic study, Burley [4] showed that female zebra finches (*Taeniopygia guttata*) paired to males with increased attractiveness produced more sons, the more beneficial sex to produce. Such an adjustment prioritizes the fitness of offspring in a current, valuable reproductive bout, given that the prospects for

future survival or optimal mate quality or environmental conditions are comparatively much less certain.

When differentially apportioning reproductive effort by mate quality, there are several putative aspects of breeding that could be adjusted. The most obvious distinction is between pre-hatch (or birth) and post-hatch investments—namely whether embryonic, clutch or brood discrimination occurs at the level of the gamete, the physiological investments prior to hatch or birth, or behavioural adjustments that parents can make for live young. In birds, it was shown that females can modulate pre-hatch reproductive investment by manipulating egg and clutch sizes or the concentration of egg internal compounds [5–8] according to paternal attractiveness (see [9] for a review). However, the differential allocation of immune factors in the eggs according to paternal quality has seldom been considered, particularly with an experimental approach [10,11].

The sexual signals displayed by mates are often recognized as the most salient cues used by females to make differential allocation decisions. These include

* Author for correspondence (giraudeau.mathieu@gmail.com).

† Present address: Centre for Ornithology, School of Biosciences, Birmingham University, Birmingham, UK.

‡ Present address: Leibniz Institute for Zoo and Wildlife Research, Alfred-Kowalke-Strasse 17, 10315 Berlin, Germany.

body size, song parameters, as well as coloration. After melanins, carotenoids are the second most prevalent pigments in the avian integument and determine yellow to red colour of many bird signals [12]. In addition, carotenoid pigments are involved in different physiological processes such as antioxidant and immuno-enhancing activities [12–17]. Thus, growing evidence suggests that individuals face a trade-off concerning the allocation of carotenoid pigments between immune function and ornamentation, which ensures the honesty of carotenoid-based sexual signals [18–21]. In this context, numerous experimental studies have been published on the signaling function of carotenoid-based coloration during the mate choice period (review in the study of Hill [22]; [23,24]), but the potential role of this coloration in maternal investment decisions has seldom been considered [4,7]. Moreover, the role of sexual attractiveness has only been experimentally considered with the use of artificial methods like coloured rings or markers for manipulating male carotenoid-based coloration and signals [2,6,7]. Follow-up studies, where researchers manipulate male carotenoid-based coloration within the natural range of coloration, would constitute a more natural test of the DAH [25].

The aim of this study was to test whether captive female mallards (*Anas platyrhynchos*) adjust current reproductive investment according to the carotenoid coloration of its mate. Male mallards have a bright yellow bill that contains carotenoid pigments [26–28], and females prefer to mate with males displaying yellower bills [29,30]. We used dietary carotenoid supplementation to experimentally manipulate male carotenoid bill coloration and randomly allocated a supplemented or unsupplemented male to each female for the entire first clutch. For the second clutch, we applied the same procedure by randomly allocating a different male, supplemented or not, to each female. This balanced experimental design allowed us to control for any individual variation in maternal reproductive investment that was not owing to mate quality (*sensu* [6,31]). From collected eggs, we quantified yolk carotenoid concentration and albumen lysozyme concentration as response variables, since these two egg parameters have been shown to influence chick development and survival [10,11,32,33]. Lysozyme is a major component of maternal innate antibacterial immunity transferred to the eggs in birds [11,34]. This enzyme acts by digesting bacterial walls [35–40]. Yolk carotenoids are crucial for the development of the embryonic immune system, given their potential protective role against oxidative stress during the first stages of life [41,42]. In addition, we considered clutch and egg size as an index of maternal investment. In mallards, where there is no parental food provisioning after hatch, increasing egg size improves offspring survival by producing heavier progeny [8].

As female mallards are known to increase egg size according to paternal attractiveness (as determined by female preference during mate choice trials; [8]), we predicted that females would lay bigger eggs when paired with supplemented males than with unsupplemented ones. In addition, based on the DAH, we predicted that females paired to more coloured males would deposit more carotenoids and lysozyme in their eggs.

2. METHODS

Experiments were carried out at the Centre d'Etudes Biologiques de Chizé (CEBC) in Western France, using 40 adult duck pairs descended from individuals caught in the wild in three different areas. Birds were kept in semi-captive conditions (grassland of 800 m²) for at least 3 years before the experiments, and were therefore accustomed to their aviary environment. Individuals were fed with an *ad libitum* diet of water and a mixture of crushed corn, wheat and commercial duck food. Experiments were carried out in accordance with French veterinary services.

(a) Carotenoid provisioning

During the mate-choice period (i.e. three months before breeding, January), 40 males were randomly assigned to one of the two treatment groups: carotenoid-supplemented ($n = 20$) or controls ($n = 20$). Supplemented birds received 2 ml of a carotenoid mixture (Oro Glo liquid, 11 mg ml⁻¹ lutein and zeaxanthin; Kemin France SRL, Nantes) every 3 days during two weeks (six treatments) using a plastic syringe that we inserted into their mouth. Unsupplemented birds received the same quantity of water with the same method. In a pilot study, we found that the quantity of carotenoids received by the treated group is the minimum required to observe a significant change in bill coloration one week after the end of the feeding period by spectrophotometry (M. Giraudeau, C. Duval, V. Bretagnolle & P. Heeb 2008, unpublished data). This carotenoid mixture was chosen because carotenoids present in the bill consist of lutein (the largest fraction) and approximately equal fractions zeaxanthin and 3-dehydrolutein [28].

Birds were captured the day before the beginning, 4 days after the end, and 10 weeks after the end of the supplementation for weighing and blood sampling. We drew 500 µl of whole blood from each bird through the alar vein with a heparinized syringe and immediately placed the sample on ice until centrifugation (10 000g for 3 min). Plasma was then frozen at -80°C for later analysis.

(b) Bill colour scoring and analyses

We scored bill colour 1 day before the beginning of the carotenoid supplementation and 10 weeks after the end of this treatment (before the breeding experiment). Bill colour was measured between 300 and 700 nm using an S-2000 spectrophotometer with a DH-2000-FHS deuterium-halogen light source (Ocean Optics, Eerbeek, The Netherlands). Inclusion of the ultraviolet (UV; 320–400 nm) is necessary since ducks are sensitive to UV light [43]. The probe was held at a 90° angle to the bill and reflectance was measured on three different standardized spots under the nostril. Reflectance was calculated relative to a white standard and the three spectra obtained for each bird were averaged and summarized over 1 nm steps. From these spectrum measurements, we extracted hue, chroma, UV-chroma and brightness as response variables to assess variation in true coloration (see Loyau *et al.* [44] for a full description of these colour variables). Computations were conducted with the AVICOL software [45].

(c) Effects of bill colour on egg investment

To examine the effects of male carotenoid provisioning on maternal investment, we used the same experimental design as Cunningham & Russell [8]. Briefly, females were housed individually in a pen of 6 m² for about two weeks. Then, females were assigned one male for their first clutch;

Table 1. Differences in mallard bill coloration between carotenoid-supplemented (CS) and unsupplemented (U) males before the supplementation and 10 weeks later (before breeding, d.f. = 38).

	<i>t</i>	mean (s.e.)CS/U	<i>n</i>	<i>p</i>
start of the experiment				
brightness	-1.04	5111(411)/5772(488)	20/20	0.3
hue	1.58	531(1.9)/528(1.9)	20/20	0.12
chroma	1.52	2.43(0.08)/2.29(0.05)	20/20	0.14
UV-chroma	-1.16	0.07(0.01)/0.08(0.01)	20/20	0.25
before breeding				
brightness	-2.54	4662(240)/5706(331)	20/20	0.01
hue	1.96	530(1.8)/516(10)	20/20	0.05
chroma	3.23	2.4(0.07)/2.07(0.07)	20/20	0.002
UV-chroma	-2.52	0.09(0.01)/0.15(0.02)	20/20	0.02

20 females received a carotenoid-supplemented male and 20 an unsupplemented one. Pairs were housed together in holding pens until females had finished laying their entire clutch. Eggs were collected each day and replaced with a dummy egg to induce females to lay a normal clutch. All the eggs of the clutch were weighed. The first egg of each clutch was split open in a sterile Petri dish. Then, yolk and albumen were separated in two different Petri dishes and each of them was thoroughly mixed. Finally, we sampled 2 ml of albumen and yolk and immediately froze (-80°C) them until the determination of yolk carotenoid concentration and albumen lysozyme concentration.

After females had completed laying and males were removed, females then stayed alone for at least 17 days, the longest period a female mallard has been recorded to store sperm [46], so that all offspring in the second clutch were expected to be fathered only by the second male. For the second clutch, we randomly assigned another supplemented and unsupplemented male to each female and the same procedure was followed during egg-laying and collection. Thus, 10 females were subjected to each of the four possible combinations of mates (i.e. supplemented then supplemented, control then control, supplemented then control or control then supplemented).

(d) Assessment of egg compounds

To measure albumen lysozyme concentration, we used the lysoplate assay method of Osseman & Lawlor [47]: 25 μl of albumen was inoculated in the test holes of a 1 per cent agar gel (A5431, Sigma) containing 50 mg per 100 ml lyophilized *Micrococcus lysodeikticus* (M3770, Sigma) bacteria, which is particularly sensitive to lysozyme concentration. Crystalline hen egg white lysozyme (L6876, Sigma) was used to prepare a standard curve in each plate. Plates were incubated at room temperature ($25-27^{\circ}\text{C}$) for 18 h; during this period, a zone of clearing developed in the area of the gel surrounding the sample inoculation site as a result of bacterial lysis. The diameters of the cleared zones are proportional to the log of the lysozyme concentration. This area was measured by using digital callipers and converted on a semi-logarithmic plot into hen egg lysozyme equivalents (expressed in microgrammes per millilitre) according to the standard curve [48].

Methods for plasma and yolk-carotenoid extractions and high-performance liquid chromatography analyses follow those described in McGraw *et al.* [49].

(e) Statistical procedures

As we weighed all eggs in each clutch, we used mean egg mass per clutch for the analyses using egg mass. We log-transformed yolk carotenoid concentration and albumen lysozyme concentration data to normalize them. To analyse maternal investment parameters (mean egg mass per clutch, clutch size, yolk carotenoid concentration and albumen lysozyme concentration), we performed generalized linear models with treatment of the male allocated to the females (supplemented or not), clutch order (clutch 1 or 2) and the interaction clutch order $\times$ male treatment as a fixed factor, female was included as a random factor. Finally, to test if egg parameters covaried, we have tested correlations between egg mass, albumen lysozyme and yolk carotenoid. Data were analysed using STATISTICA 6.0 software (Statsoft, Tulsa, OK, USA).

3. RESULTS

(a) Plasma carotenoid concentration and bill coloration

Before carotenoid provisioning, supplemented and unsupplemented birds did not differ in mass (d.f. = 38, $t = 0.76$, $p = 0.45$) or tarsus length (d.f. = 38, $t = 0.34$, $p = 0.74$). Repeated-measures analysis of variance revealed no effect of carotenoid provisioning on body mass ($F_{1,38} = 0.9$, $p = 0.35$).

Plasma carotenoid concentrations were not significantly different between supplemented and unsupplemented males at the start of the experiment (d.f. = 38, $t = 0.37$, $p = 0.71$). Repeated-measures revealed that the amounts of circulating plasma carotenoids were not significantly affected by the supplementation (treatment $\times$ change in carotenoids interaction: $F_{1,36} = 1.01$, $p = 0.33$). This absence of difference of plasma carotenoids concentration between our two groups of birds only 4 days after the end of the treatment suggests that we did not give pharmacological dosage of carotenoid during supplementation.

Ten weeks after the end of the supplementation, before the maternal investment experiment, no significant difference of plasma carotenoids was detected between the two groups of males (d.f. = 38, $t = -1.37$, $p = 0.18$).

Before the carotenoid supplementation, males of the two groups did not differ significantly in the four variables of coloration measured (all tests $p > 0.1$; table 1). Carotenoid supplementation had the expected effect of

Figure 1. Maternal investment according to male treatment (carotenoid-supplemented (black bars) and unsupplemented males (grey bars)) during the two clutches (mean $\pm$ s.e.). (a) Egg mass, (b) clutch size, (c) yolk carotenoid concentration, and (d) albumen lysozyme activity.

creating two groups of male mallards that differed significantly in their bill coloration before the maternal investment experiment. Supplemented birds had significantly lower scores of bill brightness and UV-chroma and significantly higher scores of chroma and hue than unsupplemented birds (table 1).

Thus, supplementation did not result in increased blood carotenoid levels only 4 days after supplementation and yet resulted in increased bill coloration when measured 10 weeks later. Two mechanisms could explain these results. First, absorbed carotenoids were transferred to storage organs and later released in bill tissue. Second, only 4 days after the end of the carotenoid provisioning, absorbed carotenoids were taken up directly and completely by binding proteins in bill tissue.

(b) Maternal investment

(i) Egg mass and clutch size

During the experiment, females paired to supplemented males produced heavier eggs than females paired to unsupplemented ones (figure 1a). Moreover, the mass of eggs laid tended to increase in the second clutch. The interaction between male treatment and clutch order did not affect egg mass (table 2).

The number of eggs per clutch decreased significantly from the first (13.46 eggs per clutch) to the second clutch (9.70 eggs per clutch), but no significant effect of male treatment and male treatment $\times$ clutch order interaction was found on clutch size (figure 1b and table 2).

Table 2. Effect of carotenoid provisioning on maternal investment in eggs in captive breeding mallards.

maternal investment variable		$F_{1,75}$	p
clutch size	male treatment	0.03	0.69
	clutch order	8.6	<0.001
	male treatment $\times$ clutch order	0.06	0.73
mean egg mass	male treatment	3.90	0.05
	clutch order	2.74	0.1
	male treatment $\times$ clutch order	0.68	0.41
albumen lysozyme	male treatment	3.97	0.02
	clutch order	0.1	0.98
	male treatment $\times$ clutch order	0.76	0.43
yolk carotenoid	male treatment	0.0001	0.87
	clutch order	15.98	<10 ⁻⁴
	male treatment $\times$ clutch order	1.04	0.52

(ii) Yolk carotenoids

We found that the concentration of carotenoids in the first egg increased with clutch order. However, the deposition of carotenoids in the yolk was not influenced by male treatment (figure 1c). The interaction between clutch order and male treatment did not significantly influence yolk carotenoid concentration (table 2).

(iii) *Albumen lysozyme*

Eggs laid by females paired to supplemented males during the experiment showed a higher lysozyme concentration than eggs laid by females paired to unsupplemented males (table 2). Clutch order and the male treatment $\times$ clutch order interaction did not influence albumen lysozyme concentration (figure 1d).

We found a significant effect of female identity on albumen lysozyme concentration ($F_{1,75} = 2.1$, $p < 0.05$) and egg mass ($F_{1,75} = 3.9$, $p < 0.05$) but not on clutch size ($F_{1,75} = 0.5$, $p = 0.9$) and yolk carotenoid concentration ($F_{1,75} = 1.18$, $p = 0.3$). Finally, we did not find any significant correlation between mean egg mass, carotenoid concentration and lysozyme concentration ($p > 0.15$).

4. DISCUSSION

As predicted by the DAH, our study demonstrates that female mallards modulate egg mass and lysozyme deposition according to experimental manipulations of male carotenoid-based ornamentation. In birds, and particularly mallards, egg size positively influences chick mass and survival during the few days immediately after hatching, since it reflects the overall availability of reserves for the embryos [8,50]. Moreover, albumen lysozyme is involved in the egg's chemical defence, but also in physical defence by forming a fibrous network with other proteins [38]. Thus, a high level of lysozyme in the albumen increases embryo antibacterial defence during development [10,11]. By adjusting these two egg parameters simultaneously, females may influence offspring survival, growth and phenotype [51–54].

Recently, D'Alba *et al.* [55] provided the first correlative investigation of differential maternal investment in egg antimicrobial defence in relation to male attractiveness. Female blue tits (*Cyanistes caeruleus*) increased the amount of lysozyme they transferred to their eggs when mated to a more attractive male (higher UV-chroma). Here, our results constitute, to our knowledge, the first experimental demonstration of a differential allocation of lysozyme into albumen according to paternal coloration. Shawkey *et al.* [56] proposed recently that a trade-off between egg and blood immunity may not exist concerning this enzyme for two reasons: (i) egg and adult lysozyme are produced by different, specialized cells [57–59], and (ii) production and deposition of antimicrobials may be less costly than deposition of other yolk factors, such as carotenoids, antibodies and hormones (calculations from Tristram [60]). However, our data show that the deposition of this antibiotic in albumen is probably beneficial enough in some aspect(s) that females differentially allocate it according to the perceived value of one's mate and the associated reproductive bout. More studies are now needed to determine the immune and energetic costs and benefits of depositing varying amounts of lysozyme into eggs.

Contrary to our predictions, we did not find any effect of paternal coloration on yolk carotenoid concentration. At least two hypotheses could explain these results. First, paternal carotenoid-based coloration may not influence female allocation of carotenoid into eggs in mallards. Futures studies should examine the importance of other male mallard characteristics like plumage coloration on

the differential allocation of carotenoids into eggs in this species. Second, several recent studies on passerine birds showed that results differ depending on whether yolk carotenoid concentration or quantity is considered [61,62]. Safran *et al.* [61] proposed that a higher concentration may increase the dose of a compound during a given interval of embryonic development, whereas a greater amount may mean prolonged exposure during development. Thus, further studies should examine how the total amount of yolk carotenoid varies according to the experimental manipulation of male bill coloration in mallards. We unfortunately did not weigh fresh whole yolks in this study, so we cannot comment on total carotenoid quantity in egg yolks of mallards here.

Our results raise the question of the functional significance of this differential maternal investment driven by the degree of male ornamentation in a species where males do not provide any parental care [8]. First, male mallards may use carotenoid-based coloration to signal their dominance status and thereby the feeding areas available to females and their offspring during the breeding season [8]. It has been suggested in several bird species that carotenoid-based coloration is associated with intense male–male competition for territories [63–67]. However, to our knowledge, this potential role of male mallard bill coloration has not yet been studied.

Second, only males of high genetic quality or males in good phenotypic condition may be able to develop and maintain intense carotenoid-based displays [68,69] and transmit genes for parasitic resistance and attractiveness to offspring [70]. For example, in mallards, male bill colour signals immunocompetence [28]. Thus, females could use carotenoid-based signals to evaluate male genetic quality and may invest more in those advertising good genes for immune status or antioxidant system efficiency [44,71,72]. Third, it was demonstrated in mallards that male bill coloration, and particularly relative UV reflectance, is negatively correlated with sperm velocity [28]. Thus, by modulating egg mass and lysozyme deposition according to male bill UV reflectance, females may increase their investment in a particular breeding attempt with a more fertile male.

During our experiment, clutch order strongly influenced yolk carotenoid concentration and clutch size and tended to influence egg mass, indicating that female mallards laid fewer eggs in the second clutch but produced heavier eggs that contained higher yolk carotenoid concentrations. These results are in accordance with previous studies in passerines, which found that clutch size declines [73,74] and egg size increases [75] with the advance of the breeding season. An explanation is that females may produce heavier chicks with more reserves late in the season to compensate the low availability of food at this period [76].

In conclusion, our findings contribute to the limited information on the deposition of immune compounds in eggs according to paternal attractiveness. Thus, this study provides, to our knowledge, the first experimental demonstration of a differential allocation of lysozyme into eggs. Deposition of lysozyme and egg masses depended on the expression of a paternal secondary sexual character and this may reflect adaptive maternal strategies. In addition, we provide, to our knowledge, the first experimental demonstration of the importance

of carotenoid-based paternal ornamentation on maternal investment, where signals were manipulated directly by carotenoid provisioning. Thus, mothers may alter their investment in egg production in response to male carotenoid-based ornamentation, since attractive males are more likely to sire successful offspring with attractiveness and parasite-resistance genes.

Housing conditions and the experiment were carried out in compliance with European legal recruitment and national permissions (ETS123).

We are grateful to Marco Cucco for his help in lysozyme analyses. We thank Colette Trouvé, Stephanie Dano, André Lacroix and Noël Guillon for their help in the fieldwork and preparation of the samples, and two anonymous referees for their helpful comments. This project was supported by a French research grant (ANR-05, NT05-3_42075) to P.H. During this study, G.Á.C. was supported by a PhD scholarship of the Romanian Ministry of Education and Research and by an Eiffel grant from the French Ministry of Foreign Affairs. K.J.M. was supported by the National Science Foundation (grant no. IOS-0746364).

REFERENCES

- 1 Stearns, S. C. 1992 *The evolution of life histories*. Oxford, UK: Oxford University Press.
- 2 Burley, N. 1986 Sexual selection for aesthetic traits in species with biparental care. *Am. Nat.* **127**, 415–445. (doi:10.1086/284493)
- 3 Burley, N. 1988 The differential allocation hypothesis: an experimental test. *Am. Nat.* **132**, 611–628. (doi:10.1086/284877)
- 4 Burley, N. 1981 Mate choice by multiple criteria in a monogamous species. *Am. Nat.* **117**, 515–528. (doi:10.1086/283732)
- 5 Petrie, M. & Williams, A. 1993 Peahens lay more eggs for peacocks with larger trains. *Proc. R. Soc. Lond. B* **251**, 127–131. (doi:10.1098/rspb.1993.0018)
- 6 Gil, D., Graves, J., Hazon, N. & Wells, A. 1999 Male attractiveness and differential testosterone investment in zebra finch eggs. *Science* **286**, 126–128. (doi:10.1126/science.286.5437.126)
- 7 Velando, A., Beamonte-Barrientos, R. & Torres, R. 2006 Pigment-based skin colour in the blue-footed booby: an honest signal of current condition used by females to adjust reproductive investment. *Oecologia* **149**, 535–542. (doi:10.1007/s00442-006-0457-5)
- 8 Cunningham, E. J. A. & Russell, A. F. 2000 Egg investment is influenced by male attractiveness in the mallard. *Nature* **404**, 74–76. (doi:10.1038/35003565)
- 9 Sheldon, B. C. 2000 Differential allocation: tests, mechanisms and implications. *Trends Ecol. Evol.* **15**, 397–402. (doi:10.1016/S0169-5347(00)01953-4)
- 10 Saino, N., Bertacche, V., Ferrari, R. P., Martinelli, R., Møller, A. P. & Stradi, R. 2002 Carotenoid concentration in barn swallow eggs is influenced by laying order, maternal infection and paternal ornamentation. *Proc. R. Soc. Lond. B* **266**, 1729–1733. (doi:10.1098/rspb.2002.2088)
- 11 Saino, N., Ferrari, R. P., Martinelli, M. R., Rubolini, D. & Møller, A. P. 2002 Early maternal effects mediated by immunity depend on sexual ornamentation of the male partner. *Proc. R. Soc. Lond. B* **269**, 1005–1009. (doi:10.1098/rspb.2002.1992)
- 12 McGraw, K. J. 2006 The mechanics of carotenoid coloration in birds. In *Bird coloration. I. Mechanisms and measurements* (eds G. E. Hill & K. J. McGraw), pp. 177–242. Cambridge, MA: Harvard University Press.
- 13 Burton, G. W. 1989 Antioxidant action of carotenoids. *J. Nutr.* **119**, 109–111.
- 14 Krinsky, N. I. 1989 Antioxidant functions of carotenoids. *Free Radic. Biol. Med.* **7**, 617–635.
- 15 Blount, J. D. & McGraw, K. J. 2008 Control and function of carotenoid coloration in birds: a review of case studies. In *Carotenoids: physical, chemical, and biological functions and properties* (ed. J. T. Landrum), pp. 213–236. Boca Raton, FL: CRC Press.
- 16 El-Agamey, A., Lowe, G. M., McGarvey, D. J., Mortensen, A., Phillip, D. M., George Truscott, T. & Young, A. J. 2004 Carotenoid radical chemistry and antioxidant/pro-oxidant properties. *Arch. Biochem. Biophys.* **430**, 37–48.
- 17 Bendich, A. 1993 Biological functions of dietary carotenoids. *Ann. NY Acad. Sci.* **691**, 61–67. (doi:10.1111/j.1749-6632.1993.tb26157.x)
- 18 Lozano, G. A. 1994 Carotenoids, parasites, and sexual selection. *Oikos* **70**, 309–311. (doi:10.2307/3545643)
- 19 Von Schantz, T., Bensch, S., Grahm, M., Hasselquist, D. & Wittzell, H. 1999 Good genes, oxidative stress and condition-dependent sexual signals. *Proc. R. Soc. Lond. B* **266**, 1–12. (doi:10.1098/rspb.1999.0597)
- 20 Faivre, B., Gregoire, A., Preault, M., Cezilly, F. & Sorci, G. 2003 Immune activation rapidly mirrored in a secondary sexual trait. *Science* **300**, 29–31. (doi:10.1126/science.1081802)
- 21 McGraw, K. J. & Ardia, D. R. 2003 Carotenoids, immunocompetence, and the information content of sexual colours: an experimental test. *Am. Nat.* **162**, 704–712. (doi:10.1086/378904)
- 22 Hill, G. E. 1999. Mate choice, male quality, and carotenoid-based plumage coloration. In *Proc. 22nd Int. Ornithological Congress, Johannesburg, 1999* (eds N. Adams & R. Slotow), pp. 1654–1668. Durban: University of Natal.
- 23 Hill, G. E. 2002 *A red bird in a brown bag: the function and evolution of ornamental plumage coloration in the house finch*. Oxford, UK: Oxford University Press.
- 24 Møller, A. P., Biard, C., Blount, J. D., Houston, D. C., Ninni, P., Saino, N. & Surai, P. F. 2000 Carotenoid dependent signals: indicators of foraging efficiency, immunocompetence, or detoxification ability? *Avian Poultry Biol. Rev.* **11**, 137–159.
- 25 Hill, G. E. 2006 Female mate choice for ornamental coloration. In *Bird coloration. Vol. 2: function and evolution* (eds G. E. Hill & K. J. McGraw), pp. 137–200. Cambridge, MA: Harvard University Press.
- 26 Lonnberg, E. 1938 The occurrence and importance of carotenoid substances in birds. *Int. Orn. Congr.* **8**, 410–424.
- 27 Cramp, S. & Simmons, K. E. L. 1977 *Birds of the Western Palearctic*, vol. 1, pp. 505–519. London, UK: Oxford University Press.
- 28 Peters, A., Denk, A. G., Delhey, K. & Kempenaers, B. 2004 Carotenoid-based bill colour as an indicator of immunocompetence and sperm performance in male mallards. *J. Evol. Biol.* **17**, 1111–1120. (doi:10.1111/j.1420-9101.2004.00743.x)
- 29 Omland, K. E. 1996 Female mallard mating preferences for multiple male ornaments. I. Natural variation. *Behav. Ecol. Sociobiol.* **39**, 353–360. (doi:10.1007/s002650050300)
- 30 Omland, K. E. 1996 Female mallard mating preferences for multiple male ornaments. II. Experimental variation. *Behav. Ecol. Sociobiol.* **39**, 361–366. (doi:10.1007/s002650050301)
- 31 McGraw, K. J., Adkins-Regan, E. & Parker, R. S. 2005 Maternally derived carotenoid pigments affect offspring survival, sex ratio, and sexual attractiveness in a colourful

- songbird. *Naturwissenschaften* **92**, 375–380. (doi:10.1007/s00114-005-0003-z)
- 32 Romano, M., Caprioli, M., Ambrosini, R., Rubolini, D., Fasola, M. & Saino, N. 2008 Maternal allocation strategies and differential effects of yolk carotenoids on the phenotype and viability of yellow-legged gull (*Larus michahellis*) chicks in relation to sex and laying order. *J. Evol. Biol.* **21**, 1826–1840.
 - 33 Biard, C., Surai, P. F. & Møller, A. P. 2005 Effects of carotenoid availability during laying on reproduction in the blue tit. *Oecologia* **144**, 32–44. (doi:10.1007/s00442-005-0048-x)
 - 34 Saino, N., Dall'ara, P., Martinelli, R. & Møller, A. P. 2002 Early maternal effects and antibacterial immune factors in the eggs, nestlings and adults of the barn swallow. *J. Evol. Biol.* **15**, 735–743. (doi:10.1046/j.1420-9101.2002.00448.x)
 - 35 Sato, Y. & Watanabe, K. 1976 Lysozyme in hen blood serum. *Poultry Sci.* **55**, 1749–1756.
 - 36 Clark, A. G. & Bueschgens, D. H. 1986 Survival and growth of *Campylobacter jejuni*. *J. Food Prot.* **49**, 135–141.
 - 37 Grinde, B. 1989 Lysozyme from rainbow trout, *Salmo gairdneri* as an antibacterial agent against fish pathogens. *J. Fish Dis.* **12**, 95–104. (doi:10.1111/j.1365-2761.1989.tb00281.x)
 - 38 Trziszka, T. 1994 Lysozyme and its functions in the egg. *Arch. Geflügelk* **58**, 49–54.
 - 39 Braun, P. & Fehlhaber, K. 1996 Studies of the inhibitory effect of egg albumen on gram-positive bacteria and on *Salmonella enteritidis* strains. *Arch. Geflügel* **60**, 203–207.
 - 40 Kudo, S. 2000 Enzymes responsible for the bactericidal effect in extracts of vitelline and fertilisation envelopes of rainbow trout eggs. *Zygote* **8**, 257–265. (doi:10.1017/S0967199400001052)
 - 41 Pastoret, P., Gabriel, P., Bazin, H. & Govaerts, A. 1998 *Handbook of vertebrate immunology*. San Diego, CA: Academic Press.
 - 42 Saino, N., Ferrari, R. P., Martinelli, S. R., Romano, M., Rubolini, D. & Møller, S. 2003 Early maternal effects mediated by immunity depend on sexual ornamentation of the male partner. *Proc. R. Soc. Lond. B* **269**, 1005–1009. (doi:10.1098/rspb.2002.1992)
 - 43 Parrish, J., Benjamin, R. & Smith, R. 1981 Near-ultraviolet light reception in the mallard. *Auk* **98**, 627–628.
 - 44 Loyau, A., Gomez, D., Moureau, B., Thery, M., Hart, N. S., Saint Jalme, M., Bennett, A. T. D. & Sorcie, G. 2007 Iridescent structurally based coloration of eyespots correlates with mating success in the peacock. *Behav. Ecol.* **18**, 1123–1134. (doi:10.1093/beheco/arm088)
 - 45 Gomez, D. 2007 AVICOL v. 2, a program to analyse spectrometric data. Available upon request from the author at dodogomez@yahoo.fr.
 - 46 Elder, W. H. & Weller, M. W. 1954 Duration of fertility in the domestic mallard hen after isolation from the drake. *J. Wildl. Manage.* **18**, 495–502. (doi:10.2307/3797084)
 - 47 Osserman, E. F. & Lawlor, D. P. 1966 Serum and urinary lysozyme (muramidase) in monocytic and monomyelocytic leukaemia. *J. Exp. Med.* **124**, 921–951. (doi:10.1084/jem.124.5.921)
 - 48 Cucco, M., Guasco, B., Malacarne, G. & Ottonelli, R. 2007 Effects of β -carotene on adult immune condition and antibacterial concentration in the eggs of the grey partridge, *Perdix perdix*. *Comp. Biochem. Physiol. A* **147**, 1038–1046.
 - 49 McGraw, K. J., Gregory, A. J., Parker, R. S. & Adkins-Regan, E. 2003 Diet, plasma carotenoids and sexual coloration in the zebra finch (*Taeniopygia guttata*). *Auk* **120**, 400–410.
 - 50 Rhymer, J. M. 1988 The effect of egg size variability on thermoregulation of mallard *Anas platyrhynchos* offspring and its implications for survival. *Oecologia* **75**, 20–24. (doi:10.1007/BF00378809)
 - 51 Kaplan, R. H. 1985 Maternal influences on offspring development in the California newt, *Taricha torosa*. *Copeia* **85**, 1028–1035.
 - 52 Mousseau, T. A. & Fox, C. W. 1998 *Maternal effects as adaptations*. New York, NY: Oxford University Press.
 - 53 Price, T. 1998 Maternal and paternal effects in birds: effects on offspring fitness. In *Maternal effects as adaptations* (eds T. A. Mousseau & C. H. Fox), pp. 202–226. Oxford, UK: Oxford University Press.
 - 54 Wade, M. J. 1998 The evolutionary genetics of maternal effects. In *Maternal effects as adaptations* (eds T. A. Mousseau & C. H. Fox), pp. 5–21. Oxford, UK: Oxford University Press.
 - 55 D'Alba, L., Shawkey, M., Korsten, P., Vedder, O., Kingma, S. A., Komdeur, J. & Beissinger, S. R. 2010 Differential deposition of antimicrobials proteins in blue tit (*Cyanistes caeruleus*) clutches by laying order and male attractiveness. *Behav. Ecol. Sociobiol.* **64**, 1037–1045.
 - 56 Shawkey, M. D., Kosciuch, K. L., Liu, M., Rohwer, F. C., Loos, E. R., Wang, J. M. & Beissinger, S. R. 2008 Do birds differentially distribute antimicrobial proteins within clutches of eggs? *Behav. Ecol.* **19**, 920–927. (doi:10.1093/beheco/arn019)
 - 57 Mandeles, S. & Ducay, E. D. 1962 Site of egg white protein formation. *J. Biol. Chem.* **237**, 3196–3199.
 - 58 Shutz, G., Nguyen-Huu, M. C., Giesecke, K., Hynes, N. E., Groner, B., Wurtz, T. & Sippel, A. E. 1978 Hormonal control of egg-white protein messenger RNA synthesis in the chicken oviduct. *Cold Spring Harb. Symp. Quant. Biol.* **42**, 617–624.
 - 59 Moen, R. C. & Palmiter, R. D. 1980 Changes in hormone responsiveness of chicken oviduct during primary stimulation with estrogen. *Dev. Biol.* **78**, 450–463. (doi:10.1016/0012-1606(80)90346-2)
 - 60 Tristram, G. R. 1953 Amino acid composition of the proteins. In *The proteins* (eds H. Neuarth & K. Bailly), pp. 181–233. New York, NY: Academic Press Inc.
 - 61 Safran, R. J., Pilz, K. M., McGraw, K. J., Correa, S. M. & Schwabl, H. 2008 Are yolk androgens and carotenoids in barn swallow eggs related to parental quality? *Behav. Ecol. Sociobiol.* **62**, 427–438. (doi:10.1007/s00265-007-0470-7)
 - 62 Pilz, K. M., Smith, H. G., Sandell, M. & Schwabl, H. 2003 Inter-female variation in egg yolk androgen allocation in the European starling: do high quality females invest more? *Anim. Behav.* **65**, 841–850. (doi:10.1006/anbe.2003.2094)
 - 63 Roskaft, E. & Rohwer, S. 1987 An experimental study of the function of the red epaulettes and black body colour of male red-winged blackbirds. *Anim. Behav.* **35**, 1070–1077. (doi:10.1016/S0003-3472(87)80164-1)
 - 64 Evans, M. R. & Hatchwell, B. J. 1992 An experimental study of male adornment in the scarlet-tufted Malachite sunbird: II. The role of the elongated tail in mate choice and experimental evidence for a handicap. *Behav. Ecol. Sociobiol.* **29**, 421–427.
 - 65 Pryke, S. R., Anderson, S., Lawes, M. J. & Piper, S. E. 2002 Carotenoid status signalling in captive and wild red-collared widowbirds: independent effects of badge size and colour. *Behav. Ecol.* **13**, 622–631. (doi:10.1093/beheco/13.5.622)
 - 66 Pryke, S. R. & Anderson, S. 2003 Carotenoids-based epaulettes reveal male competitive ability; experiments with resident and floater red-shouldered window-birds. *Anim. Behav.* **66**, 217–224. (doi:10.1006/anbe.2003.2193)

- 67 Pryke, S. R. & Anderson, S. 2003 Carotenoid-based status signalling in red-shouldered widow-birds (*Euplectes axillaris*): epaulet size and redness affect captive and territorial competition. *Behav. Ecol. Sociobiol.* **53**, 393–401.
- 68 Trivers, R. L. 1972 Parental investment and sexual selection. In *Sexual selection and the descent of man* (ed. B. Campbell), pp. 136–179. Chicago, IL: Aldine.
- 69 Zahavi, A. 1975 Mate selection: a selection for a handicap. *J. Theor. Biol.* **53**, 205–214. (doi:10.1016/0022-5193(75)90111-3)
- 70 Hamilton, W. D. & Zuk, M. 1982 Heritable true fitness and bright birds: a role for parasites. *Science* **218**, 384–387. (doi:10.1126/science.7123238)
- 71 Bendich, A. 1989 Carotenoids and the immune response. *J. Nutr.* **119**, 112–115.
- 72 Blount, J. D., Metcalfe, N. B., Birkhead, T. R. & Surai, P. F. 2003 Carotenoid modulation of immune function and sexual attractiveness in zebra finches. *Science* **300**, 125–127. (doi:10.1126/science.1082142)
- 73 Klomp, H. 1970 The determination of clutch size in birds. A review. *Ardea* **58**, 1–124.
- 74 Nilsson, J. A. 1991 Clutch size determination in the marsh tit (*Parus palustris*). *Ecology* **72**, 1757–1762. (doi:10.2307/1940974)
- 75 Verhulst, S., Van Balen, J. H. & Tinbergen, J. N. 1995 Seasonal decline in reproductive success of the great tit: variation in time or quality? *Ecology* **76**, 2392–2403. (doi:10.2307/2265815)
- 76 Nilsson, J. A. 2000 Time-dependent reproductive decisions in the blue tit. *Oikos* **88**, 351–361. (doi:10.1034/j.1600-0706.2000.880214.x)