Published in final edited form as:

Tetrahedron Lett. 2011 March 16; 52(11): 1149–1150. doi:10.1016/j.tetlet.2011.01.047.

Chemoenzymatic Synthesis of 2-Arachidonoylglycerol, An Endogenous Ligand for Cannabinoid Receptors

Subramanian K Vadivel^{*}, Kyle M Whitten, and A. Makriyannis

Center for Drug Discovery, 116 Mugar Hall, 360 Huntington Avenue, Northeastern University, Boston, MA 02115

Abstract

A simple and efficient synthesis of 2-arachidonoyl glycerol, an endogenous agonist for cannabinoid receptors was achieved using Novozym 435, immobilized lipase from *Candida Antarctica*.

Keywords

Cannabinoid; endogenous ligand; 2-arachidonoylglycerol; biocatalysis

2-Arachidonoylglycerol (2-AG) is an endogenous cannabinergic ligand that interacts with both the CB1 and CB2 receptors. Although 2-AG synthesis involves several candidate enzymes,¹ 2-AG is inactivated principally by monoacylglycerol lipase²(MGL), although fatty acid amide hydrolase(FAAH) may contribute to its degradation.³ 2-AG was shown to possess various biological activities such as binding to the CB1 and CB2 cannabinoid receptors, inhibition of adenylyl cyclase in mouse spleen cells, and inducing hypothermia, reducing spontaneous activity, analgesia, and immobility in mice.⁴ 2-AG acts as a full cannabinergic agonist, and the structure of 2-AG is strictly recognized by the cannabinoid receptors (CB1 and CB2). Thus, 2-AG-rather than anandamide may represent the true natural ligand for cannabinoid receptors.^{5–7}

The major problem in the synthesis of pure 2-AG is the rapid migration of the arachidonoyl group from secondary to the primary hydroxyl group, resulting in the formation of more stable 1-arachidonoyl glycerol. This migration is catalyzed by water, acid, base or heat.⁸ Earlier synthetic methods utilized coupling of 1,3-silyl⁹ or benzylidene¹⁰ protected glycerol with arachidonic acid and followed by deprotection and separation of the isomeric arachidonoyl glycerols. All these methods suffer from extended reaction time, acidic conditions required for the removal of the protecting groups as well as extensive work up and purification. Another interesting method also appeared in the literature which utilizes regioselective transformation of glycidyl arachidonate into 2-arachidonoyl-1,3-bis(trifluoroacetyl)glycerol followed by the cleavage of trifluoroacetyl group with pyridine.¹¹

Searching for an alternative, green and efficient methodology that would circumvent these problems during 2-arachidonoyl glycerol synthesis, we have developed an enzyme catalyzed

^{*}To whom correspondence should be addressed Center for Drug Discovery, Northeastern University, 360 Huntington Avenue, 116 Mugar LifeSciences Building, Boston, MA, 02115, USA. Tel.: +1 617 373 7620; Fax: +1 617 373 7493; s.vadivel@neu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

efficient and highly regioselective synthesis. The advantage of biocatalysis is that the reactions are carried out at ambient temperature, nearly neutral pH and the reactions are often highly regio- and stereoselective.^{12–13} Enzymes derived from microbial cells can be immobilized and reused for many cycles. It was known that enzymes can show selectivity during glycerolysis of fatty acids as well as transesterification of symmetrical triglycerides.^{14–18} Herein, we report an improved, practical application of the above methodology in the synthesis of 2-arachidonoyl glycerol utilizing unsymmetrical triglyceride.

The synthesis is based on a two-step enzymatic process (Scheme-1). The first involves the synthesis of 1,3-diacyl glycerol using the method reported by Halldorsson *et al.*¹⁹ Enzymatic acylation of glycerol was carried out in anhydrous dichloromethane using vinylbutyrate as an acyl transfer agent. The reaction proceeded smoothly at 0°C to provide exclusively 1,3-butyroyl glycerol(**1**) with greater than 90% yield.²⁰ The 1,3-diacylglycerol was coupled with arachidonic acid using 4-dimethylaminopyridine (DMAP) and 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI) in dichloromethane at room temperature for 12 h provided the required triglyceride(**2**) in 84% yield.²¹ Triglyceride(**2**) was subjected to immobilized *Candida antarctica* (Novozym 435) known for its high 1,3-regioselective glycerolysis of triglycerides. The reaction was facile and afforded the required 2-arachidonoyl glycerol(**3**) exclusively (67%) and the by-product butylacetate was easily removed under vacuum. We also noted the formation of ethyl arachidonate (27%) as a side product. It is noteworthy that there was no formation of 1-arachidonoylglycerol observed in the reaction. The crude product was purified by a small filter column of boric acid impregnated silica.²²

In summary, the improved and practical synthesis of 2-AG was successfully carried out without any isomerization of the more stable 1-arachidonoyl glycerol. The mild neutral conditions, easy scalability and removal of the volatile byproduct provide added advantages to the current method for the synthesis of 2-monoacylglycerides.

Acknowledgments

One of the authors (SKV) acknowledges the financial support for this research from NIDA (R03 DA029184-01).

References and Notes

1. Di Marzo V, Petrosino S. *Curr Opin Lipidol* 2007;18:129. [PubMed: 17353660]
2. Lambert DM, Fowler CJ. *J Med Chem* 2005;48:5059. [PubMed: 16078824]
3. Goparaju SK, Ueda N, Yamaguchi H, Yamamoto S. *FEBS Lett* 1998;422:69. [PubMed: 9475172]
4. Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminski NE, Schatz AR, Gopher A, Almog S, Martin BR, Compton DR, et al. *Biochem Pharmacol* 1995;50:83. [PubMed: 7605349]
5. Sugiura T, Kishimoto S, Oka S, Gokoh M. *Prog Lipid Res* 2006;45:405. [PubMed: 16678907]
6. Sugiura T, Kodaka T, Nakane S, Miyashita T, Kondo S, Suhara Y, Takayama H, Waku K, Seki C, Baba N, Ishima Y. *J Biol Chem* 1999;274:2794. [PubMed: 9915812]
7. Sugiura T, Kondo S, Kishimoto S, Miyashita T, Nakane S, Kodaka T, Suhara Y, Takayama H, Waku K. *J Biol Chem* 2000;275:605. [PubMed: 10617657]
8. Martin JB. *J Amer Chem Soc* 1953;75:5483.
9. Han LN, Razdan RK. *Tetrahedron Lett* 1999;40:1631.
10. Seltzman HH, Fleming DN, Hawkins GD, Carroll FI. *Tetrahedron Lett* 2000;41:3589.
11. Stamatov SD, Stawinski J. *Tetrahedron Lett* 2002;43:1759.
12. Nakamura K, Yamanaka R, Matsuda T, Harada T. *Tetrahedron Asymmetry* 2003;14:2659.
13. Matsuda T, Yamanaka R, Nakamura K. *Tetrahedron: Asymmetry* 2009;20:513.
14. Monteiro JB, Nascimento MG, Ninow JL. *Biotechnol Lett* 2003;25:641. [PubMed: 12882159]

15. Pfeffer J, Freund A, Bel-Rhliid R, Hansen CE, Reuss M, Schmid RD, Maurer SC. *Lipids* 2007;42:947. [PubMed: 17623117]
16. Damstrup ML, Jensen T, Sparso FV, Kiil SZ, Jensen AD, Xu X. *J Am Oil Chem Soc* 2005;82:559.
17. Damstrup ML, Jensen T, Sparso FV, Kiil SZ, Jensen AD, Xu X. *J Am Oil Chem Soc* 2006;83:27.
18. Irimescu R, Iwasaki Y, Hou CT. *J Am Oil Chem Soc* 2002;79:879.
19. Halldorsson A, Magnusson CD, Haraldsson GG. *Tetrahedron* 2003;59:9101. NMR data for compound **1**: ^1H NMR (CDCl_3 , 500 MHz) δ 4.12–4.27 (m, 4H), 4.03–4.12 (m, 1H), 2.20–2.41 (t, $J = 7.4$ Hz, 4H), 1.67 (sxt, $J = 7.42$ Hz, 4H), 0.96 (t, $J = 7.32$ Hz, 6H).
21. Preparation of **2**: To a solution of 1,3-dibutanoylglycerol (**1**) (0.500g, 2.45mmol) and arachidonic acid (0.74g, 2.45mmol) in anhydrous dichloromethane (10mL) was added DMAP (2.4g, 19.6mmol) and EDCI (1.88g, 9.8 mmol). The resulting solution was stirred for 24h. Cold water was added to the reaction mixture and washed with 5% HCl. The aqueous layer was extracted with dichloromethane and the combined organic extracts were washed with water, brine and dried over sodium sulfate. The reaction mixture was subjected to flash chromatography (hexanes/acetone; 7/3) to afford pure triglyceride as a colorless oil (1.06g, 88%). IR (neat) 3012, 2931, 1741, 1167, 1094 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz) δ 5.30–5.45 (m, 8H), 5.23–5.30 (m, 1H), 4.10–4.37 (m, 4H), 2.76–2.91 (m, 6H), 2.34 (t, $J = 7.57$ Hz, 2H), 2.30 (t, $J = 7.57$ Hz, 4H), 2.12 (q, $J = 7.16$ Hz, 2H), 2.06 (q, $J = 7.32$ Hz, 2H), 1.68–1.75 (m, 2H), 1.60–1.68 (m, 4H), 1.23–1.42 (m, 6H), 0.91–0.99 (m, 6H), 0.84–0.91 (m, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 173.3, 172.8 (2C), 130.7, 129.2, 128.9, 128.8, 128.5, 128.3, 128.0, 127.7, 69.2, 62.3 (2C), 36.1 (2C), 33.8, 31.7, 29.5, 27.4, 26.7, 25.8 (3C), 24.9, 22.8, 18.5 (2C), 14.3, 13.8 (2C).
22. Preparation of 2-arachidonoyl glycerol (**3**): Triglyceride **2** (0.050g, 0.102mmol) and anhydrous ethanol (0.7mL) were stirred at room temperature and the reaction was started by addition of Novozym 435 (75mg) and the reaction mixture was stirred for 1 h and the starting material was completely consumed. An additional 50mg was added and the reaction mixture was stirred for another 1 h. After completion of the reaction the enzyme was filtered off and washed with ether and the solvent was evaporated and the crude product was purified by plug of boric acid impregnated silica gel (hexanes/acetone; 9/1-3/2) to afford analytically pure 2-AG (0.026g, 67%) as a colorless oil. IR (neat) 3210, 3012, 2927, 1736, 1475, 1417, 1377, 1277, 1177 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 5.37–5.49 (m, 8H), 4.93 (p, 5.0 Hz, 1H), 3.84 (t, $J = 5.0$ Hz, 4H), 2.79–2.88 (m, 6H), 2.39 (t, $J = 7.5$ Hz, 2H), 2.13 (q, $J = 7.0$ Hz, 2H), 2.06 (q, $J = 7.0$ Hz, 2H), 2.00 (brt, $J = 5.5$ Hz, 2H), 1.72 (p, $J = 7.2$ Hz, 2H), 1.30–1.40 (m, 6H), 0.92 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 173.2, 130.7, 129.3, 128.9, 128.8, 128.5, 128.3, 128.1, 127.8, 67.51, 65.0, 64.9, 33.8, 31.7, 29.6, 27.4, 26.7, 25.8, 25.8, 25.8, 24.9, 22.8, 14.3.

Scheme 1.