Transcatheter Embolization in Pelvic Trauma

Scott R. Broadwell, M.D.,¹ and Charles E. Ray, Jr., M.D.^{1,2}

ABSTRACT

This article outlines the clinical and radiological approach to patients with pelvic trauma. It reviews the clinical descriptions and classifications of pelvic trauma as well as the management of patients with stable and unstable fractures. A review of embolization techniques in these types of patients is provided.

KEYWORDS: Embolization, trauma, fractures

Objectives: Upon completion of this article, the reader will understand which patients are best suited for pelvic embolization, and be able to describe general techniques used in such procedures.

Accreditation: Tufts University School of Medicine (TUSM) is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians. TUSM takes responsibility for the content, quality, and scientific integrity of this CME activity.

Credit: TUSM designates this educational activity for a maximum of 1 Category 1 credit toward the AMA Physicians Recognition Award. Each physician should claim only those credits that he/she actually spent in the activity.

Since it was first described 30 years ago,¹ percutaneous transcatheter embolization has been used successfully in the management of patients with hemorrhage from pelvic fractures. Pelvic fractures are associated with significant mortality risk from uncontrolled bleeding, and for this reason aggressive, expedient, and streamlined treatment of patients with known or suspected pelvic hemorrhage is crucial to improve their survival. Many, if not most, level I trauma centers in this country and elsewhere have adopted algorithm-based clinical management pathways to optimize the outcomes in this potentially lethal condition. Within these algorithms, transcatheter embolization (TCE) for pelvic arterial hemorrhage has established itself as a standard of treatment in the trauma patient with evidence of ongoing blood loss.²⁻¹² This article reviews current management concepts for the hemodynamically unstable patient with blunt trauma to the pelvis; it also discusses the role of arterial embolization and the

techniques, as well as the limitations, complications, and pitfalls associated with TCE.

PELVIC FRACTURES AND MORTALITY

The osseous and ligamentous structure of the pelvic ring provides an inherently stable configuration that requires a great deal of energy to disrupt. McCoy et al¹³ estimated that the energy required to disrupt this ring is 6300 J, or that generated in the pelvis of a motor vehicle occupant instantaneously decelerated from 30 to 0 miles per hour. Grimm et al¹⁴ found that the mean force required to disrupt the pelvis (in their cadaveric open-book pelvic fracture model) was 3481 N. It is therefore not surprising that most pelvic fractures are associated with high-energy trauma. Motor vehicle collisions, pedestrians struck by a motor vehicle, motorcycle accidents, falls from a height, and industrial accidents account for the vast majority of pelvic fractures encountered in

Interventional Radiology on Embolization; Editor in Chief, Peter R. Mueller, M.D.; Guest Editors, Matthew A. Mauro, M.D., and Charles Ray, M.D. *Seminars in Interventional Radiology*, volume 21, number 1, 2004. Address for correspondence and reprint requests: Charles E. Ray, M.D., Denver Health Medical Center, 777 Bannock Street, Mail Code 0024, Denver, CO, 80204. E-mail: cray@dhha.org. ¹Department of Diagnostic Radiology, University of Colorado Health Sciences Center, Denver, Colorado; ²Department of Interventional Radiology, Denver Health Medical Center, Denver, Colorado. Copyright © 2004 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662. 0739-9529,p;2004,21,01,023,036,ftx,en;sir00224x.

Table 1 Mechanism of Injury in Pelvic Fracture from Blunt Trauma in Three Recent Large Studies

Study	n	MVC	PV	FFH	MC	Other
Poole, Ward ¹⁹	348	228 (66%)	46 (13%)	16 (5%)	19 (5%)	39 (11%)
Demetriades et al ²³	1545*	605 (39%)	647 (42%)	180 (12%)	47 (3%)	28 (2%)
Henry et al ²⁴	234 [†]	150 (64%)	44 (19%)	22 (9%)		17 (7%)
Totals	2127	983 (46%)	948 (45%)	218 (10%)	66 (3%)	84 (4%)

n, number of patients in study with pelvic fracture; MVC, motor vehicle collision; PV, pedestrian struck by motor vehicle; FFH, fall from height; MC, motorcycle crash.

*Mechanism of injury not accounted for in 38 patients.

[†]Mechanism of injury not accounted for in one patient.

civilian trauma.^{15–24} Table 1 reviews the reported mechanism of injury for pelvic fractures in three large studies. Interestingly, motor vehicle collision may produce fewer pelvic fractures per incident than other common causes.²³

The massive force required to fracture the pelvis is rarely isolated to this anatomic region; injuries to other sites and organ systems are noted frequently. As a result, pelvic fracture patients frequently will manifest severe injuries to the central nervous system, thorax, abdomen, and extremities. These associated injuries have been shown to be the leading cause of death in patients with severe pelvic fractures.^{17–19,22–24} However, the mortality from hemorrhagic complications of pelvic fractures is not insignificant, and hemorrhage from pelvic fractures is a particularly important cause of early deaths.^{15,25,26}

Mortality rates in the literature vary widely and are summarized in Table 2. The large variation is multifactorial and may be due to different classification systems used, variability in inclusion criteria with respect to fracture severity, variable numbers of patients with significant associated trauma, and inclusion of patients treated before the advances in modern trauma care that

have resulted in improved outcomes overall.²⁷ Further complicating the data, most researchers reporting mortality data have excluded patients who die in the prehospital setting. A notable recent exception is provided by Adams et al,²⁶ who upon reviewing 392 fatalities from motor vehicle collision, found that 82 (23%) had sustained pelvic fractures, and that the fracture was ruled the cause of death in 63 (77%) of these patients. They conclude that excluding those individuals who do not survive long enough to reach the hospital might result in the underreporting of mortality from pelvic fractures. Their conclusion may have significant implications as prehospital care continues to improve; it is not unreasonable to expect that more patients will reach trauma centers alive and with hemodynamically significant pelvic fractures requiring emergent treatment.

In our review of the literature, we would summarize that although the mortality of patients with pelvic fractures appears to be less than previously reported, hemorrhage from pelvic fracture is not insignificant and—as suggested by Adams et al²⁶—may be underreported. Furthermore, the decrease in mortality has occurred concurrently with advances in trauma care, which include more widespread use of TCE. In addition, recognition of associated injuries as the leading cause of death has significant implications for the angiographer, who can play an integral role in the diagnosis and treatment of patients with hemodynamically significant hemorrhage from injuries outside of the pelvis. This role is discussed elsewhere in this issue of *Seminars in Interventional Radiology*.

Table 2 Pelvic Fracture Mortality

Study	No. of Patients	Deaths	Mortality (%)
Rothenberger et al ¹⁵	604	72	12
Gilliland et al ²⁵	100	30	30
Evers et al ¹⁶	245	35	14
Poole et al ¹⁷	236	18	7.6
Gruen et al ¹⁸	36	4	11
Poole, Ward ¹⁹	348	28	8
Muir et al ²⁰	111*	56	50
Eastridge, Burgess ²⁴	111	26	23.4
Jones et al ²¹	39	10	25.6
Parreira et al ²²	103	20	19
Demetriades et al ²³	1545	208	13.5
Mucha et al ⁴	533	34	6.4 [†]

*153 patients with blunt trauma sustained a pelvic fracture. Of these, 42 died prior to reaching a hospital. If these patients were to be included, percent mortality in this study would increase to 64.

[†]When categorizing patients into initially hemodynamically stable versus unstable, mortality rates were 3.4 and 42%, respectively.

TYPES OF HEMORRHAGE IN PELVIC FRACTURES

Hemorrhage from pelvic fractures occurs through a variety of mechanisms. As well-stated by Eastridge et al,²⁴ “Pelvic fractures from high-energy blunt mechanisms of injury are at considerable risk for major hemorrhagic sequela secondary to disruptions of the intimate relationship between the arterial vascular supply and the pelvic sidewall, disruption of the rich presacral venous plexus (retroperitoneal hemorrhage), and from fracture of the vascular cancellous bone within the pelvis.” In addition to bleeding from cancellous bone,

vessels can be injured from one or more of the following mechanisms: shearing of vessels against fixed ligamentous structures, avulsion of vessels attached to a displaced pelvic segment, penetration injury from a shard of bone, and iatrogenic lesions arising from resuscitative interventions.²⁸ Contributing to further blood loss, hemorrhage may be complicated by coagulopathic states induced by hypothermia or fluid resuscitation.

PELVIC FRACTURE CLASSIFICATION

There are several classification schemes for pelvic fractures. One of their primary values lies in standardizing the terminology used by the orthopedic surgeon, trauma surgeon, and interventional radiologist. The result is improved communication among the trauma team. At our institution, we use the Pennal and Tile system as modified by Young and Burgess.²⁹ As will be discussed below, certain fracture classes in this system have been shown to correlate with increased prevalence of arterial injury. Additionally, this classification system should be commended for its descriptive, logical naming scheme. Eponymous classification systems do not aid in our understanding of mechanisms of injury and in our opinion should be abandoned. The Young-Burgess classification system and the anatomy of the bony pelvis will be briefly reviewed.

The bony pelvis consists of the sacrum, coccyx, and paired iliac, ischial, and pubic bones. The stability of the pelvis is provided primarily by the strong pelvic ligaments, which include interosseous, anterior, and posterior sacroiliac ligaments; sacrotuberous ligaments; sacrospinous ligaments; with additional support provided by the iliolumbar and lateral lumbosacral ligaments.²⁹ The bony pelvis contains four articulations; two are synovial, the sacroiliac joints, and two are secondary cartilaginous joints, the symphysis pubis and the sacrococcygeal joint.³⁰ High-energy trauma to the pelvis will result in fracture or disruption of these structures to different degrees, which in turn will result in varying levels of instability.

In their retrospective radiographic analysis of 142 patients with fractures of the pelvis, Young and Burgess²⁹ found that the patterns of fracture and ligament disruption could be categorized based on the four principal types of force that created them: anteroposterior compression (APC), lateral compression (LC), vertical shear (VS), and combined mechanical (CM). They found that 94% of fractures were correctly diagnosed using the radiographic AP projection alone; however, using a three-view approach (AP, inlet, and outlet views) of the pelvis increases the accuracy of plain radiography even further.

APC injuries result when the force vector acts to rotate externally one hemipelvis. This type of mechanism is one of distraction at the symphysis and ipsilateral

sacroiliac joint (SIJ), and can be recognized by the vertical fractures of the pubic rami and/or disruption of ligaments of the symphysis associated with it.²⁹ As the magnitude of the external rotating force increases, the SIJ becomes increasingly unstable from disruption of the anterior sacroiliac, sacrospinous, sacrotuberous, and the posterior sacroiliac ligaments. Tile³¹ showed that diastasis of the symphysis pubis up to 2.5 cm could occur without disruption of posterior ligaments; however, a diastasis of less than this does not guarantee the integrity of these same ligamentous structures.

Under the Young-Burgess system,²⁹ APC injuries are further subdivided into types I, II, and III based on the degree of posterior injury. In type I fractures, no SIJ diastasis occurs, and the symphysis pubis may be widened up to 2.5 cm. Type II fractures are associated with anterior SIJ diastasis but posterior SIJ ligaments are intact, whereas type III injuries are associated with variable symphysis pubis diastasis and anterior and posterior SIJ diastasis resulting from complete ligamentous compromise.³²

The second major class of force vectors resulting in recognizable fracture patterns is the LC type. Energy is transmitted laterally from the iliac wing medially to the pubic rami, the SIJ, and sacrum. These injuries can be recognized by their horizontally or coronally-oriented pubic rami fractures.²⁹ Under the Young-Burgess system,²⁹ LC injuries are also subdivided into three types. In the LC type I class, the transmitted injury is to the posterior portion of the pelvis. The compressive nature seen in this type of injury may result in crush fractures of the sacrum but the ligaments remain intact, and therefore pelvic stability is not significantly compromised. If the force moment is anterior, as with type II and III fractures, the result is internal rotation of the ipsilateral hemipelvis, with the sacrum or anterior SIJ acting as a fulcrum. Compression fractures of the sacrum and fractures of the posterior iliac wing can therefore be seen. If the force is great enough, it can be transmitted to the contralateral hemipelvis where its action is one of external rotation, causing anterior sacroiliac ligament injury and SIJ diastasis to occur. This latter situation typifies the LC type III class.

The third class of injuries is the VS. These injuries occur when a vertically directed force is transmitted to the pelvis lateral to the midline. These injuries frequently occur in patients falling from a height (and landing feet first), where downward forces are transmitted through the torso to the midline sacrum, and opposing forces are transmitted from lower extremities upward lateral to the midline. This type of injury results in severe derangement of pelvic ligaments and pelvic instability. Vertically oriented fractures may occur in pubic rami, sacrum, or iliac bones. The pelvic outlet radiograph is the best projection for visualizing the degree of fracture displacement.²⁹

APC, LC, and VS vectors can occur in isolation from each other or in a combined fashion. This combined injury is the case in the fourth class of fractures in the Young-Burgess system.²⁹ When combined mechanisms are involved, the injuries are classified as complex pattern or CM, and some combination of the above fracture patterns may occur. Again, recognition of the fracture pattern (as a means to classify) is made somewhat easier by noting that vertical fractures through the pubic rami are associated with APC vectors, whereas horizontal or coronal fractures of the pubic rami are associated with LC vectors.²⁹

HEMORRHAGE RISK BASED ON TYPE/CLASS OF FRACTURE

Another utility of the Young-Burgess system²⁹ might be that certain classes are associated with an increased risk of hemorrhage. Those classes are APC II and III, LC III, VS, and CM.³³ This concept was recently supported by the work of Hamill et al,³⁴ who tested the hypothesis that these classes are associated with a greater need for pelvic arterial embolization. They found that although a correlation existed, it was not strong enough to warrant a change in their indications for angiography. Of special note, pelvic fractures in the elderly are more likely to cause significant hemorrhage, to require transfusion, to require angiography, and to be fatal, regardless of the force vectors.³⁵

The above statements are not without controversy in the literature. Many authors would argue that fracture type or class in determining outcome or site of vascular injury is somewhat limited. Specifically, Cook et al¹² found in their analysis of 150 consecutive patients with unstable fractures of the pelvis that morphology of the fracture was not a reliable guide to the associated vascular injury. Although they did not use the Young-Burgess classification system, Parreira et al²⁴ found that the outcome of patients with pelvic fractures due to blunt trauma correlates with the severity of the associated injuries rather than with the characteristics of the fracture type.

MANAGEMENT OF UNSTABLE PELVIC FRACTURES/HEMORRHAGE

The management of the hemodynamically unstable patient with pelvic fractures remains a diagnostic and therapeutic challenge. An aggressive, multispecialty approach is indicated by most authors. In addition to the host of ancillary support and nursing staff required, the team should be composed of anesthesiologists, emergency medicine physicians, interventional radiologists, and orthopedic and trauma surgeons. The major sources of bleeding need to be identified and controlled as quickly as possible. This will require an organized,

expedient, and coordinated effort from the resuscitative team. An algorithm approach might help achieve this goal.

Using our algorithm and experience as an example,¹¹ the interventional radiologist can be notified of the patient potentially requiring angiography and/or embolotherapy for ongoing pelvic arterial bleeding after the diagnosis of pelvic fracture has been made. This diagnosis will have been reached through a combination of physical examination and radiographic findings. Assessment for thoracic and/or abdominal sources of significant hemorrhage can be performed with a portable chest radiograph and either abdominal ultrasound or diagnostic peritoneal lavage (DPL). Based on the results of their 1989 study, Evers et al¹⁶ recommended pelvic angiography should be performed before laparotomy in hemodynamically unstable patients with pelvic fracture, unless the DPL is grossly positive.

Contrast Enhanced Computed Tomography

As mentioned above, identification of patients likely to benefit from TCE remains a technical and logistical challenge. With the improved throughput of modern multislice scanners, contrast-enhanced computed tomography (CECT) may be useful. In their retrospective study of 290 patients with pelvic fractures who had CECT performed during the early resuscitative period, Pereira et al³⁶ found that lack of contrast extravasation during CT scanning had a 99.6% negative predictive value in determining the need for angiographic embolization. Therefore, lack of contrast extravasation during early CECT (in patients stable enough to undergo this examination) may be an excellent means to rule out patients likely to require TCE.

Contradictory to the above findings by Pereira et al,³⁶ we have not found CECT to be a valuable tool in assessing which patients will require TCE in patients who present with **hemodynamic instability**. If contrast extravasation is visualized, active bleeding is occurring and TCE is immediately indicated. However, if contrast extravasation is not visualized, we believe that an arterial source is not excluded and the patient undergoes exploratory angiography. Therefore, in the **unstable patient**, the results of the CECT do not dictate whether or not the patient will undergo angiography; the CECT, therefore, is not typically obtained in hemodynamically unstable patients in our institution.

Diagnostic Angiography

Digital subtraction angiography (DSA) remains the gold standard for the diagnosis of arterial injury. As will be discussed below, diagnosis through the use of DSA has the unique added benefit of providing the opportunity

Figure 1 A 24-year-old unhelmeted motorcyclist involved in a high speed motor vehicle accident. (A) Digital subtraction aortogram, early arterial phase, demonstrating contrast extravasation arising from the left obturator artery. (B) Digital subtraction aortogram, late arterial phase, during the same injection as shown in (A). Note the pooling of contrast near the pubic rami on the left. (C) Selective injection following embolization of the anterior division of the left internal iliac artery. The anterior division was quickly embolized because of the bleeding from the obturator branch. Unfortunately, the patient had an obturator artery replaced to the posterior division: persistent bleeding is still noted from the replaced artery. (D) Final angiogram demonstrating occlusion of both the anterior and posterior divisions of the internal iliac artery. Gelfoam slurry was used in both cases. Adapted from Biffl et al.¹¹

for concomitant, definitive treatment through the use of TCE. In hemodynamically unstable patients with pelvic fractures, the yield of angiography in identifying a pelvic source of bleeding ranges from 43 to 78%.^{8,9,11,12}

At angiography, the following might be encountered: active extravasation of contrast, pseudoaneurysm, arteriovenous fistulae, intimal tears, vessel occlusion, spasm, and/or thrombosis. In a series of 63 patients, contrast extravasation—evidence for transection or laceration—was the most common finding.³ The same study demonstrated that the following arteries were most commonly injured (in decreasing order of frequency): superior gluteal, internal pudendal, obturator, and lateral sacral arteries.³ In a separate study, Velmahos et al⁹ found that the most frequently embolized vessels in decreasing frequency were the internal iliac, superior gluteal, internal pudendal, and lateral sacral arteries; these were followed by the obturator and ilio-lumbar arteries, which demonstrated equal frequency of embolization.

TREATMENT MODALITIES

Military Antishock Trousers/Pneumatic Antishock Garment

The use of military antishock trousers/pneumatic antishock garment (MAST/PASG) was once widespread; however, it is not used frequently in the urban, civilian trauma setting where relatively short transit times are the norm. To the authors' knowledge, the largest prospective, randomized study evaluating the use of MAST in the prehospital setting was conducted by Mattox et al,³⁷ who showed no survival benefit in their study of 911 patients. On the contrary, MAST application was associated with increased mortality in various subsets of the study group.³⁷ Other authors also have shown no survival benefit from the use of MAST/PASG.^{38,39} MAST application is also associated with other undesirable qualities, including additional training requirements and expenditures, compromised evaluation of the lower abdomen, pelvis, genitourinary tract, and lower

extremities; restricted access to the femoral vessels; and possible contribution to metabolic acidosis.⁴⁰

External Fixation

External fixation has been advocated as a means for control of pelvic hemorrhage.^{6,7,11,16,41,42} Proposed mechanisms of action are (1) the apposition of fracture fragments, which decreases bone-edge and cancellous bleeding; (2) the stabilization of an early clot, making it less likely to dislodge during transport and/or transfer; and (3) a decrease in the pelvic volume, which increases intrapelvic pressure and thereby affects tamponade of venous and small arterial bleeding. Grimm et al¹⁴ call into question the last of these mechanisms. Their cadaveric study suggests that the true volume of the retroperitoneum is >20 L, and that an expanding retroperitoneal hematoma has little respect for the margins of the bony pelvis as defined by the anatomist. Frequently quoted in the literature, a 3-cm diastasis of the symphysis pubis converts the average pelvic volume from 4000 to 8000 mL, which is 1.5 times the circulating blood volume of a normal-sized adult. Compared with 20,000 mL, either of these volumes is trivial. In addition, Grimm et al¹⁴ also showed little added increase in retroperitoneal pressures after the placement of an external fixation device, and therefore suggest it is unlikely that significant additional tamponade is achieved. Their findings require the questioning of our understanding of the possible mechanisms through which external fixation provides control of hemorrhage in pelvic fractures.

Control of bone-edge bleeding might be the greatest benefit provided. This type of blood loss might be the most significant source of bleeding in pelvic fractures.⁴³ Although the exact mechanisms may not be well understood, pelvic stabilization is effective and should be addressed early in the treatment pathway. The protocol used at our institution addresses the need for timely pelvic stabilization through the early application of a folded sheet wrapped tightly around the patient's pelvis. This is performed in conjunction with tapping the knees and ankles together. These maneuvers provide a rapid, easily applied, temporary solution and are applied prior to transporting the patient to the angiography suite.

Laparotomy and Pelvic Packing

At our institution, laparotomy with pelvic packing can be used to control pelvic hemorrhage and generally is used when the trauma ultrasound (or DPL) is grossly positive, indicating significant hemoperitoneum. Sites of bleeding in the abdomen are addressed and the pelvis is packed with laparotomy pads prior to closure. Anticipation is for early return to the operating room for further intervention and/or repacking.

Celiotomy with Bilateral Hypogastric Artery Ligation

This method of treatment is really a historical footnote because its use is no longer in favor. The nature of the procedure requires violation of the intact peritoneum, which results in release of the natural tamponade provided by this membrane. Furthermore, violation of the retroperitoneum exposes this space to possible sepsis.

Transcatheter Embolotherapy

Transcatheter embolotherapy has become a mainstay of management in the unstable patient with pelvic fractures and is successful in controlling hemorrhage in 85 to 95% of patients who undergo this potentially lifesaving procedure. Prior to discussion of patient selection, access, and techniques, anatomic considerations are reviewed.

ANATOMIC CONSIDERATIONS

The pelvis derives the majority of its blood supply from the paired internal iliac arteries (IIAs). The IIAs (previously named the hypogastric arteries) bifurcate just beyond their origin to form anterior and posterior divisions. Hemorrhage can occur from virtually any arterial branch, and as discussed above, predicting which artery is bleeding based on fracture patterns often is not possible. Other vessels, with origins distinct from the IIA, can also supply the pelvis in the normal setting. Branches from the external iliac (namely the inferior epigastric and deep circumflex iliac arteries) can provide significant blood supply to the pelvis. Superior hemorrhoidal branches, which arise from the inferior mesenteric artery, also supply the deep pelvis.

Vascular anomalies within the pelvis are commonplace. "Normal" pelvic anatomy, in fact, with predictable patterns of branching, is probably seen in the minority of patients. Some common anomalies that may be pertinent to the trauma population undergoing TCE are outlined in Table 3⁴⁴ and Fig. 1.

The most important aspect of pelvic arterial anatomy is the degree of collateralization of vessels noted within the pelvis. Collaterals can form from one side of the pelvis to the other, and can form on the same side of the pelvis from anterior to posterior division branches. It is because of these extensive collateral networks that

Table 3 Pelvic Vascular Anomalies⁴⁴

Common trunk of both divisions (no anterior and posterior divisions), 10%
Obturator artery: arising from inferior epigastric or external iliac artery, 22 and 2%, respectively
Inferior gluteal artery: arising from superior gluteal artery, 25%
Accessory internal pudendal artery, 6 to 10%
Lateral sacral arteries: may be multiple

embolization of large feeding branches to the pelvis is considered safe. Although complete embolization of both IIAs might be associated with an increased risk of pelvic ischemia, no study to date has proven this hypothesis. In clinical practice, the authors believe that it is safe to embolize three of the four trunks of the IIAs without placing the patient at undue risk for pelvic ischemia.

INDICATIONS/PATIENT SELECTION

In the trauma setting, the decision to perform angiography and/or embolotherapy must be weighed against the limitations of the procedures. One clinically significant limitation is the often extensive investment in time. Given that time to definitive treatment is critical, assessment must be made whether the patient can “afford” these possibly lengthy procedures. Location of the angiography suite in close proximity to the trauma bay might aid in ongoing resuscitative measures and crisis management, as well as minimize wasted minutes in transport. Despite the added time needed to perform TCE, rapid transit of suitable patients to the angiography suite might decrease mortality. Agolini et al⁸ found that patients who received embolization within 3 hours of arrival had a significantly greater survival rate than those embolized after 3 hours (14 versus 75%, respectively).

Ideally, the consideration of TCE should be made early and as part of a comprehensive, algorithm-directed treatment strategy. Patients likely to benefit from TCE include, but are not limited to, those with ongoing arterial bleeding despite surgical intervention, bleeding from injuries managed nonoperatively, and those with vascular complications (e.g., pseudoaneurysm or arteriovenous fistula). It should be remembered that although TCE has been shown to be very effective in controlling arterial hemorrhage associated with pelvic fractures, severe fractures occur infrequently, and patients with bleeding from arterial injury represent a small subset of those with hemodynamically significant blood loss.²³ In their recent study, Demetriades et al²³ found that angiography was used in 4.7% of pelvic trauma patients, and only half of these required embolization. Other studies have suggested the requirement for embolization in 10% or more of all patients with pelvic fractures.⁷

ACCESS

If possible, common femoral artery (CFA) access through either groin should be obtained on the side opposite any unilateral pelvic or lower extremity injury. When the nature of the patient's injuries prevents use of this route (e.g., with large groin hematomas or open anterior fractures), the left brachial or axillary approach can be used. Once access is obtained, a 5-Fr sheath should be placed to aid with catheter exchanges, reduce

overall trauma at the arteriotomy site, and provide more secure access. A sheath will also allow for maintained access in the event that a catheter becomes occluded by the embolic agent.

DIAGNOSTIC ANGIOGRAPHY

Certain guidelines should be used when performing diagnostic angiography in trauma patients. First, initial nonselective angiography is performed to visualize arteries adjacent to, but not directly supplying, the injured area. The reason for this is that adjacent vessels, such as the lumbar arteries, might be injured due to proximity injury, either from blunt or penetrating trauma. Second, higher injection rates are typically necessary in the trauma patient population due to the hyperdynamic status of such patients. Cardiac outputs of ≥ 15 L per minute are not unusual in this population. Finally, as with diagnostic angiography performed for other causes of hemorrhage, filming should be performed long into the venous phase to adequately assess sites of contrast extravasation.

In patients with unstable pelvic fractures, an aortogram should be performed expeditiously for visualization of the patient's lower abdomen and pelvis. The catheter is placed just below the renal arteries, and the largest field of view is obtained to visualize lumbar arteries as well as the pelvic supply. Particular attention should be paid to the lumbar arteries, and external iliac branches. In addition, any vascular anomalies should be investigated with the nonselective aortogram. If questions persist, selective angiography of those vessels in question should be obtained.

Occasionally, active contrast extravasation can be observed on the nonselective aortogram (Figs. 1, 2). In this setting, selective catheterization of the bleeding vessel should be performed next. It is important to note, however, that lack of contrast extravasation on the aortogram does not preclude active bleeding; superselective catheterization of both IIAs is always indicated. Superselective catheterization can be performed on both sides with a Cobra-shaped catheter; the curve of the catheter allows easy selection of the contralateral IIA, whereas the ipsilateral IIA can be selected with the same catheter once a Waltman loop is formed. Again, the diagnostic angiogram should be performed forcefully to ensure adequate filling of the artery in a potentially hyperdynamic patient. If reflux into the common iliac artery is not noted, the authors repeat the angiogram at a higher injection rate. When selecting catheters, an end-hole catheter is preferred over a side-hole catheter to aid in more selective deployment of the embolic agent and to decrease the likelihood of nontarget embolization. Rather than exchanging catheters if a bleeding source is found, the authors use an end-hole catheter for the diagnostic angiogram.

Figure 2 A 15-year-old female who presented to the emergency department following a rollover motor vehicle accident. The patient was pinned under the truck for hours (the engine block was cold). As soon as the rescue workers removed the truck, she became hemodynamically unstable. At presentation to the emergency room, she had a systolic blood pressure of < 60 mmHg and a core temperature 27°C. She was immediately transferred to the interventional suite. (A) Digital subtraction aortogram demonstrates marked attenuation of all visualized vessels secondary to the patient's severe hypotension. Also demonstrated is active contrast extravasation arising from the left internal iliac distribution. (B) Selective injection of the left internal iliac artery demonstrates active contrast extravasation from the proximal internal iliac artery. (C) Postembolization angiography following embolization of the proximal internal iliac artery with coils and Gelfoam slurry. There is complete cessation of flow into the internal iliac distribution.

EMBOLIZATION TECHNIQUE: SELECTIVE EMBOLOTHERAPY

In the best of circumstances, a single bleeding source is identified, and selective embolization of the feeding vessel is performed. In some cases, however, multiple bleeding sites are visualized, or the injured vessel cannot be selected due to either technical or time constraints (Fig. 3). In these instances, nonselective ("shotgun") embolotherapy may be necessary (discussed below). This deployment of embolic agents proximal to the bleeding source is reserved only for those cases in which superselective embolization cannot be performed. Nonselective embolotherapy such as this usually is performed only in the pelvis, and generally is considered safe because of the extensive collateralization seen with pelvic vessels, as noted above.

Embolic Agents

The choice of embolic agent depends on the size of the vessel to be embolized, the desired level of permanence, and the familiarity and preferences of the interventionalist.

Posttraumatic bleeding typically involves previously healthy vessels that undergo acute changes from their traumatic insult. Temporizing measures may be all that is required to give the vessel time to heal, or to stabilize the patient for them to go to the operating room.⁴⁵

Autologous blood clot was the first agent used for trauma embolotherapy.¹ Although autologous clot causes a satisfactory occlusion, it tends to be temporary because the injected clot will lyse over time. In many cases, the occluded vessel bleeds again within several hours following embolization with autologous clot. It is for this reason, plus the availability of other commercially available agents, that autologous clot currently is used rarely, if ever.

In the trauma population, because temporary occlusions are usually considered adequate, agents such as Gelfoam pledgets or slurry (Upjohn, Kalamazoo, MI) are preferred (Fig. 4). Gelfoam pledgets and slurry have the added benefits of being readily available and inexpensive, easy to work with, compatible with a microcatheter system, and mechanically obstructive (see Bauer and Ray,⁴⁶ this issue). The Gelfoam slurry, which should approximate the consistency of pudding, is

Figure 3 A 72-year-old male struck by an automobile. The patient was taken directly to the operating room for repair of a grade III liver laceration. During the operation, an expanding pelvic hematoma was noted. After packing of the liver lesion, the patient was transferred to the interventional suite. (Courtesy of Kimi Kondo, D.O.) (A) Selective injection of the right iliac artery demonstrating multiple collections of contrast, indicating active bleeding. (B) Later image following the same injection as shown in (A). Note the multiple sites of contrast extravasation. Because of the multiple distal sites of bleeding, the entire internal iliac distribution was embolized with Gelfoam slurry (shotgun embolization). (C) Final angiogram demonstrating resolution of the multiple bleeding sites.

injected slowly to prevent reflux of agent into more proximal branches, which could result in nontarget embolization. The injection proceeds until there is significant slowing of flow; however, it is important to cease the injection prior to complete cessation of flow because this can result in proximal reflux of agent.⁴⁷ Angiograms performed intermittently during the embolization procedure are mandatory, and the procedure can be stopped once cessation of contrast extravasation is noted.

Coils are a frequently used embolic agent. A property of coils of which one should be constantly aware is that they are dependent on an intact coagulation pathway for effective occlusion. The coils themselves rarely cause vessel occlusion; rather, the fibers on the coil induce a thrombogenic response, leading to clot formation and vessel occlusion. In general, any underlying coagulopathy should be corrected as soon as possible when coils are used.

In addition to the standard push variety, detachable coils have the added benefit of ease of recovery and repositioning in those instances when coil deployment is not as intended (e.g., when the coil is incorrectly sized or positioned). In the trauma population, however, the luxury of the added time needed for detachable coils might not be available.

Coils can provide rapid, permanent occlusion of vessels. As mentioned previously, they are dependent on an intact clotting cascade, and may require reinforcement with additional agents. It should be noted that although a patient might not be coagulopathic at the time of embolization, their clinical course might be uncertain. Coil occlusion can fail if the patient becomes coagulopathic following embolization and the initial clot induced by the coil lyses and a new one cannot be formed. It is the authors' opinion that coils should not be used as the only embolic agent in patients who are receiving many blood products (≥ 8 units packed red blood cells unmatched with fresh frozen plasma), or who are hypothermic (both are causes of coagulopathy). If a coil is used, the authors use it in combination with Gelfoam; the coil can act as a strut to hold the Gelfoam in place, and the Gelfoam causes a mechanical occlusion of the bleeding vessel.

Embolic agents used for other indications, such as tumor embolization, are needed infrequently for pelvic trauma embolization. These agents, such as particles or silk suture, cause a permanent occlusion and tend to cause a more distal occlusion than is necessary. With the exception of using particles to embolize beyond a pseudoaneurysm neck to prevent retrograde filling of the

Figure 4 A 60-year-old male who presented to the angiography suite following a motor vehicle accident. An open-book pelvic fracture was noted. Although the patient was hemodynamically stable, over a 6-hour period the patient's hematocrit level decreased from 36 to 24, required transfusion with five units of packed red blood cells. A large scrotal hematoma was recognized, and the patient was transferred to the interventional radiology suite. (A) Selective right internal iliac angiogram, which demonstrates active contrast extravasation from the right internal pudendal artery. (B) Delayed image from the same injection demonstrating marked contrast pooling at the site of contrast extravasation. (C) Unsubtracted image demonstrating contrast collecting in the patient's scrotum. Note also the marked diastasis of the pubic symphysis resulting from the patient's open-book fracture. (D) Selective injection of the bleeding internal pudendal artery. The artery was embolized with Gelfoam slurry. (E) Final postembolization angiogram demonstrating cessation of bleeding following Gelfoam embolization.

pseudoaneurysm, the authors do not find it necessary to use particles or suture material during pelvic trauma embolization procedures. Other embolic materials, such as liquid agents, have little to no role in the acutely traumatized patient.

Shotgun Embolotherapy

If subselective TCE cannot be performed, and the decision has been made to deploy the embolic agent

more proximally within the internal iliac vessels, it is important to place the catheter near the origin of the vessel to embolize effectively the ilio lumbar and lateral sacral branches, both arising from the proximal posterior division of the IIA. Advancing the catheter too far could result in failure to embolize the bleeding branches.⁴² Although not the first-line approach in TCE, temporary arterial embolization of the bilateral internal iliac arteries (TAEBIIA) has been advocated as an effective, rapid, and safe alternative.¹⁰ Velmahos et al¹⁰ found that

angiographic and clinical control of bleeding was obtained in 27 of 30 patients (90%) undergoing TAE/BIIA; with inclusion of two additional patients who were taken for repeat embolization, the success rate increased to 97%. One death (of 10) occurred after failure of TAE/BIIA to control ongoing hemorrhage; the remainder of deaths occurred because of associated injuries.¹⁰ Although such shotgun embolizations are not optimal, given the option of no intervention, they may still prove to be the most appropriate therapy for these very unstable patients (Fig. 3).

Interventions Other Than Embolization

In the setting of an unstable patient with pelvic fractures, interventions other than embolization are often possible. In particular, temporary balloon occlusion of a massively bleeding trunk, particularly one in which embolotherapy has failed or cannot be performed because of technical reasons, should be considered. By temporarily tamponading the bleeding vessel, the severely injured patient might be able to go to the operating room for a definitive procedure. This setting would be very uncommon in the pelvic trauma population, however. Any vessel that is bleeding at such a rate would probably force the surgeons to the operating room before the patient even presented to the angiography suite.

Stents and covered stents have been used effectively in treating acutely or chronically traumatized vessels.^{48,49} Although typically used in larger vessels such as the aorta, stents have been used effectively to treat traumatic injury to cervical vessels as well.^{50,51} With the present state of technology, however, deployment of stents, covered stents, or stent grafts would be exceedingly difficult given the tortuosity of the pelvic vasculature. The authors know of no published report documenting successful use of such devices in the pelvic trauma population.

COMPLICATIONS AND LIMITATIONS

The use of TCE in the management of hemodynamically unstable patients with pelvic fractures is not without inherent risks. Reported short- and long-term complications include nontarget embolization of lower extremity vessels, rhabdomyolysis that may be worsened by ischemia, acute renal failure, colonic and ureteric infarctions, perineal wound sepsis, sacral nerve injury, sexual dysfunction (impotence), rectal dysfunction (difficulty defecating), avascular necrosis of the femoral head, and death.^{12,52-55} The direct impact that TCE has on mortality in hemodynamically unstable patients with pelvic fracture is difficult to assess given the relatively high mortality these patients experience using any treatment algorithm. The lack of prospective, randomized data to determine the optimal timing of TCE

within various treatment algorithms is unfortunate; however, designing a study that would withhold TCE from some patients would be unethical and likely will never be performed.³⁴ Even those authors who favor TCE as a last resort recognize that patients requiring angiography have much higher injury severity scores than hemodynamically stable patients.¹² When TCE is employed as a heroic measure after all other modalities have failed, it is not surprising that mortality remains high. In the reviewed literature, mortality in association with TCE ranges from 17 to 89%,^{5,6,8,12,16,17,25} with a mean of 43%.¹² Further complicating matters, the patients most likely to benefit from TCE are those with significant ongoing arterial hemorrhage. Ideally, this condition would be identified and corrected as soon as possible. Unfortunately, this diagnosis might be reached by the process of elimination, which can result in a significant delay in reaching the angiography suite, thus subjecting the patient to increased blood loss and resulting in increased mortality.⁴⁵

CONCLUSION

Hemorrhage from pelvic fractures remains a significant source of mortality in the polytraumatized patient. The potential lethal character of intrapelvic hemorrhage demands urgent diagnosis and treatment. The use of an algorithm-directed, expedient protocol will increase patient survival. Within these algorithms, angiography should be performed as soon as possible after excluding significant intra-abdominal and thoracic sources of blood loss. Angiography is accurate in identifying sites of arterial hemorrhage, and TCE is highly effective in providing definitive control at these sites. Rapid recognition and correction of underlying coagulopathy is helpful in maximizing the efficacy of TCE.

REFERENCES

1. Margolies MN, Ring EJ, Waltman AC, Kerr WS, Baum S. Arteriography in the management of hemorrhage from pelvic fractures. *N Engl J Med* 1972;287:317-321
2. Flint LM, Brown A, Richardson JD, Polk HC. Definitive control of bleeding from severe pelvic fractures. *Ann Surg* 1979;189:709-714
3. Kam J, Jackson H, Ben-Menachem Y. Vascular injuries in blunt pelvic trauma. *Radiol Clin North Am* 1981;19:171-186
4. Mucha P, Farnell MB. Analysis of pelvic fracture management. *J Trauma* 1984;24:379-385
5. Panetta T, Sclafani SJ, Goldstein AS, Phillips TF, Shaftan GW. Percutaneous transcatheter embolization for massive bleeding from pelvic fractures. *J Trauma* 1985;25:1021-1028
6. Moreno C, Moore EE, Rosenberger A, Cleveland HC. Hemorrhage associated with major pelvic fracture: a multi-specialty challenge. *J Trauma* 1986;26:987-993
7. Ben-Menachem Y, Coldwell DM, Young JW, Burgess AR. Hemorrhage associated with pelvic fractures: causes,

- diagnosis, and emergent management. *AJR Am J Roentgenol* 1991;157:1005-1014
8. Agolini SF, Shah K, Jagge J, Newcomb J, Rhodes M, Reed JF. Arterial embolization is a rapid and effective technique for controlling pelvic fracture hemorrhage. *J Trauma* 1997;43:395-399
9. Velmahos GC, Chahwan S, Falabella A, Hanks SE, Demetriades D. Angiographic embolization for intraperitoneal and retroperitoneal injuries. *World J Surg* 2000;24:539-545
10. Velmahos GC, Chahwan S, Hanks SE, et al. Angiographic embolization of bilateral internal iliac arteries to control life-threatening hemorrhage after blunt trauma to the pelvis. *Am Surg* 2000;66:858-862
11. Biffl WL, Smith WR, Moore EE, et al. Evolution of a multidisciplinary clinical pathway for the management of unstable patients with pelvic fractures. *Ann Surg* 2001;233:843-850
12. Cook RE, Keating JF, Gillespie I. The role of angiography in the management of haemorrhage from major fractures of the pelvis. *J Bone Joint Surg (Br)* 2002;84-B:178-182
13. McCoy GF, Johnstone RA, Kenright J. Biomechanical aspects of pelvic and hip injuries in road traffic accidents. *J Orthop Trauma* 1989;3:118-123
14. Grimm MR, Vrahas MS, Thomas KA. Pressure-volume characteristics of the intact and disrupted pelvic retroperitoneum. *J Trauma* 1998;44:454-459
15. Rothenberger DA, Fischer RP, Strate RG, Velasco R, Perry JF. The mortality associated with pelvic fractures. *Surgery* 1978;84:356-359
16. Evers BM, Cryer HM, Miller FB. Pelvic fractures: priorities in management. *Arch Surg* 1989;124:422-424
17. Poole GV, Ward EF, Muakkassa FF, Hsu HSH, Griswold JA, Rhodes RS. Pelvic fracture from major blunt trauma: outcome is determined by associated injuries. *Ann Surg* 1991;213:532-538
18. Gruen GS, Leit ME, Gruen RJ, Peitzman AB. The acute management of hemodynamically unstable multiple trauma patients with pelvic ring fractures. *J Trauma* 1994;36:706-711
19. Poole GV, Ward EF. Causes of mortality in patients with pelvic fractures. *Orthopedics* 1994;17:691-696
20. Muir L, Boot D, Gorman DF, Teanby DN. The epidemiology of pelvic fractures in the Mersey Region. *Injury* 1996;27:199-204
21. Jones AL, Powell JN, Kellam JF, McCormack RG, Dust W, Wimmer P. Trauma to the adult pelvis and hip. *Orthop Clin North Am* 1997;28:345-350
22. Parreira JG, Cimbra R, Rasslan S, Oliveira A, Fregoneze M, Mercadante M. The role of associated injuries on outcome of blunt trauma patients sustaining pelvic fractures. *Injury* 2000;31:677-682
23. Demetriades D, Karaiskakis M, Toutouzas K, Alo K, Velmahos G, Chan L. Pelvic fractures: epidemiology and predictors of associated abdominal injuries and outcomes. *J Am Coll Surg* 2002;195:1-10
24. Eastridge BJ, Burgess AR. Pedestrian pelvic fractures: 5-year experience of a major urban trauma center. *J Trauma* 1997;42:695-700
25. Gilliland MD, Ward RE, Barton RM, Miller PW, Duke JH. Factors affecting mortality in pelvic fractures. *J Trauma* 1982;22:691-693
26. Adams JE, Davis GG, Heidepriem RW, Alonso JE, Alexander CB. Analysis of the incidence of pelvic trauma in fatal automobile accidents. *Am J Forensic Med* 2002;23:132-136
27. Reed RA, Teitelbaum GP, Katz MD, Pentecost MJ. Early management of the trauma patient with pelvic fracture: a medical perspective. *Semin Interv Radiol* 1992;9:1-3
28. Baum S. *Abrams' Angiography: Vascular and Interventional Radiology*, 4th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 1997:1648-1650
29. Young JW, Burgess AR, Brumback RJ, Poka A. Pelvic fractures: value of plain radiography in early assessment and management. *Radiology* 1986;160:445-451
30. Moore KL. *Clinically Oriented Anatomy*, 3rd ed. Baltimore: Williams and Wilkins; 1992:243-252
31. Tile M. *Fractures of the Pelvis and Acetabulum*. Baltimore: William and Wilkins, 1984
32. Young JW, Resnik CS. Fracture of the pelvis: current concepts of classification. *AJR Am J Roentgenol* 1990;155:1169-1175
33. Burgess AR, Eastridge BJ, Young JW, et al. Pelvic ring disruptions: effective classification system and treatment protocols. *J Trauma* 1990;30:848-856
34. Hamill J, Holden A, Paice R, Civil I. Pelvic fracture pattern predicts pelvic arterial haemorrhage. *Aust N Z J Surg* 2000;70:338-343
35. Henry SM, Pollack AN, Jones AL, Boswell S, Scalea TM. Pelvic fracture in geriatric patients: a distinct clinical entity. *J Trauma* 2002;53:15-20
36. Pereira SJ, O'Brien DP, Luchette FA, et al. Dynamic helical computed tomography scan accurately detects hemorrhage in patients with pelvic fracture. *Surgery* 2000;128:678-685
37. Mattox KL, Bickell W, Pepe PE, Burch J, Feliciano D. Prospective MAST study in 911 patients. *J Trauma* 1989;29:1104-1112
38. Chang FC, Harrison PB, Beech RR, Helmer SD. PASG: does it help in the management of traumatic shock? *J Trauma* 1995;39:453-456
39. Mackersie RC, Christensen JM, Lewis FR. The prehospital use of external counterpressure: does MAST make a difference? *J Trauma* 1984;24:882-888
40. Schneider PA, Mitchell JM, Allison EJ. The use of military antishock trousers in trauma: a reevaluation. *J Emerg Med* 1989;7:497-500
41. Agnew SG. Hemodynamically unstable pelvic fractures. *Orthop Clin North Am* 1994;25:715-721
42. Sclafani SJ, Becker JA. Traumatic presacral hemorrhage: angiographic diagnosis and therapy. *AJR Am J Roentgenol* 1982;138:123-126
43. Huittinen VM, Slatis P. Postmortem angiography and dissection of the hypogastric artery in pelvic fractures. *Surgery* 1973;73:454-462
44. Kadir S. *Diagnostic Angiography*, 2nd ed. Philadelphia: W.B. Saunders; 1991:98-100
45. Ray CE, Waltman AC. General principles of embolization and chemoembolization. In: Bakal CW, Silberzweig JE, Cynamon J, et al, eds. *Vascular and Interventional Radiology: Principles and Practice*. New York: Thieme; 2002:93-94
46. Bauer JR, Ray CE Jr. Transcatheter arterial embolization in the trauma patient: a review. *Semin Interv Radiol* 2004;21:11-22
47. Kidney DD. The endovascular approach to trauma. In: Dyet JF, Ettles DF, Nicholson AA, et al, eds. *Textbook of Endovascular Procedures*. Philadelphia: Churchill Livingstone; 2000:314

48. Fattori R, Napoli G, Lovato L, et al. Indications for, timing of, and results of catheter-based treatment of traumatic injury to the aorta. *AJR Am J Roentgenol* 2002;179:603–609
49. Hoffer EK, Karmy-Jones R, Bloch RD, et al. Treatment of acute thoracic aortic injury with commercially available abdominal aortic stent-grafts. *J Vasc Interv Radiol* 2002;13:1037–1041
50. Ray CE, Biffi WL. Neurovascular interventions in the trauma patient. *Semin Interv Radiol* 2002;19:377–387
51. Duke BJ, Ryu RK, Coldwell DM, Brega KE. Treatment of blunt injury to the carotid artery by using endovascular stents: an early experience. *J Neurosurg* 1997;87:825–829
52. Perez JV, Hughes TM, Bowers K. Angiographic embolization in pelvic fracture. *Injury* 1998;29:187–191
53. Rose SC, Moore EE. Emergency trauma angiography: accuracy, safety, and pitfalls. *AJR Am J Roentgenol* 1987;148:1243–1246
54. Hare WS, Holland CJ. Paresis following internal iliac artery embolization. *Radiology* 1983;146:47–51
55. Vassiliu P, Sava J, Toutouzas KG, Velmahos GC. Is contrast as bad as we think? Renal function after angiographic embolization of injured patients. *J Am Coll Surg* 2002;194:142–146