Published in final edited form as:

J Org Chem. 2010 November 19; 75(22): 7636–7643. doi:10.1021/jo1015135.

Reactivity and Selectivity in Hydrovinylation of Strained Alkenes

Wang Liu and T. V. RajanBabu*

Department of Chemistry, The Ohio State University, 100 W. 18th Ave., Columbus, Ohio 43210, USA

Abstract

Scope of Ni(II)-catalyzed hydrovinylation has been extended to strained alkenes such as heterobicyclic- [2.2.1]heptanes and cylobutenes. Reactions involving the heterobicyclic compounds are rare examples for this class of compounds where the metal-catalyzed C-C bond-forming reactions proceed *without* a concomitant ring-opening process. While the enantioselectivity in these systems remains modest, hydrovinylation of *endo*-5,6--bis-benzyloxymethylbicyclo[2.2.1]hept-2-ene gives excellent yield (>90%) of the product with one of the highest enantioselectivities (95–99 %ee) reported for a C-C bondforming reaction of norbornenes.

Introduction

Heterodimerization of alkenes is a reaction with a huge potential for the synthesis of valuable intermediates since the starting materials are often readily available, or can easily be synthesized.^{1,2} Advantages of alkenes over other conventional carbon feedstocks such as CO or HCN include their lack of toxicity and ease of handling while possessing sufficient reactivity to permit activation by transition metal complexes. In addition, depending on the substitution pattern, an alkene could be prochiral, and thus can serve as a cheap source for enantiomerically pure intermediates. However, since the two starting materials and the expected product(s) in a dimerization necessarily carry the same functional group, viz., an alkene, finding successful reaction conditions without concomitant side-reactions such as homodimerizations, oligomerizations and isomerizations is more challenging. The success of the reaction depends on judicious choice of two alkenes, where there are significant differences in their reactivities. Such differences might result from either electronic or steric reasons. Proper choice of a catalyst can augment such differences, and a number of these dimerization reactions have been developed.³ Hydrovinylation (addition of ethylene) of alkenes (eq 1) is one such reaction where significant progress has been achieved in several areas including the development of enantioselective processes. Codimerization of ethylene with more reactive partners like vinylarenes and 1,3-dienes have been carried out with excellent overall selectivity, and advances in broadening the scope of this reaction with respect to substrates and catalysts continue unabated.⁴

rajanbabu.1@osu.edu.

SUPPORTING INFORMATION. Spectroscopic and chromatographic data for characterization of all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Results and Discussion

Norbornene represents a class of substrates where the viability of the reaction depends presumably on the enhanced reactivity of the strained bicyclic system as compared to ethylene and the dimerization product (eq 2). Indeed hydrovinylation of norbornene was among the first metal-catalyzed asymmetric carbon-carbon bond-forming reactions ever reported,⁵ even though the enantioselectivity was unacceptable by current standards. Other reports of codimerization of norbornene with ethylene include the use of $[\text{Ni}(2,4,6\text{-Me}_3\text{C}_6\text{H}_2)(\text{CH}_3\text{CN})(\text{phosphane})]^+ [\text{BF}_4]^-$,⁶ $(\text{PCy}_3)_2(\text{CO})\text{RuHCl}/\text{HBF}_4\cdot\text{Et}_2\text{O}$,⁷ and $\text{Co}(\text{pyridineimine})\text{Cl}_2/\text{MAO}$.⁸ In 2003 we reported a remarkable ligand effects on the course of the Ni-catalyzed hydrovinylation of norbornene (Scheme 1). It was shown that under our then newly developed reaction conditions a ligand with a smaller cone angle (Ph_3P , 145°) gave a 2:1 adduct (norbornene:ethylene) whereas a larger ligand (Cy_3P , cone angle 180°) gave a 1:1 adduct.⁹ Among several chiral ligands examined, a phosphoramidite ligand (**L4**, Figure 1) derived from 1,1'-binaphthol gave quantitative yield, giving the 1:1 adduct (**1**) in ~80% ee. Even though this represents one of the highest enantioselectivity reported for a carbon-carbon bond-forming reaction of norbornene,¹⁰ the generality of this reaction, or the broader question of whether reactivity differences brought about by strain can be used to effect a selective heterodimerization has not been addressed. In this paper we disclose the first experiments that deal with these aspect of hydrovinylation.

Hydrovinylation of norbornene derivatives

Our studies started with a detailed examination of the hydrovinylation of *endo*-5,6-bis-5,6(benzyloxymethyl)bicyclo[2.2.1]hept-2-ene (**3**) using the ligands **L1-L6**. This substrate was chosen as a prototypical bicyclo[2.2.1]-alkene since the enantiomers of the product can be detected by UV absorption, and, thus directly analyzed on a chiral stationary phase HPLC column. The choice of ligands for this study is based on several previous observations from which we concluded that hemilabile ligands, when used in conjunction with a highly dissociated counter ion (e.g. 3,5- $[(\text{CF}_3)_2\text{C}_6\text{H}_3]_4\text{B}^-$, BARF⁻) gave the highest selectivity in the reactions of vinylarenes^{11,12} and 1,3-dienes.^{13,14} It was also known that in ligands **L1-L3** the location of the 'hemilabile' oxygen in relation to the phosphorus atom is crucial for obtaining high regio- and stereoselectivity for specific classes of substrates. Thus we have shown that ligand **L1** is the most suitable one for the hydrovinylation of certain classes of 1,3-dienes,¹³ ligand **L2** is best for vinylarenes¹⁵ and ligand **L3** gave low selectivities for all classes of substrates. The phosphoramidites, popularly known as the Feringa ligands, are by far the most successful ligands for this reaction.¹⁶⁻²¹ In our work the three ligands shown **L4-L6** have been found to be broadly applicable for several of the hydrovinylation reactions.²¹

(2)

(3)

The Ni(II)-catalyzed hydrovinylation of **3** with the ligands carrying a hemilabile benzyloxy substituent (**L1-L3**) parallels our observations in the related reaction with vinylarenes (Table 1). Nickel complex formed from the ligand **L1**, allyl nickel bromide dimer and Na BARF is competent to effect the hydrovinylation of this substrate at low temperature (entry 1), but at higher temperatures, extensive isomerization of the product (to give a mixture of **4** and **5**) is observed (entry 2). Ligands **L2** and **L3**, on the other hand, did not isomerizes the initially formed product up to several hours. We had previously found that **L2** was one of the few ligands capable of effecting hydrovinylation of several vinylarenes at room temperature with *no isomerization of the initially formed 3-aryl-1-butene*.¹⁵ Compound **5** is formed in varying amounts (0–10%) depending on the catalyst and reaction conditions (and its structure is presumed on the basis of ¹H NMR which shows a distinct olefinic H at δ 5.04 (q, J = 6 Hz) and C_{sp2}-CH₃ signals at δ 1.81 (d J = 6 Hz). As with the other substrates, finely tuned phosphoramidites are the best ligands for the hydrovinylation of this substrate. Thus ligand **L4** and its enantiomer, *ent*-**L4**, and a simplified analog containing the biphenyl core (**L6**) gave the best overall yield and selectivity (>90% yield, 93–99% ee) for this reaction (entries 5, 6 and 8). This result represents the *highest enantioselectivity ever observed* for an asymmetric catalyzed C-C bond-forming reaction of a norbornene derivative. Surprisingly, the ligand **L5**, which gave the highest yields and enantioselectivities (up to 99% yield and >95% ee) for a wide spectrum of vinylarenes,²¹ was only moderately selective (66% ee) for hydrovinylation of **3**, even though the yield of the reaction was excellent (entry 7). The enantioselectivities of the products were determined by chiral stationary phase HPLC separation of the primary products on Chiracel-AD-H column using hexane and isopropanol (99.6:0.4) as the mobile phase. The ratios of enantiomers were further confirmed by conversion of the dibenzylether **4** into a THF derivative **6** (eq 3). The enantiomers of the THF derivative show baseline resolution in gas chromatography on a Cyclodex B column. A racemic authentic sample of **6** was prepared via hydrovinylation using ligand **L3**.

Hydrovinylation of Azabicyclo[2.2.1]alkenes

Even though carbametallation and related reactions of norbornene and similar [2.2.1]-bicyclic molecules are well documented, reactions of the corresponding heterocyclic analogs where the bicyclic motif is preserved are rare.^{22–24} Such metalation reactions involving Ni, Pd and Rh intermediates most often lead to ring opening reactions.^{24–31} In earlier work we had noticed that substrates carrying hetero-atoms, especially nitrogen, are slow to undergo hydrovinylation under moderate conditions.¹⁹ In light of this observation it is surprising that the azabicyclic alkene **7** undergoes hydrovinylation in very good yields, even though, not unexpectedly, the product is obtained as a mixture of two regio-isomers (Scheme 2, Table 2). For this reaction, the achiral ligands **L1-L3** are much less effective (entries 1–3) compared to the phosphoramidite ligands **L4-L6**, the Ni-complexes of the former requiring room temperature to get reasonable yields. Yields up to 89% are obtained using the latter class of ligands at –47 °C (entries 4–6). In these hydrovinylation, ligand **L1** is the least reactive and at temperatures where full conversion is observed (–10 °C), the primary products are completely isomerized to the ethylidene derivatives **10–13**. The compounds **12** and **13** were not separable by column chromatography, and were identified as the ketone **14**,

prepared by Ru-mediated oxidation of the alkenes **12** and **13**. Note that the enantioselectivity of these reactions are among the lowest we have seen even with the highly successful phosphoramidite ligands (entries 4–6, column 6).

Two other [2.2.1]-heterobicyclic molecules that undergo hydrovinylation are shown in eq 4 and 5. In general, these substrates react only sluggishly with most ligands, and, invariably higher temperatures are required to get acceptable conversion to the products. Curiously for both these substrates ligand **L1** works best. Attempts to achieve complete conversion leads to extensive isomerization of the initial products and other side reactions. The hydrovinylation reaction of **17** using a Ni(II)-**L4** complex gave a poor yield of the product **18**, yet with an unusually high enantioselectivity (>87% ee) as determined by chiral stationary phase GC, where base-line separation of the enantiomers was observed. The configuration of the major product has not been confirmed independently, but the proposed structure is based on the previous results from hydrovinylation of norbornene.⁹

(4)

(5)

Finally, the bicyclic heterocycles shown in Figure 2 failed to undergo the Ni-catalyzed hydrovinylation under a variety of conditions. The exact reason why these substrates fail to undergo the reaction is not obvious. One might speculate that the first two substrates are Lewis basic and probably forms stable Ni(II)-intermediates. The second and third substrates, because of the disposition of the substituents on N (over the lone double bond) is sterically encumbered. Recall that a reactive Ni(II)intermediate, possibly a hydride, has to initially coordinate with the double bond before the reaction can take place.

Hydrovinylation of a Cyclobutene (**19**)

Except for a few notable examples that involve metathesis reactions (Ru)^{32–36} and rearrangements (Rh),³⁷ activation of the strained double bond in cyclobutenes for metal-catalyzed carbon-carbon bond forming reactions have not been explored.³⁸ We find that the cyclobutene double bond is sufficiently reactive to take part in a heterodimerization reaction with ethylene even at low temperatures. Thus under the standard reaction conditions described earlier, the compound **19** undergoes hydrovinylation reactions giving the expected product in moderate yields (Scheme 3 and Table 3). Attempts to isolate of **20** from a mixture **19** and **20** of were unsuccessful, and this mixture was subjected to hydroboration in order to

isolate the hydrovinylation product as a primary alcohol derivative **21**. As with other substrates, the phosphoramidites, in particular, the ligand **L6** gave the best selectivities for this substrate. While the absolute configuration of the product has not been determined, the relative configuration of **20** was established by nOe measurements on the hydroboration product **21**. We have not optimized these reactions except for examining the viability of these ligands.

Conclusions

An *endo*-5,6-bis(benzyloxymethyl)bicyclo[2.2.1]hept-2-ene derivative undergoes Ni(II)-catalyzed asymmetric hydrovinylation giving the highest enantioselectivities (~95% ee) reported for a C-C bond-forming reaction of norbornene derivatives. Other results reported in this paper clearly demonstrate that the scope of Ni(II)-catalyzed hydrovinylation extends to strained alkenes such as heterobicyclic-[2.2.1]heptenes and cyclobutenes. Reactions involving the hetero-bicyclic compounds are rare examples for this class of compounds where the metal-catalyzed C-C bond-forming reactions proceed *without* a concomitant ring-opening process. While the enantioselectivity in these systems remains modest, we hope that these results will stimulate further work in this area, especially since the products of this reaction can potentially be transformed into highly substituted cyclohexane derivatives.

Experimental Section

General methods

Reactions requiring air-sensitive manipulations were conducted under an inert atmosphere of nitrogen by Schlenk techniques or with the aid of a Vacuum Atmospheres glovebox. Methylene chloride and 1,2-dichloroethane were distilled from calcium hydride under a dry atmosphere and stored over molecular sieves. Tetrahydrofuran was distilled under nitrogen from sodium/benzophenone ketyl. The ligands **L1-L3**³⁹, **L4-L6**⁴⁰ and Na⁺{[3,5-(CF₃)₂C₆H₃]₄B}⁻ (NaBARF)⁴¹ were prepared according to the methods described in the literature. Ethylene (99.5%) was purchased from Matheson and passed through Drierite before use. Flash column chromatography was carried out on silica gel 40 (). Enantiomeric excesses of volatile chiral compounds were determined by gas chromatographic analysis on chiral stationary phase GC columns (Cyclodex B, 25 m × 0.25 mm ID, 0.12 mm film thickness or Cyclosil 30 m X 0.25 mm ID, 0.25 μm thickness). Enantiomeric excesses of other compounds were determined by HPLC using a Chiralcel OJ-H or a Chiralcel AD-H column with hexane/isopropanol as solvents. Conditions are described under specific compounds. Optical rotations were recorded at the sodium D line in chloroform.

Synthesis of *endo*-5,6-bis(benzyloxymethyl)bicyclo[2.2.1]hept-2-ene (**3**)

To a solution of 2,3-*bis*(hydroxymethyl)bicyclo[2.2.1]hept-2-ene⁴² (1.79 g, 11.6 mmol) in THF (30 mL) was added NaH (1.39 g, 60 wt % in mineral oil, 34.8 mmol) in one portion at 0°C under argon. The resulting suspension was stirred at 0°C for 60 min and then benzyl bromide (4.4 mL, 37 mmol) was added dropwise at 0°C. The mixture was allowed to warm to room temperature and stirred for 18 h. Water was added carefully to quench the reaction, and the mixture was extracted with ether and the organic layers were combined, washed with brine, dried and concentrated. The resulting residue was purified by column chromatography on silica gel (eluting with hexanes/ethyl acetate = 30/1) to get 3.4 g (88%) of **3** as an oil: ¹H NMR (500 MHz, CDCl₃): δ 1.24 (d, *J* = 8.2 Hz, 1 H), 1.39 (d, *J* = 8.2 Hz, 1 H), 2.46 – 2.47 (m, 2 H), 2.89 (s, 2 H), 2.95 – 2.99 (m, 2 H), 3.21 – 3.24 (m, 2 H), 4.32 and 4.38 (AB quartet, *J* = 12.0 Hz, 4 H), 5.96 (s, 2 H), 7.17 – 7.27 (m, 10 H), ¹³C NMR (125 MHz, CDCl₃): δ 41.6, 45.6, 49.0, 70.5, 73.0, 127.5, 127.6, 128.3, 135.3, 138.7; IR (neat) cm⁻¹: 2920, 2855, 1496, 1454, 1365, 1095; HRMS (ESI) Calcd for C₂₃H₂₆O₂Na ([M + Na]⁺) 357.1830, observed 357.1832.

Typical procedure for catalytic hydrovinylation (Table 1, entry 8)

The pre-catalyst was prepared as follows in a glovebox: To **di-η-allyl-di-μ-bromonickel(II)** (1.0 mg, 0.00278 mmol) in CH₂Cl₂ (1 mL) was added a solution of ligand **L6** (2.5 mg, 0.0057 mmol) in CH₂Cl₂ (1 mL) at ambient temperature. The resulting solution was added to a suspension of Na BARF (5.0 mg, 0.00564 mmol) in dichloromethane (1 mL) and the mixture was stirred at ambient temperature in a septum-sealed 25 mL round bottom flask for 10 min affording a dark brown solution containing a small amount of fine particulate (NaBr). The flask was removed from the glovebox and was then cooled to -78 °C (acetone-dry ice bath), creating a small vacuum. Dry ethylene (passed through a 0.5" × 4" column of Drierite®) was introduced *via* needle through the serum stopper and the vessel atmosphere was slowly evacuated 3 times with a 60 mL syringe. At 78 °C, a solution of **3** (60 mg, 0.18 mmol) in 1 mL dry CH₂Cl₂ is introduced dropwise into the solution of the pre-catalyst over a one minute period via a syringe. The vessel was then maintained at -78 °C for a period of 6 h. At the end of this period the ethylene line is removed and the reaction was quenched by addition of a saturated aqueous NH₄Cl solution. The product was extracted with ether. The organic layers are combined, dried over MgSO₄ and concentrated to get an yellow oil. The resulting residue was purified by column chromatography on silica gel (eluting with hexanes/ethyl acetate = 40/1) to get 60 mg (92%) of **4** as an oil.

Hydrovinylation of Substrate 3, Products 4 and 5

¹H NMR (500 MHz, CDCl₃): δ 1.08–1.12 (m, 1 H), 1.22 (d, *J* = 9.8 Hz, 1 H), 1.42 (d, *J* = 9.8 Hz, 1 H), 1.59–1.61 (m, 1 H), 2.14 (s, 1 H), 2.18 (m, 3 H), 2.25 (s, 1 H), 3.30 (t, *J* = 8.7 Hz, 1 H), 3.37 (t, *J* = 8.7 Hz, 1 H), 3.43–3.46 (m, 1 H), 3.48–3.51 (m, 1 H), 4.35–4.44 (m, 4 H), 4.80–4.85 (m, 2 H), 5.65–5.72 (m, 1 H), 7.17–7.27 (m, 10 H); ¹³C NMR (125 MHz, CDCl₃): δ 30.1, 36.7, 37.6, 39.5, 40.2, 40.7, 45.6, 67.8, 68.5, 73.1, 112.0, 127.5, 127.6, 127.6, 128.3, 138.6, 138.6, 144.0; IR (neat) cm⁻¹: 2954, 2871, 1634, 1496, 1454, 1366, 1097; HRMS (ESI) Calcd for C₂₅H₃₀O₂Na ([M + Na]⁺) 385.2144, observed 385.2139; [α]_D²² = -9.1 (*c* 0.33, CHCl₃) (**L6** at -35 °C, 82% ee by HPLC); [α]_D²² = +13.1 (*c* 0.42, CHCl₃) (*ent*-**L4** at -78 °C, 95% ee by HPLC; > 99% by GC of the corresponding THF derivative **6**, see next experiment); HPLC conditions and retention times (Chiracel AD-H): solvent hexanes:isopropanol 99.6:0.4; flow rate 0.3 mL/min, retention times (min): 25.53 (+)-isomer, 29.54 (-)-isomer. The retention time for starting material: 32.30 min. (see attached chromatograms). Chiral stationary phase GC for **6** (Cyclodex B column, programmed run: 10 min at 80 °C, 1 °C per min to 110 °C, 60 min at 110 °C; retention time: 43.91 min, (-)-isomer: >99% (only one isomer seen in the gas chromatogram). Retention times of authentic mixture (-)-isomer: 43.75 min; (+)-isomer: 44.45 min.

Conversion of 4 into the THF Derivative 6

To a solution of **4** (131 mg, 0.36 mmol, prepared using **L6** at -78 °C) in CH₂Cl₂ (20 mL) was added a hexane solution of BCl₃ (3.6 mL, 1 mol L⁻¹) dropwise at -20 °C under nitrogen. The resulting solution was stirred for 2 h and then quenched with MeOH (10 mL) at the same temperature. The mixture is stirred for 40 min. The solvent was then removed under reduced pressure. The resulting residue was purified by column chromatography on silica gel (eluting with pentane/ethyl ether = 50/1) to get 59 mg (90%) of **6** as an oil. ¹H NMR (500 MHz, CDCl₃): δ 1.05–1.09 (m, 1 H), 1.31 (d, *J* = 10 Hz, 1 H), 1.52 (d, *J* = 10 Hz, 1 H), 1.76–1.81 (m, 1 H), 1.99–2.00 (m, 1 H), 2.13–2.15 (m, 1 H), 2.38–2.53 (m, 3 H), 3.31 (dd, *J* = 9.6 Hz, 6.8 Hz, 2 H), 3.80 (d, *J* = 9.5 Hz, 1 H), 3.80 (d, *J* = 9.5 Hz, 1 H), 4.78 (d, *J* = 10.4 Hz, 1 H), 4.85 (d, *J* = 17.1 Hz, 1 H), 5.65–5.72 (m, 1 H); ¹³C NMR (125 MHz, CDCl₃): δ 30.6, 38.5, 39.7, 40.7, 45.4, 46.1, 46.2, 68.6, 69.0, 111.8, 144.1; IR (neat) cm⁻¹: 2945, 2848, 1636, 1090, 998, 908. HRMS (ESI) Calcd for C₁₁H₁₆ONa ([M + Na]⁺) 187.1099, observed 187.1104. [α]_D²² = -31.7 (*c* 0.54, CHCl₃). GC (methylsilicone)

conditions: 110°C (isothermal), retention time: 29.88 min. GC Cyclodex B column, retention times (min) 43.81 min, confirmed by comparison to authentic product (see next experiment)

Synthesis of Racemic **6** via Hydrovinylation of **4** followed by THF formation

A racemic sample was prepared by hydrovinylation of **4** using the ligand **L3** (4 mol% catalyst, 23 °C, 2 h, 72% yield) followed by the THF formation. GC retention times (min): 43.76 min (~ 50%), 44.45 min (~50%).

An alternate synthesis of THF derivative **6** via hydrovinylation of cyclopentadiene/maleic anhydride adduct

The general hydrovinylation procedure was employed for the anhydride adduct (50 mg, 0.30 mmol) with ligand **L4** (13.2 mg, 0.024 mmol, 8 mol% catalyst) at 23 °C over 13 h. The crude product was passed through a short pad of silica gel and redissolved in THF (10 mL) at 0°C, and LiAlH₄ (40 mg, 1.05 mmol) was added to this solution. The reaction mixture was warmed to room temperature and stirred for 14 h and then quenched with saturated NH₄Cl. The aqueous layer was extracted with ether four times. The organic layer was washed with brine, dried over MgSO₄, and concentrated. The residue was purified by column chromatography (eluting with hexanes/ethyl acetate = 1.0/1.2) to get 24 mg of 1.0:2.8 mixture of two diols, one corresponding to the unreacted starting material and one to the hydrovinylation product, in 13% and 36% yields respectively. To an ice-cooled solution of the mixture of diols (27 mg) and pyridine (27.6 mg, 0.35 mmol) in CH₂Cl₂ (2 mL) was added *p*-TsCl (33.3 mg, 0.17 mmol). The reaction solution was stirred at room temperature for 21 h before addition of H₂O. The organic solution was separated, washed with 1 M aqueous HCl solution, saturated aqueous NaHCO₃ solution, and brine, and dried (Na₂SO₄). After concentration, the residue was purified by silica gel column chromatography (pentane/ether, 50:1) to get 14 mg mixture of **6** and a THF from the starting material. The mixture analyzed by chiral stationary phase GC on the Cyclodex B column revealed that the product was nearly racemic (<3% ee).

Synthesis of *N*-(4-toluensulfonyl)-2-aza-bicyclo[2.2.1]hept-5-ene (**7**)

This compound was prepared according to a literature procedure.⁴³ Cyclopentadiene (4 mL, 49.0 mmol) was added to a solution of NH₄Cl (1.325 g, 24.8 mmol), 36% aqueous formaldehyde (2.6 mL, 34.0 mmol) and MeOH (5 mL). The reaction mixture was vigorously stirred for 12 h at room temperature before diluting with an equal volume of water and was subsequently washed with diethyl ether (2 × 10 mL). The aqueous phase was made basic with saturated NaOH. The 2-azabicyclo[2.2.1]hept-5-ene was isolated by extraction with ether (3 × 10 mL). To a mixture of this extract and 10% NaOH (10 mL) was added a solution of TsCl (4 g, 21.0 mmol) in a mixed solvent (ether: 6 mL, CH₂Cl₂: 3 mL) over a period of 5 min at room temperature. The mixture was stirred for 12 h. The organic layer was separated, dried with Na₂SO₄, filtered and concentrated in vacuo. The resulting oil was purified by chromatography (EtOAc: hexanes = 1:7) to get the product (2.77 g, 53%) as a white solid: mp 67–70°C. ¹H NMR (500 MHz, CDCl₃): δ 1.40 – 1.46 (m, 2 H), 2.39 (s, 3 H), 2.52 (dd, *J* = 8.5 Hz, 1.3 Hz, 1 H), 3.11 (broad s, 1 H), 3.31 (dd, *J* = 8.5 Hz, 2.8 Hz, 1 H), 4.62 (broad s, 1 H), 5.97 – 5.99 (m, 1 H), 6.07 – 6.09 (m, 1 H), 7.25 (d, *J* = 8 Hz, 2 H), 7.65 (d, *J* = 8 Hz, 2 H); ¹³C NMR (125 MHz, CDCl₃): δ 21.5, 43.8, 47.1, 64.2, 127.7, 129.5,

133.5, 136.3, 136.7, 143.1. IR cm^{-1} : 1654, 1458, 1330, 1155, 1092, 662. HRMS (ESI) Calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_2\text{SNa}$ ($[\text{M} + \text{Na}]^+$) 272.0716, observed 272.0715.

Typical procedure for catalytic hydrovinylation of heterobicyclic alkenes (Table 2, entry 4)

The pre-catalyst was prepared as follows in a glovebox: To **di- η -allyl-di- μ -bromonickel(II)** (2.9 mg, 0.0080 mmol) in CH_2Cl_2 (1 mL) was added a solution of ligand **L4** (8.7 mg, 0.016 mmol) in CH_2Cl_2 (1 mL) at ambient temperature. The resulting solution was added to a suspension of Na BARF (14.3 mg, 0.016 mmol) suspended in dichloromethane (1 mL) and the mixture was stirred at ambient temperature in a septum-sealed 25 mL round bottom flask for 10 min affording a dark brown solution containing a small amount of fine particulate (NaBr). The flask was removed from the glovebox. The flask was then cooled to -47°C acetonitrile-dry ice bath, creating a small vacuum. Dry ethylene (passed through a $0.5'' \times 4''$ column of Drierite®) was introduced *via* needle through the serum stopper and the vessel atmosphere was slowly evacuated 3 times with a 60 mL syringe. After cooling the solution to -47°C , a solution of **7** (50 mg, 0.20 mmol) in 1 mL dry CH_2Cl_2 is introduced dropwise into the solution of the pre-catalyst over a one minute period via syringe. The vessel was then maintained at -47°C for a period of 5 h. At the end of this period the ethylene line was removed and the reaction was quenched by addition of a saturated aqueous NH_4Cl solution. The product was extracted with ether. The organic layers are combined, dried over MgSO_4 and concentrated to give yellow oil. The resulting residue was purified by column chromatography on silica gel (eluting with hexanes/ethyl acetate = 9/1) to get 45 mg (80%) of an inseparable mixture of **8** and **9** as an oil.

Products of Hydrovinylation of **7** Using Ligand **L4**

^1H NMR (500 MHz, CDCl_3): (**8**) δ 0.84 – 0.86 (m, 1 H), 1.27 – 1.29 (m, 1 H), 1.35 – 1.44 (m, 1 H), 1.62 – 1.77 (m, 1 H), 2.40 (s, 3H), 2.44 (broad s, 1 H), 2.66 – 2.70 (m, 1 H), 2.98 – 3.07 (m, 2 H), 3.96 (broad s, 1 H), 4.89 – 4.98 (m, 2 H), 5.53 – 5.60 (m, 1 H), 7.28 (d, J = 8.2 Hz, 2 H), 7.68 (d, J = 8.2 Hz, 2 H); (**9**) δ 0.89 – 0.91 (m, 1 H), 1.29 – 1.31 (m, 1 H), 1.35 – 1.44 (m, 1 H), 2.01 – 2.06 (m, 1 H), 2.29 (broad s, 1 H), 2.30 – 2.36 (m, 1 H), 2.40 (s, 3 H), 2.98 – 3.07 (m, 2 H), 4.15 (broad s, 1 H), 4.89 – 4.98 (m, 2 H), 5.63 – 5.69 (m, 1 H), 7.28 (d, J = 8.2 Hz, 2 H), 7.68 (d, J = 8.2 Hz, 2 H); (**8+9**): ^{13}C NMR (125 MHz, CDCl_3): δ 21.5, 33.8, 34.0, 34.2, 37.4, 38.4, 43.0, 43.7, 47.0, 53.7, 54.0, 60.2, 64.0, 113.4, 114.8, 127.4, 127.4, 129.6, 135.8, 135.9, 139.8, 141.5, 143.2, 143.2. IR (neat) cm^{-1} : 3430, 2977, 2880, 1638, 1597, 1457, 1341, 1161, 1093, 818. HRMS (ESI) Calcd for $\text{C}_{15}\text{H}_{19}\text{NO}_2\text{SNa}$ ($[\text{M} + \text{Na}]^+$) 300.1029, observed 300.1041. HPLC conditions and retention times (Chiracel AD-H): solvent (hexanes:isopropanol) 98.5 : 1.5; flow rate 0.4 mL/min, retention times (min): for **8**: 43.06, 57.81; for **9**, 53.11, 60.77. (see attached chromatograms)

Hydrovinylation of **7** Using Ligand **L1**

The products of the reaction were separated into **10**, **11** and a mixture of **12** and **13** by column chromatography on silica gel (eluting with hexanes/ethyl acetate = 9/1). The mixture of **12** and **13** was identified after conversion to a ketone **14** (see later).

10: ^1H NMR (500 MHz, CDCl_3): δ 1.30 (d, J = 10.1 Hz, 1 H), 1.38 (d, J = 10.1 Hz, 1 H), 1.69 (d, J = 6.9 Hz, 3 H), 1.75 (d, J = 16 Hz, 1 H), 2.14 – 2.18 (m, 1 H), 2.39 (s, 3H), 2.51 (broad s, 1 H), 2.94 (d, J = 8.8 Hz, 1 H), 3.23 – 3.26 (m, 1 H), 4.71 (broad s, 1 H), 5.04 (q, J = 6.9 Hz, 1 H), 7.24 (d, J = 8.2 Hz, 2 H), 7.67 (d, J = 8.2 Hz, 2 H); ^{13}C NMR (125 MHz, CDCl_3): δ 14.8, 21.5, 35.2, 37.0, 38.0, 53.1, 59.5, 116.8, 127.6, 129.2, 136.1, 138.6, 143.0. White solid: mp $98 - 102^\circ\text{C}$; IR (neat) cm^{-1} : 3448, 2980, 1597, 1452, 1338, 1159, 1094, 667. HRMS (ESI) Calcd for $\text{C}_{15}\text{H}_{19}\text{NO}_2\text{SNa}$ ($[\text{M} + \text{Na}]^+$) 300.1029, observed 300.1037.

11: ^1H NMR (500 MHz, CDCl_3): δ 1.30 (d, $J = 6.9$ Hz, 3 H), 1.33 (d, $J = 10.4$ Hz, 1 H), 1.44 – 1.46 (m, 1 H), 1.51 (d, $J = 16.1$ Hz, 1 H), 2.07 – 2.11 (m, 1H), 2.39 (s, 3 H), 2.55 (broad s, 1 H), 2.88 (d, $J = 8.8$ Hz, 1 H), 3.27 – 3.30 (m, 1 H), 4.33 (broad s, 1 H), 5.34 – 5.38 (m, 1 H), 7.22 (d, $J = 8$ Hz, 2 H), 7.64 (d, $J = 8$ Hz, 2 H); ^{13}C NMR (125 MHz, CDCl_3): δ 13.4, 21.5, 32.5, 37.1, 38.4, 53.2, 64.9, 116.8, 127.9, 129.1, 129.2, 135.7, 138.6, 142.8. White solid: mp 95 – 98 °C; IR (neat) cm^{-1} : 3427, 2921, 1593, 1449, 1323, 1154, 1090, 1052, 668. HRMS (ESI) Calcd for $\text{C}_{15}\text{H}_{19}\text{NO}_2\text{SNa}$ ($[\text{M} + \text{Na}]^+$) 300.1029, observed 300.1044.

Oxidation of 12 and 13

The inseparable mixture of the two isomers **12** and **13** was oxidized to the ketone **14**. A flask is charged with a magnetic stirrer, 1 mL of carbon tetrachloride, 1.5 mL of water and 8.7 mg (0.031 mmol) of a mixture of **12** and **13**. To this mixture, sodium metaperiodate (33.6 mg, 0.157 mmol) was added. To this biphasic solution an acetonitrile solution (1 mL) of ruthenium trichloride hydrate (0.65 mg, 0.0031 mmol) was added. The entire mixture was stirred vigorously for 16 h at room temperature. The phases were separated and the aqueous phase was extracted 3 times with CH_2Cl_2 . The combined organic extracts were dried (MgSO_4) and concentrated. The crude product was purified by chromatography (EtOAc: hexanes = 1.0:1.5) to give the product (4.9 mg, 59%) as an oil. The structure was confirmed by comparison of ^1H NMR with the data reported in the literature.⁴³ ^1H NMR (500 MHz, CDCl_3): δ 1.46 – 1.49 (m, 1 H), 1.75 (d, $J = 10.7$ Hz, 1 H), 2.09 – 2.35 (ABX system, $\nu_a = 2.11$, $\nu_b = 2.32$, $J_{AB} = 18$ Hz, $J_{AX} = 4.4$ Hz, $J_{BX} = 3$ Hz, 2 H), 2.42 (s, 3 H), 2.82 (broad s, 1 H), 3.32 – 3.33 (m, 2 H), 4.54 (broad s, 1 H), 7.32 (d, $J = 8.2$ Hz, 2 H), 7.72 (d, $J = 8.2$ Hz, 2 H); ^{13}C NMR (125 MHz, CDCl_3): δ 21.6, 36.4, 46.0, 48.4, 50.8, 59.1, 127.4, 130.0, 135.0, 144.0, 211.8; IR (neat) cm^{-1} : 3352, 2918, 2852, 1708, 1679, 1366, 1161. HRMS (ESI) Calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_3\text{SNa}$ ($[\text{M} + \text{Na}]^+$) 288.0665, observed 288.0674.

Synthesis of Diisopropyl 2,3-diazabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylate (15)

The substrate was prepared following a procedure reported in the literature.⁴⁴ To a freshly distilled cyclopentadiene (0.62 mL, 7.60 mmol) solution in CH_2Cl_2 kept at 0 °C, was added the diisopropyl azodicarboxylate (1 mL, 5.08 mmol). The reaction was allowed to warm up to room temperature and stirred until full consumption of the starting material, as estimated by TLC. The solvent was evaporated under reduced pressure. The crude clear oil was purified by silica-gel chromatography with 1:3 EtOAc/hexanes as eluent. The diazabicyclo was obtained quantitatively as a white solid: mp 95 – 98 °C; ^1H NMR (500 MHz, CDCl_3): δ 1.22 (d, $J = 6.3$ Hz, 12 H), 1.65 – 1.70 (m, 2 H), 4.90 – 4.95 (m, 2 H), 5.10 (broad s, 2 H), 6.46 (broad s, 2 H); ^{13}C NMR (125 MHz, CDCl_3): δ 21.9, 47.9, 65.2, 70.1, 134.2, 138.4, 158.5; IR (neat) cm^{-1} : 3411, 2981, 1743, 1700, 1373, 1307, 1174, 1100; HRMS (ESI) Calcd for $\text{C}_{13}\text{H}_{20}\text{N}_2\text{O}_4\text{Na}$ ($[\text{M} + \text{Na}]^+$) 291.1315, observed 291.1311.

Hydrovinylation of 15. 16

^1H NMR (500 MHz, CDCl_3): δ 1.24 (d, $J = 6.3$ Hz, 12 H), 1.58 (broad s, 1 H), 1.61 – 1.64 (m, 2 H), 2.05 (broad s, 1 H), 2.68 (broad s, 1 H), 4.33 (broad s, 1 H), 4.51 (broad s, 1 H), 4.93 – 4.97 (m, 2 H), 5.02 – 5.05 (m, 2 H), 5.63 – 5.70 (m, 1 H); ^{13}C NMR (125 MHz, CDCl_3): δ 22.0, 35.0, 43.3, 60.3, 64.4, 69.9, 115.3, 139.0, 157.3; IR (neat) cm^{-1} : 2980, 2937, 2878, 1697, 1468, 1374, 1105, 918, 770; HRMS (ESI) Calcd for $\text{C}_{15}\text{H}_{24}\text{N}_2\text{O}_4\text{Na}$ ($[\text{M} + \text{Na}]^+$) 319.1628, observed 319.1627.

Hydrovinylation of 7-tert-Butoxycarbonyl-7-azabicyclo[2.2.1]hept-2-ene (17)

The starting material was prepared according to a procedure reported in the literature.⁴⁵ **18:** ^1H NMR (500 MHz, CDCl_3): δ 1.35 – 1.56 (m, 3 H), 1.42 (s, 9 H), 1.64 – 1.73 (m, 3 H),

2.29 – 2.33 (m, 1 H), 4.00 (broad s, 1H), 4.22 (broad s, 1 H), 4.88 (d, $J = 10.1$ Hz, 1 H), 4.94 (d, $J = 17.0$ Hz, 1 H), 5.70 – 5.77 (m, 1 H); ^{13}C NMR (125 MHz, CDCl_3): δ 28.3, 29.7, 47.4, 79.3, 113.0, 142.0. IR (neat) cm^{-1} : 3365, 2920, 2852, 1704, 1454, 1366, 801; HRMS (ESI) Calcd for $\text{C}_{13}\text{H}_{21}\text{NO}_2\text{Na}$ ($[\text{M} + \text{Na}]^+$) 246.1465, observed 246.1462; GC (Cyclodex B, programmed run, 5 min at 110°C , $0.5^\circ\text{C}/\text{min}$, and finally 5 min at 130°C). Retention times (min): 39.6, 40.3. Retention time for starting material 17 (min): 28.7.

cis-3,4-bis(benzyloxymethyl)-3,4-dimethylcyclobutene (**19**). To a solution of *cis*-1,2-dimethylcyclobut-3-ene-1,2-dimethanol ⁴⁶ (1.06 g, 7.4 mmol) in THF (30 mL) was added NaH (888 mg, 60 wt % in mineral oil, 22.2 mmol) in one portion at 0°C under argon. The resulting suspension was stirred at 0°C for 60 min and then benzyl bromide (2.83 mL, 23.8 mmol) was added dropwise at 0°C . The mixture was allowed to warm to room temperature and stirred for another 18 hours. Water was carefully added to quench the reaction and the mixture was extracted with ether and the organic layers were combined, washed with brine, dried and concentrated. The resulting residue was purified by column chromatography on silica gel (eluting with hexanes/ethyl acetate = 40/1) to get 2.16 g (90%) of *cis*-3,4-bis(benzyloxymethyl)-3,4-dimethylcyclobutene (**19**) as an oil: ^1H NMR (400 MHz, CDCl_3): δ 1.15 (s, 6 H), 3.41 (d, $J = 9.3$ Hz, 2 H), 3.54 (d, $J = 9.3$ Hz, 2 H), 4.40 (d, $J = 12.4$ Hz, 2 H), 4.44 (d, $J = 12.4$ Hz, 2H), 6.08 (s, 2 H), 7.21 – 7.31 (m, 10 H); ^{13}C NMR (100 MHz, CDCl_3): δ 19.3, 51.8, 73.2, 75.4, 127.3, 127.5, 128.2, 138.8, 141.1. IR (neat) cm^{-1} : 3029, 2852, 1496, 1453, 1362, 1094, 1075; HRMS (ESI) Calcd for $\text{C}_{22}\text{H}_{26}\text{O}_2\text{Na}$ ($[\text{M} + \text{Na}]^+$) 345.1830, observed 345.1833.

Hydrovinylation of **19**. **20**

^1H NMR (400 MHz, CDCl_3): δ 1.07 (s, 3 H), 1.20 (s, 3 H), 1.64 (dd, $J = 8.3$ Hz, 11.1 Hz, 1 H), 1.92 (t, $J = 10.4$ Hz, 1 H), 2.62 – 2.68 (s, 1H), 3.25 (d, $J = 9.0$ Hz, 1 H), 3.35 (d, $J = 9.0$ Hz, 1 H), 3.42 (d, $J = 9.0$ Hz, 1 H), 3.51 (d, $J = 9.0$ Hz, 1 H), 4.37 – 4.45 (m, 4 H), 4.90 – 4.94 (m, 2 H), 5.87 – 5.96 (m, 1 H), 7.24 – 7.32 (m, 10 H); ^{13}C NMR (100 MHz, CDCl_3): δ 20.2, 21.1, 32.8, 40.2, 43.7, 46.5, 73.1, 73.6, 76.4, 114.0, 127.2, 127.3, 127.3, 127.4, 128.2, 128.2, 138.7, 138.9, 139.1; IR (neat) cm^{-1} : 2959, 2926, 2854, 1454, 1360, 1095; HRMS (ESI) Calcd for $\text{C}_{24}\text{H}_{30}\text{O}_2\text{Na}$ ($[\text{M} + \text{Na}]^+$) 373.2144, observed 373.2142;

Determination configuration and enantiomeric excess of the hydrovinylation product **20**

To a solution of **20** (48 mg, 0.14 mmol) in THF (4 mL) was added 9-BBN (0.51 mL, 0.5 M, 0.6 mmol) at 0°C under nitrogen. The mixture was allowed to warm to room temperature and stirred for 3 hours. To the solution was added 1 mL of 2M NaOH and 0.4 mL of 30% aqueous H_2O_2 were added successively at 0°C and the resulting mixture was stirred at room temperature for 30 min. Water was added and the mixture was extracted with ether. The organic layer was washed with brine, dried over MgSO_4 and concentrated. The residue was purified by column chromatography (eluting with hexanes/ethyl acetate = 10/1) to afford 35 mg (70%) of **21**. ^1H NMR (400 MHz, CDCl_3): δ 1.02 (s, 3H), 1.11 (s, 3 H), 1.34 – 1.42 (m, 1 H), 1.45 – 1.56 (m, 2 H), 1.63 – 1.72 (m, 1 H), 1.96 – 2.06 (m, 1 H), 2.11 (broad s, 1 H), 3.07 (d, $J = 9.0$ Hz, 1 H), 3.26 (d, $J = 9.0$ Hz, 1 H), 3.40 (d, $J = 9.0$ Hz, 1 H), 3.37 – 3.43 (m, 1 H), 3.50 – 3.54 (m, 1 H), 3.59 (d, $J = 9.0$ Hz, 1 H), 4.29 – 4.37 (m, 4 H), 7.17 – 7.27 (m, 10 H); The configuration of the product is assigned by the strong $n\text{Oe}$ observed between the methyl hydrogens and the ring hydrogen shown. ^{13}C NMR (125 MHz, CDCl_3): δ 20.3, 21.1, 33.8, 34.2, 37.7, 40.2, 44.7, 62.4, 72.7, 73.1, 73.4, 76.1, 127.3, 127.4, 127.5, 127.6, 128.2, 128.3, 138.3, 138.9; IR (neat) cm^{-1} : 3364, 2930, 2866, 1717, 1454, 1361, 1072, 736; HRMS (ESI) Calcd for $\text{C}_{24}\text{H}_{32}\text{O}_3\text{Na}$ ($[\text{M} + \text{Na}]^+$) 391.2249, observed 391.2252; $[\alpha]_{\text{D}}^{22} = -16.9$ (c 0.51, CHCl_3) (**L6**); HPLC (chiracel AD-H) conditions: hexanes:isopropanol 99:1, 0.3 mL/min, retention time (min): 73.43 (–)-isomer, 77.38 (+)-isomer. The retention times were

confirmed by comparison to an authentic racemic **21** prepared using nickel complex of **L2** (see attached chromatograms).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial assistance for this research by NSF (CHE-0610349) and NIH (General Medical Sciences, R01 GM075107) is gratefully acknowledged.

References

1. Chauvin, Y.; Olivier, H. Dimerization and Codimerization. In: Cornils, B.; Herrmann, WA., editors. Applied Homogeneous Catalysis with Organometallic Compounds. Vol. 1. VCH; New York: 1996. p. 258-268.
2. Su ACL. Adv Organomet Chem. 1979; 17:269.
3. For seminal studies and reviews, see: Hata G. J Am Chem Soc. 1964; 86:3903. Wittenberg D. Angew Chem, Int Ed Engl. 1964; 3:153. Alderson T, Jenner EL, Lindsey RV Jr. J Am Chem Soc. 1965; 87:5638. Bogdanovi'c B, Henc B, Meister B, Pauling H, Wilke G. Angew Chem, Int Ed Engl. 1972; 11:1023. Bogdanovi'c B. Adv Organomet Chem. 1979; 17:105. Fe: Ehlers J, König WA, Lutz S, Wenz G, tom Dieck H. Angew Chem, Int Ed Engl. 1988; 27:1556. Moreau B, Wu JY, Ritter T. Org Lett. 2009; 11:337. [PubMed: 19093845] Use of Ni: Wilke G. Angew Chem, Int Ed Engl. 1988; 27:185. RajanBabu TV. Chem Rev. 2003; 103:2845. [PubMed: 12914483] For more recent reports, see: Ru: Gooßen LJ. Angew Chem, Int Ed Engl. 2007; 46:7544. [PubMed: 17879243] Ura Y, Tsujita H, Mitsudo T-a, Kondo T. Bull Korean Chem Soc. 2007; 28:2139. Co: Hilt G, du Mesnil FX, Lüers S. Angew Chem, Int Ed Engl. 2001; 40:387. [PubMed: 11180333]
4. For representative examples, see: Jolly PW, Wilke G, Cornils B, Herrmann WA. Hydrovinylation. Applied Homogeneous Catalysis with Organometallic Compounds. VCH New York 1996; 2:1024–1048. Nomura N, Jin J, Park H, RajanBabu TV. J Am Chem Soc. 1998; 120:459. Gooßen LJ. Angew Chem, Int Ed Engl. 2002; 41:3775. [PubMed: 12386846] RajanBabu TV, Nomura N, Jin J, Radetich B, Park H, Nandi M. Chem Eur J. 1999; 5:1963. He Z, Yi CS, Donaldson WA. Org Lett. 2003; 5:1567. [PubMed: 12713325] Diez-Holz CJ, Böing C, Franció G, Hölscher M, Leitner W. Eur J Org Chem. 2007; 2995. Tsujita H, Ura Y, Matsuki S, Wada K, Mitsudo T, Kondo T. Angew Chem, Int Ed. 2007; 46:5160. Ho C-Y, Ohmiya H, Jamison TF. Angew Chem Int Ed. 2008; 47:1893. Zhang Q, Zhu SF, Qiao XC, Wang LX, Zhou QL. Adv Synth Catal. 2008; 350:1507. Grutters MMP, van der Vlugt JI, Pei Y, Mills AM, Lutz M, Spek AL, Müller C, Moberg C, Vogt D. Adv Synth Catal. 2009; 351:2199. RajanBabu TV. Synlett. 2009:853. [PubMed: 19606231] Sharma RK, RajanBabu TV. J Am Chem Soc. 2010; 132:3295. [PubMed: 20163120]
5. Bogdanovi'c B, Henc B, Lösler A, Meister B, Pauling H, Wilke G. Angew Chem Int Ed Engl. 1973; 12:954. Nozaki's Cu-catalyzed cyclopropanation of styrene with ethyl diazoacetate is the only other example of an asymmetric C-C bond-forming reaction that precedes this discovery. Nozaki H, Moriuti S, Takaya H, Noyori R. Tetrahedron Lett. 1966:5239.
6. Muller G, Ordinas JI. J Mol Catal, A: Chem. 1997; 125:97.
7. Yi CS, He Z, Lee DW. Organometallics. 2001; 20:802.
8. Bianchini C, Giambastiani G, Meli A, Toti A. Organometallics. 2007; 26:1303.
9. Kumareswaran R, Nandi N, RajanBabu TV. Org Lett. 2003; 5:4345. [PubMed: 14601996]
10. For other related metal-catalyzed C-C bond-forming reactions of norbornene, see: Hydrocyanation: Hodgson M, Parker D, Taylor RJ, Ferguson G. Organometallics. 1988; 7:1761. Baker MJ, Pringle PG. J Chem Soc, Chem Commun. 1991:1292. Hydroformylation: Horiuchi T, Shirakawa E, Nozaki K, Takaya H. Tetrahedron: Asymmetry. 1997; 8:57. Yan M, Xu QY, Chan ASC. Tetrahedron: Asymmetry. 2000; 11:845. Hydrosilylation: Hayashi T. Acta Chem Scand. 1996; 50:259. Hydroalkenylation: Ozawa F, Kabatake Y, Kubo A, Hayashi T. J Chem Soc, Chem Commun. 1994:1323. Sakuraba S, Awano K, Achiwa K. Synlett. 1994:291. Wu XY, Xu HD, Tang

- FY, Zhou QL. *Tetrahedron: Asymmetry*. 2001; 12:2565. Aufdenblatten R, Diezi S, Togni A. *Monat Chem*. 2000; 131:1345.
11. Nandi M, Jin J, RajanBabu TV. *J Am Chem Soc*. 1999; 121:9899.
 12. Zhang A, RajanBabu TV. *Org Lett*. 2004; 6:1515. [PubMed: 15101781]
 13. Zhang A, RajanBabu TV. *J Am Chem Soc*. 2006; 128:54. [PubMed: 16390118]
 14. Saha B, Smith CR, RajanBabu TV. *J Am Chem Soc*. 2008; 130:9000. [PubMed: 18570419]
 15. For example, hydrovinylation of styrene using [(allyl)Ni]Br₂/ligand **L2**/NaBARF (0.007 equiv catalyst) at room temperature in an atmosphere of ethylene gave >99% yield of 3-phenyl-1-butene with >99% selectivity for this isomer. Under these conditions most other ligands leads to extensive isomerization of the initial product into (Z)- and (E)-2-phenyl-2-butenes.
 16. Franció G, Faraone F, Leitner W. *J Am Chem Soc*. 2002; 124:736. [PubMed: 11817933]
 17. Park H, Kumareswaran R, RajanBabu TV. *Tetrahedron*. 2005; 61:6352.
 18. Smith CR, Mans DJ, RajanBabu TV. *Org Synth*. 2008; 85:238. [PubMed: 19655040]
 19. Smith CR, Lim HJ, Zhang AB, RajanBabu TV. *Synthesis*. 2009:2089. [PubMed: 19763244]
 20. Lassauque N, Francio G, Leitner W. *Adv Synth Catal*. 2009; 351:3133.
 21. Smith CR, RajanBabu TV. *Org Lett*. 2008; 10:1657. [PubMed: 18351772]
 22. Larock RC, Johnson PL. *J Chem Soc, Chem Commun*. 1989:1368.
 23. Ozawa F, Kobatake Y, Kubo A, Hayashi T. *J Chem Soc, Chemical Commun*. 1994:1323.
 24. Menard F, Lautens M. *Angew Chem Int Ed*. 2008; 47:2085.
 25. Lautens, M.; Rovis, T. Hydroalumination of Carbon-Carbon Double Bonds. In: Jacobsen, EN.; Pfaltz, A.; Yamamoto, H., editors. *Comprehensive Asymmetric Catalysis*. Vol. 1. Springer; Berlin: 1999. p. 337-348.
 26. Lautens M, Renaud JL, Hiebert S. *J Am Chem Soc*. 2000; 122:1804.
 27. Lautens M, Dockendorff C, Fagnou K, Malicki A. *Org Lett*. 2002; 4:1311. [PubMed: 11950350]
 28. Li LP, Rayabarapu DK, Nandi M, Cheng CH. *Org Lett*. 2003; 5:1621. [PubMed: 12735736]
 29. Rayabarapu DK, Cheng CH. *Acc Chem Res*. 2007; 40:971. [PubMed: 17591745]
 30. McManus HA, Fleming MJ, Lautens M. *Angew Chem Int Ed*. 2007; 46:433.
 31. Feng CC, Nandi M, Sambaiah T, Cheng CH. *J Org Chem*. 1999; 64:3538. [PubMed: 11674478]
 32. Zuercher WJ, Hashimoto M, Grubbs RH. *J Am Chem Soc*. 1996; 118:6634.
 33. Tallarico JA, Randall ML, Snapper ML. *Tetrahedron*. 1997; 53:16511.
 34. Limanto J, Snapper ML. *J Am Chem Soc*. 2000:122.
 35. Bassindale MJ, Hamley P, Harrity JPA. *Tetrahedron Lett*. 2001; 42:9055.
 36. Nicolaou KC, Vega JA, Vassilikogiannakis G. *Angew Chem, Int Ed*. 2001; 40:4441.
 37. Huffman MA, Liebeskind LS. *J Am Chem Soc*. 1993; 115:4895.
 38. Namyslo JC, Kaufmann DE. *Chem Rev*. 2003; 103:1485. [PubMed: 12683789]
 39. Zhang A, RajanBabu TV. *J Am Chem Soc*. 2006; 128:54. [PubMed: 16390118]
 40. Smith CR, Mans DJ, RajanBabu TV. *Org Synth*. 2008; 85:238. [PubMed: 19655040]
 41. Smith CR, Zhang A, Mans DJ, RajanBabu TV. *Org Synth*. 2008; 85:248. [PubMed: 19672483]
 42. Setzer WN, Brown ML, Yang X-j, Thompson MA, Whitaker KW. *J Org Chem*. 1992; 57:2812.
 43. Gleim RD, Spurlock LA. *J Org Chem*. 1976; 41:1313.
 44. (a) Diels O, Bolm JH, Knoll W. *Justus Liebigs Ann Chem*. 1925; 443:242. (b) Menard F, Lautens M. *Angew Chem Int Ed*. 2008; 47:2085.
 45. Carroll FI, Liang F, Navarro HA, Briecaddy LE, Abraham P, Damaj MI, Martin BR. *J Med Chem*. 2001; 44:2229. [PubMed: 11405659]
 46. Inoue M, Lee N, Kasuya S, Sato T, Hiramama M, Moriyama M, Fukuyama Y. *J Org Chem*. 2007; 72:3065. [PubMed: 17355151]

Figure 1.
Ligands for Hydrovinylation of Strained Alkenes

Figure 2.
Heterocyclic Compounds that Failed to Undergo Hydrovinylation

Scheme 1.

Ligand Dependence on the Ni-Catalyzed Hydrovinylation of Norbornene

Scheme 2.

Hydrovinylation of a [2.2.1]-Heterobicyclic Amide (7)

Scheme 3.
Hydrovinylation of a Cyclobutene (**19**)

Table 1

Hydrovinylation of **3**

entry	ligand	conditions (mol% cat, temp, time h)	yield of 4 (%) selec. (% ee) sign of [α] for major
1.	L1	3/-22 °C/3	70 ^{b,c} /NA
2.	L1	4/23 °C/2	<5/NA
3.	L2	3/23 °C/2	48 ^{b,c} /NA
4.	L3	4/23 °C/2	72 ^{b,c,d} /NA
5.	L4	3/-78 °C/5	92/93 (-)
6.	ent-L4	3/-78 °C/6	93/95-99 (+) ^e
7.	L5	3/-78 °C/6	95/66 (-)
8.	L6	3/-78 °C/6	92/94 (-)

^a See eq 2 and Experimental Section for details of the procedure.

^b Product isolated as a mixture of **3** and **4**.

^c rest starting material.

^d After 10 h, no starting material, only isomerization product, **5** along with other contaminants.

^e Product *ent-4*

Table 2

Ni-Catalyzed Hydrovinylation of **7**^a

entry	ligand	conditions (mol% cat, temp, time in h)	yield of HV (%)	products (%)	enantioselectivity for 8 , 9 (%ee)
1.	L1	14/-10 °C/4	78 ^b	10 (25), 11 (22) [12 + 13] (31)	--
2.	L2	8/20 °C/21	79	8 (46), 9 (33)	--
3.	L3	8/20 °C/21	70	8 (40), 9 (30)	--
4.	L4	8/-47 °C/5	80	8 (38), 9 (42)	8 (21), 9 (33)
5.	L5	8/-47 °C/6	89	8 (53), 9 (36)	8 (7), 9 (0)
6.	L6	8/-47 °C/6	89	8 (37), 9 (52)	8 (20), 9 (31)

^a See Scheme 2 and Experimental Section for procedure.

^b at -10 °C, **8** and **9** fully isomerized to **10**, **11**, **12** and **13**; at -47 °C, 22% yield of [**8**+**9**] with rest starting material.

Table 3**Ni-Catalyzed Hydrovinylation of Cyclobutene **19****

entry	ligand	conditions (mol% cat, temp, time in h)	yield of 2 (%); (20 : 19) ^a	% ee (20) 0 sign of [α]
1.	L1	14/−30 °C/15	32; 20:1	--
2. ^b	L2	14/43 °C/12	67; 5:1	--
4.	L3	14/23 °C/18	0 ^c	--
5.	L4	10/−50 °C/8	47; 4.3:1.0	55 (−)
6.	L4	14/−45 °C/6.5	40; 32:1	--
8.	L5	10/−50 °C/3.5	46; 1.4:1.0	52 (−)
7.	L6	10/−50 °C/5	48; 4.7:1.0	82 (−)

^aRatio and yield of product calculated based on ¹H NMR of a mixture of **19** and **20**.

^bin ClCH₂CH₂Cl

^ccomplex mixture