Published in final edited form as:

Mol Ecol. 2006 October ; 15(12): 3849–3861. doi:10.1111/j.1365-294X.2006.03044.x.

Diversity of microcystin genotypes among populations of the filamentous cyanobacteria *Planktothrix rubescens* and *Planktothrix agardhii*

Rainer Kurmayer¹ and Marlies Gumpenberger²

Austrian Academy of Sciences, Institute for Limnology, Mondseestraße 9, 5310 Mondsee, Austria

Abstract

Microcystins (MCs) are toxic heptapeptides that are produced by filamentous cyanobacteria *Planktothrix rubescens* and *P. agardhii* via non-ribosomal peptide synthesis. MCs share a common structure cyclo (-D-Alanine¹-L-X²-D-erythro-β-iso-aspartic acid³-L-Z⁴-Adda⁵-D-Glutamate⁶-N-methyl-dehydroalanine⁷) where X² and Z² are variable L-amino acids in positions 2, 4 of the molecule. Part of the *mcyB* gene (1,451 bp) that is involved in the activation of the X² amino acid during MC synthesis was sequenced in 49 strains containing different proportions of arginine, homotyrosine, and leucine in position 2 of the MC molecule. Twenty-five genotypes were found that consisted of eight genotype groups (A-H, comprising 2–11 strains) and 17 unique genotypes. *P. rubescens* and *P. agardhii* partly consisted of the same *mcyB* genotypes. The occurrence of numerous putative recombination events that affected all of the genotypes can explain the conflict between taxonomy and *mcyB* genotype distribution. Genotypes B (homotyrosine and leucine in X²) and C (arginine in X²) showed higher nonsynonymous/synonymous (d_N/d_S) substitution ratios implying a relaxation of selective constraints. In contrast other genotypes (arginine, leucine, homotyrosine) showed lowest d_N/d_S ratios implying purifying selection. Restriction fragment length polymorphism (RFLP) revealed the unambiguous identification of *mcyB* genotypes, which are indicative of variable X² amino acids in eight populations of *P. rubescens* in the Alps (Austria, Germany, Switzerland). The populations were found to differ significantly in the proportion of specific genotypes and the number of genotypes that occurred over several years. It is concluded that spatial isolation might favour the genetic divergence of microcystin synthesis in *Planktothrix* spp.

Keywords

microcystin synthesis; microevolution; geographic isolation; population genetics; plankton ecology; toxicity

¹Corresponding author: Rainer Kurmayer, Austrian Academy of Sciences, Institute for Limnology, Mondseestraße 9, 5310 Mondsee, Austria, Phone 0043-6232-312532, Fax 0043-6232-3578, rainer.kurmayer@oeaw.ac.at.

²Present address: Potsdam Institute for Climate Impact Research, Telegrafenberg, P.O. Box 60 12 03, D-14412 Potsdam, Germany

Author information

Our research aims towards understanding the evolution and maintenance of the impressive diversity of secondary metabolites in cyanobacteria. Apart from the problems involved in using water bodies affected by toxic cyanobacteria, the evolution of genes involved in secondary metabolite synthesis is also interesting from a biological perspective. By linking molecular insights with ecological research, we hope to identify the evolutionary mechanisms affecting the synthesis of natural products on a molecular and ecosystem level.

Introduction

Toxic heptapeptides microcystins are produced by various genera of the phylum of cyanobacteria, i.e. the genera *Microcystis*, *Anabaena*, and *Planktothrix*. The filamentous cyanobacteria *Planktothrix* spp. are one of the most important microcystin producers and can be found in freshwater habitats in the temperate region of the Northern Hemisphere (Fastner *et al.* 1999). *P. agardhii* co-occurs with *Limnolthrix redekei* and other filamentous cyanobacteria, e.g. *Pseudoanabaena limnetica* in high abundance in shallow and eutrophic lakes (Rücker *et al.* 1997). In contrast *P. rubescens* forms metalimnetic layers in deep stratified and less eutrophic lakes, and often monopolizes resources and dominates phytoplankton completely (Anneville *et al.* 2004).

Microcystins (MCs) are cyclic heptapeptides that are produced by cyanobacteria and share a common structure cyclo (-D-Ala¹ -L-X² -D-MeAsp³ -L-Z⁴ -Adda⁵ -D-Glu⁶ -Mdha⁷), where X and Z are variable L-amino acids (e.g. LR refers to leucine and arginine in the variable positions), D-MeAsp is D-erythro-β-iso-aspartic acid, Adda is (2*S*, 3*S*, 8*S*, 9*S*-3-amino-9-methoxy-2,6,8-trimethyl-10-phenyldeca-4,6-dienoic acid), and Mdha is N methyl-dehydroalanine (Carmichael *et al.* 1988). MCs are synthesized by thiotemplate mechanism such as other non-ribosomal peptides produced by bacteria and fungi (Marahiel *et al.* 1997). The large enzyme complex encoded by the *mcy* gene cluster is composed of peptide synthetases, polyketide synthases, and tailoring enzymes (Tillett *et al.* 2000, Christiansen *et al.* 2003). It has a modular structure, each module of which contains specific functional domains for activation = aminoacyl adenylation (A) domains, and thioesterification (thiolation domains) of the amino acid substrate and for the elongation (condensation domains) of the growing peptide (Tillett *et al.* 2000). Adenylation domains show a high degree of conservation (core motifs) enabling the definition of general rules for the structural basis of substrate recognition of non-ribosomal peptide synthetases (Marahiel *et al.* 1997). The determination of substrate specificity is based on the crystallization of the adenylation domain of gramicidin synthetase (*GrsA*) (Conti *et al.* 1997). This enabled the identification of core motifs constituting a putative binding pocket for the activation of the amino acid substrate during peptide synthesis and the *in silico* identification of critical binding pocket residues that are located between core motifs A3-A6 (Challis *et al.* 2000, Stachelhaus *et al.* 1999).

McyBA1 is responsible for the activation of amino acids as aminoacyl adenylate followed by peptide bond formation to the growing microcystin molecule by the condensation domain located upstream of *mcyBA1* (Tillett *et al.* 2000). In *Planktothrix* spp. three different amino acids have been described in position 2 of the microcystin molecule: leucine, arginine, homotyrosine (Luukkainen *et al.* 1993, Henriksen & Moestrup 1997, Fastner *et al.* 1999, Kurmayer *et al.* 2005). In an earlier study, we described the occurrence of novel microcystin variants, which appeared to be unique and dominant for *Planktothrix* and that exclusively occurred in a specific lake, Schwarzensee in the Austrian Alps (Kurmayer *et al.* 2004). It has been speculated that the populations of *P. rubescens* may be geographically isolated, for example Schwarzensee, the lake of origin of the isolates containing those novel microcystin variants is at an altitude of 716 m above sea level, while many other lakes have a lower altitude, such as Mondsee, which is located only 11.4 km away.

Typically the distribution of microorganisms is not restricted by geographical barriers (Finlay 2002). Studies have shown that recombination and migration rates in natural populations of soil bacteria (*Bacillus* spp.) and aquatic cyanobacteria (*Microcoleus* sp., *Nodularia* sp.) were high, and were not influenced by geographic factors (Roberts & Cohan 1995, Barker *et al.* 2000, Lodders *et al.* 2005). On the other hand, the occurrence of physical isolation, which is defined as the spatial separation of two or more populations among

geographically isolated habitats (Papke & Ward 2004) cannot *a priori* be excluded. Geographic isolation has been observed among isolated populations of extreme habitats, for example, *Synechococcus* inhabiting hot springs (Papke *et al.* 2003) and hyperthermophilic Archaea (Whitaker *et al.* 2003), but also for free living *Pseudomonas* spp. in pristine soil samples (Cho & Tiedje 2000). The question of isolation is of fundamental importance to understand the evolution of microcystin synthesis, i.e. populations may diverge through neutral processes (i.e. random drift, Kimura 1983), local adaptation processes, or both.

The ability to detect and interpret genotypic variation depends on the type of the gene under investigation and the resolving power of the genetic markers. Genes showing high variability with potential functional consequences are considered highly suitable to interpret genotypic variation in potentially isolated populations (Palys *et al.* 1997). The species *P. rubescens* (*sensu* Suda *et al.* 2002) only consists of microcystin genotypes while - similar to *Microcystis* sp. - populations assigned to *P. agardhii* consisted of microcystin genotypes and genotypes without the *mcy* genes (Kurmayer *et al.* 2004). It has been shown that *Microcystis* sp. colonies that were isolated from field samples of a eutrophic lake show a high genetic variability within *mcyBA1* (Kurmayer *et al.* 2002), for example, due to domain duplication and recombination within the *mcyB* gene (Mikalsen *et al.* 2003).

It was the aim of the present study (i) to analyse the genetic variation within strains, (ii) identify genotypes that are indicative of the production of specific MC variants, (iii) and to directly detect and quantify those genotypes in various lakes in the Alps. This knowledge is necessary to understand MC net production in phytoplankton as well as the microevolution of microcystin synthesis in cyanobacteria. For this purpose, 49 *Planktothrix* strains isolated in Europe were sequenced for the first adenylation domain (A1) of the *mcyB* gene. In order to characterise populations directly (and to overcome a possible isolation bias), restriction fragment length polymorphism (RFLP) profiles were assigned to each *mcyBA1* genotype and subsequently used to distinguish the gene pools of spatially separated populations.

Materials and Methods

Cultivation and sampling of cyanobacteria

The 49 strains that were used in this study were either isolated from several European freshwater habitats, as described in Kurmayer *et al.* (2004), or acquired from international culture collections (Table 1). The isolates were assigned to the genus *Planktothrix* (*P. agardhii* and *P. rubescens*, Suda *et al.* 2002, min. 99.2% similarity in 16S rDNA) according to the morphological criteria provided by Komarek (2003). In addition all of the strains were genetically tested using *Planktothrix* specific primers binding to the intergenic spacer region within the phycocyanin operon (PC-IGS) as described in Kurmayer *et al.* (2004). Clonal strains were grown in BG-11 medium (Rippka 1988) modified to contain 2 mM NaNO₃ + 10 mM NaHCO₃ and kept in batch culture at 15°C under continuous light conditions (5–10 μmol m⁻² s⁻¹, Osram Type L30W/77 Fluora).

P. rubescens was sampled by pulling a plankton net (30 μm in mesh size) from a depth of 20 m to the surface at the deepest part of eight lakes in the Alps (Table 2). Aliquots (a few ml from net samples) were filtered onto glass fibre filters (GF/C, Whatman, Kent, Great Britain) under vacuum pressure and stored frozen (–20°C) until DNA extraction. In this study each lake was sampled a total of three times from 2001–2005.

Microcystin analysis

All strains were analysed for their composition of microcystin variants, i.e. amino acid composition in position 2 of the MC molecule by HPLC-DAD (high performance liquid chromatography with diode array detection) as described (Kurmayer *et al.* 2004, 2005).

Briefly, the cells were filtered on glass fibre filters (GF/C, Comesa, Vienna, Austria) and MCs were extracted using 75% (w/v) aqueous methanol. MC variants were quantified at 240 nm by their characteristic absorption spectra (original spectrum and first order derivative) and retention times (Fastner *et al.* 1999) using a linear gradient of acetonitrile (0.05% TFA, Trifluoroacetic acid) against water (0.05% TFA) according to Lawton *et al.* (1994).

Demethylated variants containing arginine in position 2 of the molecule, i.e. [D-Asp³, Mdha⁷]-RR and [D-Asp³, Dhb⁷]-RR eluted from 13.4-13.9 min and 14.3-14.8 min, each, those containing homotyrosine (Hty) in position 2, [Asp³]-MC-HtyR eluted from 18.3-18.5 min, and those containing leucine in position 2, [Asp³]-MC-LR eluted 19.3-19.6 min (Kurmayer *et al.* 2005). All strains from Lake Schwarzensee were found to produce new microcystin variants eluting from 23.5-24.0 min (Kurmayer *et al.* 2004). Amino acid analysis, one-dimensional, and two-dimensional NMR revealed that the two novel variants contained homotyrosine/tyrosine in position 2 of the molecule: [D-Asp³, (E)-Dhb⁷]-MC-HtyY and [D-Asp³, (E)-Dhb⁷]-MC-HtyHty (R. Kurmayer, W. Yoshida, K. Ishida, T. Hemscheidt, unpublished).

Genetic analysis

Two millilitres of each strain culture were incubated 1h on ice and centrifuged at 16,000 g for 10 min. The pellet was lyophilised in a vacuum centrifuge at 30°C. DNA extraction from the strains or field samples was performed as described in Kurmayer *et al.* (2003).

PCR amplifications were performed in a volume of 20 µl, containing 2 µl of Qiagen PCR buffer (Qiagen, VWR, Vienna, Austria), 1.2 µl MgCl₂ (25 mM, Qiagen), 0.6 µl deoxynucleotide triphosphates (10 µM each, MBI Fermentas, St Leon-Rot, Germany), 1 µl of each primer (10 pmol µl⁻¹), 0.1 µl Taq DNA polymerase (5 u µl⁻¹, Qiagen), 13.1 µl sterile Millipore water and 1.0 µl of the diluted DNA extract (1:100). The PCR thermal cycling protocol included an initial denaturation at 94°C for 3 min, followed by 35 cycles at 94°C for 30 s, with an annealing temperature of 52°C for 30 s, and elongation at 72°C for 2 min, followed by 72°C for 5 min.

Oligonucleotide primers for *mcvBA1* amplification and sequencing are published in Kurmayer *et al.* (2005). *McyBA1*totfwd and *McyBA1*totrev were used to amplify the total fragment of the first adenylation domain (A1) of *mcvB* in strains and field samples (1,693 bp, including the core motifs A1-A10; Marahiel *et al.* 1997). The primers were specific for *Planktothrix*. Amplification products for *mcvB* were purified using the Qiagen QIAquick PCR Purification Kit (Qiagen, VWR, Vienna, Austria) and sequenced directly by standard automated fluorescence techniques (Applied Biosystems, Weiterstadt, Germany). These sequence data have been submitted to the DDBJ/EMBL/GenBank databases under the accession numbers: AJ890255-AJ890282 (Table 1).

In addition *mcvBA1* PCR products were digested using *AluI* and *TseI*. The restriction profiles were compared with the *mcvBA1* sequences. Restriction analysis was performed directly from the PCR amplification product using *AluI* (MBI Fermentas, St Leon-Rot, Germany) and *TseI* (New England Biolabs, Frankfurt am Main, Germany) according to the manufacturer's instructions. Digestions were performed in a volume of 20 µl, containing 0.2 µl of the restriction enzyme (5 u µl⁻¹) and 7 µl of the PCR products and incubated at 37°C (*AluI*) and 65°C (*TseI*) for 3h. The PCR products for *mcvB* were visualized in 1% agarose in 0.5 × TBE (Tris-borate-EDTA buffer) and ethidium-bromide staining performed *via* agarose gel electrophoresis according to standard procedures (Sambrook *et al.* 1989). For the electrophoresis of restriction fragments, 2% agarose was used.

PCR products for *mcyBA1* obtained from field samples were cloned using the pDrive Cloning Vector system (Qiagen, VWR, Vienna, Austria) according to the manufacturer's instructions. The transformation efficiency was $2.1 \times 10^4 - 7.0 \times 10^4$ colonies per μg of DNA. After the transformed colonies were grown on an agar plate overnight, white colonies were randomly picked using the tip of a pipette, and re-dissolved in 10 μl of Millipore water each. Re-PCR amplification from the white clones was always successful and the subsequent restriction analysis of colony specific PCR products revealed the unambiguous identification of a specific restriction type. The sequencing of *mcyBA1* primer binding sites performed for all of the different *A1uI* restriction types observed among strains (Table 3) revealed high similarity, i.e. identical nucleotides for the binding site of the *mcyBA1*totrev primer (5'-AGA CTT GTT TAA TAG CAA AGG C-3'). One substitution of adenine by cytosine in position 6 of the *mcyBA1*totfwd primer for restriction types I, IV (5'-CAC CTA GTT GAA GAA CAA GTT CT-3') was found. Consequently the PCR amplification of the restriction types from the field samples was considered unbiased.

Phylogeny and statistical analysis

Sequences (1,451 bp) were aligned using multiple sequence alignment (Clustal W 1.8) and similarities between nucleotide sequences were calculated using DNADIST in the PHYLIP package (Version 3.6 alpha; Felsenstein 1989). Maximum likelihood analysis was used to estimate nucleotide substitution parameters under a general time-reversible nucleotide substitution model by estimating the gamma distribution for variable rates among the sites. Ambiguous sites at which at least one sequence showed an undetermined nucleotide were removed (two sites) and the discrete gamma algorithm was used to approximate a continuous gamma distribution using five categories of rates (ncatG = 5) in the programme BASEML of the PAML package (Version 3.14, Yang 1997). The best fit of the model showed (log-) likelihood $\ln L = -2871.96$ with parameter estimates for the transition/transversion ratio $\kappa = 2.07$ and the shape parameter $\alpha = 0.006$. Statistical significance of the branches was estimated by bootstrap analysis generating 100 replicates of the original data set using the PHYLIP package. Finally consensus trees following the 50% majority rule were computed.

The ratio of nonsynonymous (d_N) and synonymous (d_S) substitution rates per site was determined using the likelihood approach, which was implemented in the program CODEML of the PAML package (Yang 1998). The "one ratio" model, assuming the same ratio for all of the branches in the phylogeny and the "free ratio" model, assuming as many ratios as the number of branches in the phylogenetic tree were used. Branch-site models (Zhang *et al.* 2005) were employed to test for positive selection acting on specific branches in the phylogenetic tree showing increased d_N/d_S ratios. Branches of the tree were divided *a priori* into foreground and background lineages, and a likelihood ratio test was constructed by comparing a model that allows positive selection on the foreground lineages (the alternative model) with a model that does not allow such positive selection (the null model). The improved branch-site likelihood method for detecting positive selection implemented in the PAML package was used (Version 3.15, Anisimova *et al.* 2003, Zhang *et al.* 2005).

Following Tanabe *et al.* (2004) the runs test implemented in the programme GENECONV (v1.81, Sawyer 1999) was used to investigate whether substitutions were significantly clustered, and whether gene conversion (recombination) events occurred within *mcyBA1*. The settings used were the default (/g0 = mismatches within fragments were not allowed). The Global *P*-value calculated from 10,000 random permutations of the alignment was used to assess the significance of any unusually long fragments that were sufficiently similar to be suggestive of past gene conversion. Recombination events were independently quantified using a recombination detection programme (RDP) developed by Martin & Rybicki (2000) using the default settings (window size 10, highest acceptable *p*-value 0.05).

The frequency of the occurrence of genotypes and restriction types between the *Planktothrix* species was compared using the Yates-corrected chi-square statistic in the analyses of 2×2 contingency tables (Zöfel 1992). The occurrence of restriction types between lakes was statistically compared using chi-square statistics for 8×2 contingency tables. Because the test can produce false results when the expected frequencies are small, proportions with an expected frequency <1 were not tested (Jongman *et al.* 1995). The differences in the total number of restriction types found in a population were tested using one-way ANOVA followed by the Tukey test for pairwise comparison (overall significance $p=0.01$).

Results

Genetic variation within the microcystin *mcvB* gene

Among the *mcvBA1* sequences obtained from the 49 *Planktothrix* strains, 25 different genotypes were detected, which differed in at least one base pair. We found eight *mcvBA1* genotypes (A-H) consisting of 2-11 strains each and 17 unique genotypes (Table 1). The genetic variability was between zero and 3.6%. Genotype B (11 strains) consisted of both species *P. agardhii* and *P. rubescens* while A (8), H (2), G (2), D (2), F (3) consisted of *P. rubescens* only. In contrast C (2) and E (2) occurred significantly in *P. agardhii* only (Chi-square test, χ^2 (df=1), $p < 0.05$).

The ratio of nonsynonymous (d_N) and synonymous (d_S) substitution rates for the entire phylogenetic tree was $d_N/d_S = 0.19$. Genotypes B, C, and the clade consisting of genotypes A (D, E, H, strains No34, 64, 91/1) showed a relatively high frequency of nonsynonymous substitutions resulting in $d_N/d_S = 0.73, 0.87, 0.23$, respectively. The branch-site model was used to detect a possible positive selection acting on genotypes B, C and the clade consisting of genotypes A, D, E, H, No34, 64, 91/1. Although the log-likelihood value of the alternative model was found improved when compared with the log-likelihood value of the null model the difference between the two models was not found to be significant ($2\Delta l < 3.84$, $df = 1$, $p > 0.05$).

The runs test implemented in GENECONV revealed 146 possible recombination events, i.e. gene fragments that were sufficiently similar to be suggestive of past gene conversion within *mcvBA1* (123 – 1,379 bp, 12-68 polymorphic sites). All of the genotypes showed at least one putative recombination event with another genotype, in total 8, 85, and 53 events occurred within *P. agardhii*, *P. rubescens*, and between both species, respectively. The most frequent breaking points were from Pos. 526-883 (27), 707-890 (28) and 707-883 (10) of strain No10 (AJ890255). A high number of recombinations (250) was also detected using RDP.

Microcystin genotypes and microcystin variants

Genotypes A, C, D, H, No34, 64, 72 were found to contain exclusively arginine in position 2 of the molecule (Table 1, Fig. 1). Notably the gene region coding for the putative binding pocket, which is responsible for substrate activation during microcystin synthesis (A4-A5, Stachelhaus *et al.* 1999, position 424-735 of strain No10, AJ890255, 312 bp), showed two binding pockets only: The first arginine binding pocket genotype comprised genotype C, while all other strains (A, D, H, No34, 64, 72) containing arginine only formed another binding pocket genotype differing by 4.1% from C, $d_N/d_S = 0.704$, estimated through pairwise comparison by the method of Yang & Nielsen (2000). Strains of genotype B contained microcystin with homotyrosine (tyrosine) and leucine, but never arginine in position 2. Its putative binding pocket differed from the arginine binding pocket of genotype C by 3.2% ($d_N/d_S = 3.34$) and from the other arginine binding pocket of genotype A (D, H, strains No34, 64, 72) by 2.9% ($d_N/d_S = 0.61$). All other genotypes contained arginine and

leucine, and in some cases additional low levels of homotyrosine (G, No31/1, 75, 82). Strains No40, 62, 65, 67, 91/1, 119, 120 were found inactive in MC production, which was partly caused by transposases or gene deletions (Christiansen *et al.* 2006), however, those strains did not differ in sequence compared with active *mcy* genotypes (Table 1).

Distribution of microcystin gene restriction types among strains

Out of the 25 genotypes, 10 restriction types were obtained with *AluI* and 4 restriction types were obtained with *TseI* (Table 3, Fig. 1). In total 11 *AluI*+*TseI* combinations were observed. While some restriction types were found in both strains of *P. rubescens* and *P. agardhii* (VI+II) others were found only in *P. agardhii* (V+IV, VIII+III) or in *P. rubescens* (I+I, II+I, III+I, IX+I). The occurrence of I+I in *P. rubescens* and V+IV in *P. agardhii* was statistically significant (Chi-square test, χ^2 (df=1), $p < 0.05$). The restriction types I+I, II+I, III+I comprised a group of most closely related genotypes of *P. rubescens*, which differed in 0.5% of the base pairs only (15 strains, A, D, H, No34, 64, 91/1).

Restriction types I+I, II+I, III+I, V+IV identified those genotypes with arginine only in position 2 of the molecule. Genotype B lacking arginine was identified by restriction type VI+II. Those genotypes containing arginine/leucine (and homotyrosine) were identified by VI+I, VII+III, VIII+III, IX+I, X+I.

Distribution of microcystin gene restriction types among populations

In order to test the reproducibility of the results on the restriction type proportions from cloning libraries, *mcyBA1* genes from field samples (Mondsee, 9 Dec 03, Wörthersee, 28 Aug 03) were independently amplified by PCR threefold, cloned, and subsequently digested using *AluI* after the re-amplification of 40 clones. The results were reproducible for the dominant restriction types (I, II, IV, VI) as well as the absent restriction types (VII, VIII). The results for the subdominant genotypes were more variable (Table 4).

In total 24 *AluI*+*TseI* combinations were observed. Out of a total of 985 clones, the five most frequently detected restriction types were IV+I (23.8%), I+I (20%), VI+II (13%), X+I (10%), and VI+I (5.1%). All of the other restriction types occurred <5% (Fig. 2). Nine restriction types occurred only once (I+III, I+IV, II+III, III+II, III+III, IX+III, IV+III) or twice (VII+III, X+III). Other restriction types showed new *AluI* or *TseI* profiles that were derived from unknown organisms and that occurred at all of the sampling sites (11.9%). Two restriction types that were observed among the strains of *P. agardhii* (V+IV, VIII+III) were not detected in the field samples.

The distribution of restriction types differed significantly between sites, for example I+I and IV+I occurred frequently in many lakes and were not found in Schwarzensee. In contrast Schwarzensee was dominated by VI+II or X+I. Restriction type VI+II occurred in all of the lakes except for Irrsee and X+I was not found in Afritzersee but showed high frequency in Irrsee and Schwarzensee. In addition Schwarzensee had a significantly lower total number of restriction types when compared to the other lakes (one-way ANOVA, $df = 23$, $p = 0.01$).

In summary, many populations were found to be genetically heterogeneous. Closely located and spatially separated populations were found to differ significantly in *mcyBA1* restriction type composition. Most populations contained *mcyBA1* genotypes indicative of the production of arginine, homotyrosine, and leucine. In contrast genotypes indicative of arginine were absent in Schwarzensee, and that of homotyrosine were absent in Irrsee during the study period.

Discussion

Taxonomic distribution of microcystin genes in *Planktothrix* spp

In this study certain genotypes (C, E) and restriction types (V+IV, I+I) were found to occur exclusively either in one or the other species. On the other hand genotype B occurred in both species and genotypes E, No32, and No39 of *P. agardhii* were found to be the most closely related to the genotypes of *P. rubescens*. This conflict between taxonomy and the phylogenetic assignment of *mcyBA1* genotypes can be best explained by relatively recent recombination events within *mcyBA1*. Homologous recombination has been increasingly recognized as an important force in prokaryotic evolution (Feil 2004). For example, in pathogenic bacteria, recombination has been estimated to change alleles 5-15 fold more frequently than mutation (e.g. Feil *et al.* 2003). For cyanobacteria, Rudi *et al.* (1998) demonstrated the occurrence of recombination for *rbcL* (D-ribulose 1,5-bisphosphate carboxylase-oxygenase) and the less conserved *rbcX* gene with a possible chaperonin-like function. Based on DNA-DNA hybridization studies, Suda *et al.* (2002) concluded that *P. rubescens* and *P. agardhii* differentiated relatively recently and the relatively high genetic similarity may favour the occurrence of recombination (Roberts & Cohan 1993). Typically both species do not occur in the same freshwater system, however, in some lakes both species have been found to co-occur over the course of years (e.g. Blelham Tarn, Davis *et al.* 2003). It would be interesting to know whether this co-occurrence favours the occurrence of interspecific recombination events.

Microevolution of microcystin genes

In recent years a number of recombination events within microcystin genes were reported (Mikalsen *et al.* 2003, Tanabe *et al.* 2004, Kurmayer *et al.* 2005) suggesting that microcystin genes are under continuous modification and re-organisation. While Tanabe *et al.* (2004) observed the recombination of shorter DNA fragments (< 1,000 bp) within *mcyA* (but not within *mcyD*, *mcyG*, *mcyJ*), Kurmayer *et al.* (2005) reported the replacement of whole domains, i.e. the typical N-methyl-dehydroalanine adenylation domain (2,854 bp) of *mcyAA1* was replaced by an adenylation domain without the N-methyl transferase (1,692 bp) resulting in dehydrobutyrine in position 7 instead of the common N-methyl-dehydroalanine. The flanking regions as well as the site of recombination in *mcyAA1* were found to be identical within twelve strains containing the replaced domain implying that this recombination event happened only once (R. Kurmayer, C. Molitor, unpublished). In this study a larger number of short fragments that are indicative of recombination were observed implying that parts of the *mcyB* gene were frequently transferred between lineages of *Planktothrix* spp. It is concluded that the *mcyA* and *mcyB* genes generally show a mosaic structure rather than a bifurcating phylogenetic tree.

The “one ratio” maximum likelihood model estimated $d_N/d_S = 0.19$, which is similar to the ratios calculated by Tanabe *et al.* (2004), i.e. $d_N/d_S = 0.2$ (*mcyA*), 0.14 (*mcyD*), 0.17 (*mcyG*), 0.11 (*mcyJ*). Rantala *et al.* (2004) reported $d_N/d_S < 1$ for *mcyA*, *mcyD*, *mcyE*. An excess of synonymous substitutions over non-synonymous ones indicates that *mcy* genes are subject to purifying selection and mutations affecting the protein sequence are in general deleterious. Notably the genotypes B (indicative of homotyrosine, leucine but no arginine) and C (indicative of arginine only) showed higher d_N/d_S ratios implying either relaxation of purifying selection or positive selection. Averaging d_N/d_S rates over all of the sites of the protein typically underestimates positive selection because the ratio is overwhelmed by the ubiquitous purifying selection. To increase the power of detection branch-site models have been invented to test for positive selection acting on specific sites of specific genotypes (Yang & Nielsen 2002, Zhang *et al.* 2005) or on a subset of sites in the whole phylogenetic tree (Yang *et al.* 2000). Indeed employing site models (Yang *et al.* 2000) 3% of the sites

within the whole *mcvBA1* phylogeny were found positively selected ($d_N/d_S \sim 5.0$, unpublished data). However, no statistically significant support for positive selection in specific branches was found implying that some sites in genotypes B, C rather experienced a relaxation of selective constraints. In the future quantifying the fitness of specific microcystin ecotypes (i.e. Kurmayer *et al.* 2005) under various environmental conditions will help to elucidate whether the observed differences in d_N/d_S ratios resulted from possible adaptive speciation events or from a relaxation of selective constraints.

Divergence of microcystin genotype composition between populations

In this study genotype C (2 strains isolated from the English Lake District, UK) was never detected in field samples in the Alpine lakes. In addition populations were found to differ with regard to the occurrence of specific genotypes as well as to the number of genotypes occurring in total in a specific population. This study is the first that documents homogeneous and more heterogeneous populations of aquatic cyanobacteria existing in closely located yet spatially isolated habitats over the course of several years. According to Horner-Devine (2004) three fundamental processes, the rates of dispersal, speciation, and extinction all contribute to the geographic patterns. There is no reason to assume that *P. rubescens* has fewer dispersal capabilities than other aquatic prokaryotes. Typically *P. rubescens* occurs in depths of 9–12 m (in Mondsee) with low light intensity (i.e. neutral buoyancy in Lake Zürich occurred between 4–8 $\mu\text{mol m}^{-2} \text{s}^{-1}$, Walsby *et al.* 2004). Notably the populations in Schwarzensee and Irrsee showing the lowest genetic diversity observed in this study, also had the lowest numbers of individuals when compared with the populations of high genetic diversity (Mondsee, Wörthersee) as reported previously (Christiansen *et al.* 2006, Table 4). Re-colonization of those habitats by single genotypes after the extinction of the population during winter or other catastrophic events may contribute to the low homogeneity and the high degree of clonality observed in sparse populations. Consequently isolation combined with low numbers of individuals may lead to divergence in microcystin synthesis through genetic drift.

Acknowledgments

We are grateful to Guntram Christiansen and anonymous referees for their helpful comments on earlier versions of the manuscript. We would like to thank Martin Meixner for DNA sequencing. Michi Werndl and Johanna Schmidt provided excellent assistance in the laboratory. We are grateful to the access to samples from Hallwilersee by Arno Stöckli (Abteilung für Umwelt, Kanton Aargau) and from Zürichsee by Judith Blom (University of Zürich, Limnological Station). Günther Bruscek and Karl Mayrhofer (BAW Scharfling, Institut für Gewässerökologie) provided samples from Irrsee and Schwarzensee. This study was supported by the Austrian Science Fund (P18185).

References

- Anisimova M, Nielsen R, Yang Z. Effect of recombination on the accuracy of the likelihood method for detecting positive selection at amino acid sites. *Genetics*. 2003; 164:1229–1236. [PubMed: 12871927]
- Anneville O, Souissi S, Gammeter S, Straile D. Seasonal and inter-annual scales of variability in phytoplankton assemblages: comparison of phytoplankton dynamics in three peri-alpine lakes over a period of 28 years. *Freshwater Biology*. 2004; 49:98–115.
- Barker GLA, Handley BA, Vacharapiyasophon P, Stevens JR, Hayes PK. Allele-specific PCR shows that genetic exchange occurs among genetically diverse *Nodularia* (Cyanobacteria) filaments in the Baltic Sea. *Microbiology*. 2000; 146:2865–2875. [PubMed: 11065365]
- Carmichael WW, Beasley V, Bunner DL, et al. Naming cyclic heptapeptide toxins of cyanobacteria (blue-green algae). *Toxicon*. 1988; 26:971–973. [PubMed: 3149803]
- Challis GL, Ravel J, Townsend CA. Predictive, structure-based model of amino acid recognition by nonribosomal peptide synthetase adenylation domains. *Chemistry and Biology*. 2000; 7:211–224. [PubMed: 10712928]

- Cho J, Tiedje J. Biogeography and degree of endemicity of fluorescent pseudomonas strains in soil. *Applied and Environmental Microbiology*. 2000; 66:5448–5456. [PubMed: 11097926]
- Christiansen G, Fastner J, Erhard M, Börner T, Dittmann E. Microcystin biosynthesis in *Planktothrix*: genes, evolution, and manipulation. *Journal of Bacteriology*. 2003; 185:564–572. [PubMed: 12511503]
- Christiansen G, Kurmayer R, Liu Q, Börner T. Transposons inactivate biosynthesis of the nonribosomal peptide microcystin in naturally occurring *Planktothrix* spp. *Applied and Environmental Microbiology*. 2006; 72:117–123. [PubMed: 16391033]
- Conti E, Stachelhaus T, Marahiel MA, Brick P. Structural basis for the activation of phenylalanine in the non-ribosomal biosynthesis of gramicidin S. *The EMBO Journal*. 1997; 16:4174–4183. [PubMed: 9250661]
- Davis PA, Dent M, Parker J, Reynolds CS, Walsby AE. The annual cycle of growth rate and biomass change in *Planktothrix* spp. in Blelham Tarn, English Lake District. *Freshwater Biology*. 2003; 48:852–867.
- Fastner J, Erhard M, Carmichael WW, et al. Characterization and diversity of microcystins in natural blooms and strains of the genera *Microcystis* and *Planktothrix* from German freshwaters. *Archiv für Hydrobiologie*. 1999; 145:147–163.
- Feil E. Small change: keeping pace with microevolution. *Nature Reviews*. 2004; 2:483–495.
- Feil E, Cooper J, Grundmann H, et al. How clonal is *Staphylococcus aureus*? *Journal of Bacteriology*. 2003; 185:3307–3316. [PubMed: 12754228]
- Felsenstein J. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics*. 1989; 5:164–166.
- Finlay B. Global dispersal of free-living microbial eukaryote species. *Science*. 2002; 296:1061–1063. [PubMed: 12004115]
- Henriksen P, Moestrup Ø. Seasonal variations in microcystin contents of Danish cyanobacteria. *Natural Toxins*. 1997; 5:99–106. [PubMed: 9285913]
- Horner-Devine M, Carney K, Bohannon J. An ecological perspective on bacterial biodiversity. *Proceedings of the Royal Society London Series B*. 2004; 271:113–122.
- Jongman, R.; Ter Braak, C.; Van Tongeren, O. Data analysis in community and landscape ecology. Cambridge University Press; Cambridge: 1995.
- Kimura, M. The neutral theory of molecular evolution. Cambridge University Press; Cambridge: 1983.
- Komarek J. Planktic oscillatorial cyanoprokaryotes (short review according to combined phenotype and molecular aspects). *Hydrobiologia*. 2003; 502:367–382.
- Kurmayer R, Christiansen G, Chorus I. The abundance of microcystin-producing genotypes correlates positively with colony size in *Microcystis* and determines its microcystin net production in Lake Wannsee. *Applied and Environmental Microbiology*. 2003; 69:787–795. [PubMed: 12570996]
- Kurmayer R, Christiansen G, Fastner J, Börner T. Abundance of active and inactive microcystin genotypes in populations of the toxic cyanobacterium *Planktothrix* spp. *Environmental Microbiology*. 2004; 6:831–841. [PubMed: 15250885]
- Kurmayer R, Christiansen G, Gumpenberger M, Fastner J. Genetic identification of microcystin ecotypes in toxic cyanobacteria of the genus *Planktothrix*. *Microbiology*. 2005; 151:1525–1533. [PubMed: 15870462]
- Kurmayer R, Dittmann E, Fastner J, Chorus I. Diversity of microcystin genes within a population of the toxic cyanobacterium *Microcystis* spp. in Lake Wannsee (Berlin, Germany). *Microbial Ecology*. 2002; 43:107–118. [PubMed: 11984633]
- Lawton LA, Edwards C, Codd GA. Extraction and high-performance liquid chromatographic method for the determination of microcystins in raw and treated waters. *The Analyst*. 1994; 119:1525–1530. [PubMed: 7943743]
- Lodders N, Stackebrandt E, Nübel U. Frequent genetic recombination in natural populations of the marine cyanobacterium *Microcoleus chthonoplastes*. *Environmental Microbiology*. 2005; 7:434–442. [PubMed: 15683403]
- Luukkainen R, Sivonen K, Namikoshi M, Färdig M, Rinehart KL, Niemelä SI. Isolation and identification of eight microcystins from thirteen *Oscillatoria agardhii* strains and structure of a new microcystin. *Applied and Environmental Microbiology*. 1993; 59:2204–2209. [PubMed: 8357254]

- Marahiel MA, Stachelhaus T, Mootz HD. Modular peptide synthetases involved in nonribosomal peptide synthesis. *Chemical Reviews*. 1997; 97:2651–2673. [PubMed: 11851476]
- Martin D, Rybicki E. RDP: detection of recombination amongst aligned sequences. *Bioinformatics*. 2000; 16:562–563. [PubMed: 10980155]
- Mikalsen B, Boison G, Skulberg OM, et al. Natural variation in the microcystin synthetase operon *mcyABC* and impact on microcystin production in *Microcystis* strains. *Journal of Bacteriology*. 2003; 185:2774–2785. [PubMed: 12700256]
- Palys T, Nakamura LK, Cohan FM. Discovery and classification of ecological diversity in the bacterial world: the role of DNA sequence data. *International Journal of Systematic and Evolutionary Bacteriology*. 1997; 47:1145–1156.
- Papke RT, Ramsing NB, Bateson MM, Ward DM. Geographical isolation in hot spring cyanobacteria. *Environmental Microbiology*. 2003; 5:650–659. [PubMed: 12871232]
- Papke RT, Ward DM. The importance of physical isolation to microbial diversification. *FEMS Microbiology Ecology*. 2004; 48:293–303. [PubMed: 19712299]
- Rantala A, Fewer DP, Hisbergues M, et al. Phylogenetic evidence for the early evolution of microcystin synthesis. *Proceedings of the National Academy of Sciences of the USA*. 2004; 101:568–573. [PubMed: 14701903]
- Rippka R. Isolation and purification of cyanobacteria. *Methods in Enzymology*. 1988; 167:3–27. [PubMed: 3148836]
- Roberts M, Cohan F. The effect of DNA sequence divergence on sexual isolation in *Bacillus*. *Genetics*. 1993; 134:401–408. [PubMed: 8325477]
- Roberts M, Cohan F. Recombination and migration rates in natural populations of *Bacillus subtilis* and *Bacillus mojavensis*. *Evolution*. 1995; 49:1081–1094.
- Rücker J, Wiedner C, Zippel P. Factors controlling the dominance of *Planktothrix agardhii* and *Limnithrix redekei* in eutrophic shallow lakes. *Hydrobiologia*. 1997; 342/343:107–115.
- Rudi K, Skulberg OM, Jakobsen KS. Evolution of cyanobacteria by exchange of genetic material among phyletically related strains. *Journal of Bacteriology*. 1998; 180:3453–3461. [PubMed: 9642201]
- Sambrook, J.; Fritsch, EF.; Maniatis, T. *Molecular cloning-A laboratory manual* second edition. Cold Spring Harbor Laboratory Press; New York: 1989.
- Sawyer, S. GENECONV: A computer package for the statistical detection of gene conversion. Department of Mathematics, Washington University; St. Louis: 1999. Distributed by the author available at <http://www.math.wustl.edu/~sawyer>
- Stachelhaus T, Mootz HD, Marahiel MA. The specificity-conferring code of adenylation domains in nonribosomal peptide synthetases. *Chemistry and Biology*. 1999; 6:493–505. [PubMed: 10421756]
- Suda S, Watanabe MM, Otsuka S, et al. Taxonomic revision of water-bloom-forming species of oscillatoroid cyanobacteria. *International Journal of Systematic and Evolutionary Microbiology*. 2002; 52:1577–1595. [PubMed: 12361260]
- Tanabe Y, Kaya K, Watanabe MM. Evidence for recombination in the microcystin synthetase (*mcy*) genes of toxic cyanobacteria *Microcystis* spp. *Journal of Molecular Evolution*. 2004; 58:633–641. [PubMed: 15461420]
- Tillett D, Dittmann E, Erhard M, et al. Structural organization of microcystin biosynthesis in *Microcystis aeruginosa* PCC7806: an integrated peptide-polyketide synthetase system. *Chemistry and Biology*. 2000; 7:753–764. [PubMed: 11033079]
- Walsby AE, Ng G, Dunn C, Davis PA. Comparison of the depth where *Planktothrix rubescens* stratifies and the depth where the daily insolation supports its neutral buoyancy. *New Phytologist*. 2004; 162:133–145.
- Whitaker RJ, Grogan DW, Taylor JW. Geographic barriers isolate endemic populations of hyperthermophilic archaea. *Science*. 2003; 301:976–978. [PubMed: 12881573]
- Yang Z. PAML: a program package for phylogenetic analysis by maximum likelihood. *Computer Applications in BioSciences*. 1997; 13:555–556.
- Yang Z. Likelihood ratio tests for detecting positive selection and application to primate lysozyme evolution. *Molecular Biology and Evolution*. 1998; 15:568–573. [PubMed: 9580986]

- Yang Z, Nielsen R. Estimating synonymous and nonsynonymous substitution rates under realistic evolutionary models. *Molecular Biology and Evolution*. 2000; 17:32–43. [PubMed: 10666704]
- Yang Z, Nielsen R, Goldman N, Krabbe Pedersen A-M. Codon-substitution models for heterogeneous selection pressure at amino acid sites. *Genetics*. 2000; 155:431–449. [PubMed: 10790415]
- Yang Z, Nielsen R. Codon-substitution models for detecting molecular adaptation at individual sites along specific lineages. *Molecular Biology and Evolution*. 2002; 19:908–917. [PubMed: 12032247]
- Zhang J, Nielsen R, Yang Z. Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Molecular Biology and Evolution*. 2005; 22:2472–2479. [PubMed: 16107592]
- Zöfel, P. Statistik in der Praxis, 3. Auflage. Gustav Fischer Verlag; Stuttgart, Jena: 1992.

Fig. 1.

Phylogenetic tree based on ML analysis from *mcyBA1* sequences (1,451 bp) from 49 *Planktothrix* spp. strains. Strains assigned to *P. agardhii* are underlined. Capitals A-H in the superscript indicate the genotype groups (consisting of 2-11 strains) as indicated in Table 1. Genotype B consisted of both *P. rubescens* and *P. agardhii*. For each strain the corresponding restriction profile (*AluI*, *TseI*) and the occurrence of amino acids in position 2 of the microcystin molecule is indicated. The significant bootstrap percentages were obtained from 100 pseudo replicates. The two numbers shown in boxes are the maximum-likelihood estimates of the numbers of nonsynonymous/synonymous substitutions for the entire *mcyBA1* along that branch (only ratios higher than the background ratio of the “one-ratio” model ($d_N/d_S = 0.19$) are shown, see text).

Fig. 2.

Proportion of *AluI*+*TseI* restriction types obtained from amplified and cloned *mcvBA1* PCR products from eight different *Planktothrix* spp. populations for three sampling dates. For each sample, 40 clones of *mcvBA1* PCR products were analysed and the percentage of a specific restriction type was calculated. The proportion for the total population and for the 49 sequenced strains is provided in the lower two graphs. Note that the scales of the y-axes are different. Significant differences of the proportions of restriction profiles between the eight lakes are marked by crosses at the top (xxx, $p < 0.001$, xx, $p < 0.01$, ns, not significant). The mean $\pm$ 1 SE number of the total number of restriction types found in a population is given in parentheses. Superscripts a, ab, b indicate homogeneous subsets whose highest and lowest means in the number of different restriction types are not significantly different ($p > 0.05$).

Table 1

Variation of amino acids in position 2 of microcystin molecules (Arg, arginine, Hty, homotyrosine, Leu, leucine) produced among 49 strains of *P. rubescens* (rub) and *P. agardhii* (aga). Strains are ordered according to the proportion of each of the structural variants produced (mean $\pm$ 1SE). The number of MC measurements using HPLC analysis is indicated for each strain. Strains marked by asterisks have been shown to be inactive in MC production (Christiansen *et al.* 2006). The corresponding *AluI* (I-X) and *TseI* (I-IV) restriction types, and the sequence accession numbers of *mcvBA1* are indicated. Superscripts A-H identify identical genotypes. Strains have been isolated during the study by Kurmayer *et al.* (2004). Country codes (ISO format). SAG Culture Collection of Algae (Göttingen, Germany), PCC Pasteur Culture Collection (Institute of Pasteur, Paris, France), CCAP Culture Collection of Algae and Protozoa (Windermere, UK).

Microcystin data						Restr. type		Genotype		Strain data		
Amino acid in pos. 2	Arginine	Homotyrosine	Leucine	undetermined	N of MC measurements	AluI	TseI	PC-IGS acc. no.	mcvBA1 acc. no. ²	Species	Strain No	origin
Arg	99±1		1±1	5	1	I	I		AJ749276 ^A	rub	3	Mondsee (AT)
	100			2	1	I	I		AJ890255 ^A	rub	10	Mondsee (AT)
				2	1	I	I		AJ890263 ^B	rub	40*	Mondsee (AT)
				4	1	I	I		AJ890264	rub	91/1*	Mondsee (AT)
	99±1		1±1	2	1	I	I		AJ890256 ^A	rub	99	Mondsee (AT)
	97		3	2	1	I	I		AJ890257 ^A	rub	109	Mondsee (AT)
	97±1		3±2	4	1	I	I		AJ749282 ^A	rub	111	Mondsee (AT)
	95±2		5±2	2	1	I	I		AJ890258 ^A	rub	113	Mondsee (AT)
				2	1	I	I		AJ890259 ^A	rub	119*	Mondsee (AT)
	97±1		3±1	4	1	I	I	AJ558156	AJ890260	rub	34	Ammersee (DE)
	99±1		1±1	2	1	I	I		AJ890261 ^A	rub	81	Ammersee (DE)
	97±1		3±1	2	1	I	I		AJ890262 ^B	rub	86	Ammersee (DE)
				2	II	I	I	AJ558153	AJ890265 ^D	rub	62*	Irsee (AT)
				3	II	I	I		AJ890266 ^D	rub	65*	Irsee (AT)
	100±1		1±1	4	III	I	I		AJ749277	rub	64	Wörthersee (AT)
	100			3	V	IV	IV	AJ558160	AJ749272 ^C	aga	CCAP1459/11A	L. Windermere (UK)
	100			3	V	IV	IV	AJ558159	AJ749274 ^C	aga	CCAP1459/21	Esth. Water (UK)
	100			2	IV	I	I		AJ890267	rub	72	Zürichsee (CH)
Hty/Leu	42±5	56±6	2	2	VI	II	II		AJ863131 ^A	rub	21-	Figur (AT)
	54±3	44±1	1±1	2	VI	II	II		AJ890275 ^B	rub	21/2	Figur (AT)
	93±7 ¹		3±3	2	VI	II	II		AJ890276 ^B	rub	13	Schwarzensee (AT)
	95±5 ¹		3±4	2	VI	II	II		AJ890277 ^B	rub	18	Schwarzensee (AT)
	94±6 ¹		3±2	2	VI	II	II		AJ890278 ^B	rub	59	Schwarzensee (AT)
	88 ¹		13	2	VI	II	II		AJ890279 ^B	rub	60	Schwarzensee (AT)
	86±1 ¹	14±2	5	VI	II	II	AJ558152	AJ749278 ^B	rub	80	Schwarzensee (AT)	
	69±3	32±4	3	VI	II	II	AJ558158	AJ749273 ^B	aga	CCAP1459/16	Blelham Tarn (UK)	
	75±9	23±7	2±2	2	VI	II	II		AJ863133 ^B	aga	CCAP1459/17	Blelham Tarn (UK)
	90	10		2	VI	II	II		AJ863134 ^B	aga	CCAP1459/31	White Lough (UK)
	75±3	27±2	3	VI	II	II		AJ749275 ^B	aga	CCAP1460/5	L. Kasumigaura (JP)	
	Arg/Leu	90	9	1±1	5	IV	I	I		AJ749269	rub	39
89		8±3		2	IV	I	I		AJ890268	rub	42	Wörthersee (AT)
91±3		9±3		2	IV	I	I		AJ890269	rub	101	Mondsee (AT)
88±4		12±4		2	IV	I	I		AJ890270	rub	118	Mondsee (AT)
70±2		30±2		5	VI	I	I	AJ558147	AJ749284	rub	CCAP1459/30	Plöner See (DE)
87±4		13±4		2	VI	I	I	AJ558150	AJ890271	rub	CCAP1459/24	L. Ören (SE)
80±3		19±2	1	3	VII	III	III	AJ558137	AJ749271	rub	SAG6.89	Pinflsee, Plön (DE)
95±1		5±1		4	VIII	III	III	AJ558154	AJ749283	rub	PCC7821	L. Gjersjøen (NO)
89±1		11	1±1	4	VIII	III	III		AJ749268	aga	32	Wannsee (DE)
94		2	4	2	IX	I	I		AJ890280 ^F	rub	77	Wörthersee (AT)
93±2		3±1	5±1	2	IX	I	I		AJ890281 ^F	rub	89	Wörthersee (AT)
				2	IX	I	I		AJ890282 ^F	rub	120*	Mondsee (AT)
95±1		4±1		5	X	I	I		AJ749270 ^E	aga	79	L. Arreso (DK)
99		1		4	X	I	I		AJ749281	rub	108	Irsee (AT)
90±2		9±2	1±1	7	X	I	I	AJ558135	AJ441056 ^E	aga	CYA126/8	L. Langsjön (FI)
Arg/Hty/Leu	94±2	1±1	5±1	2	VI	I	I	AJ558151	AJ890272 ^G	rub	46	Zürichsee (CH)
				2	VI	I	I		AJ890273 ^G	rub	67*	Wörthersee (AT)
	91±3	2	7±3	2	VI	I	I		AJ890274	rub	75	Zürichsee (CH)
	83±2	1±1	14±3	2±1	4	VII	III		AJ749267	aga	31/1	Wannsee (DE)
	95±1	1±0	5±1		6	X	I		AJ749279	rub	82	Ammersee (DE)

¹ New microcystins with Hty/Tyr in position 2 of the microcystin molecule (R. Kurmayer, W. Yoshida, K. Ishida, T. Hemscheidt, unpublished)

² Accession numbers AJ749267-AJ749284, AJ863131-AJ863134 have been published by Kurmayer *et al.* (2005).

Table 2

Morphometric and limnological characteristics of lakes sampled for *Planktothrix rubescens*. Trophic data (M, mesotrophic, O, oligotrophic) are yearly minimum-mean-maximum value (sample size) of the year 2001 from the Kärntner Institut für Seenforschung, Carinthia (Afritzersee, Wörthersee), Bayrisches Landesamt, Wielenbach, Germany (Ammersee), Department for Environment, Kanton Aargau (Hallwilersee), Federal Agency for Water Management, Scharfling, Upper Austria (Irrsee, Schwarzensee) and Zurich Water Supply (Zürichsee). Chlorophyll *a* values were integrated over the watercolumn. As inferred from microscopical inspection of net samples (30 µm mesh size) the abundance of *P. rubescens* is given: D, dominant; A, abundant, R, rare. M a SL, meter above sea level.

Lake	N(°)	E(°)	M a SL	Area (km ²)	Trophic status	Secchi depth (m)	Chlorophyll <i>a</i> (µg L ⁻¹)	Abundance <i>Planktothrix</i>
Afritzer See, AT	46°44	13°46	750	0.5	M	3.3-4.0-5.2 (4)	2.4-4.4-6.8 (4)	A
Ammersee, DE	47°16	11°4	533	47	M	1.6-4.4-6.5 (11)	1.5-5.1-9 (11)	A
Hallwilersee, CH	47°20	8°10	449	10	M	1-2.7-5 (12)	8.8-18.9-30 (12)	D
Irrsee, AT	47°56	13°19	553	3.6	M	2.2-5.1-8.2 (12)	1.4-2.8-4.3 (7)	R
Mondsee, AT	47°48	13°22	481	14.2	M	2.4-3.8-5 (24)	1.8-4.3-9.6 (24)	A
Schwarzensee, AT	47°45	13°30	716	0.5	O	5.5-6.5-7.5 (3)	0.5-0.9-1.5 (3)	R
Wörthersee, AT	46°36	14°3	440	19.4	M	3.5-4.9-7.5 (7)	5.4-9-13.3 (7)	D
Zürichsee, CH	47°22	8°32	406	65	M	2.3-3.9-5.9(18)	5.9-5-15.7 (12)	D

Table 3

Schematic representation of *AclI* and *TseI* restriction profiles within *mycBA1* (1,693 bp) of *Planktothrix* spp. The size of the restriction fragments in bp was calculated from the sequences.

Fragments (bp)	<i>AclI</i>							<i>TseI</i>			
	I	II	III	IV	V	VI	VII	VIII	IX	X	I II III IV
1693											—
865											—
828											—
798											—
735											—
328	—	—	—	—	—	—	—	—	—	—	
289	—	—	—	—	—	—	—	—	—	—	
272	—	—	—	—	—	—	—	—	—	—	
263	—	—	—	—	—	—	—	—	—	—	
243				—	—	—	—	—	—	—	
225					—	—	—	—	—	—	
208		—			—	—	—	—	—	—	
201	—	—	—	—	—	—	—	—	—	—	
179					—	—	—	—	—	—	
132	—	—	—	—	—	—	—	—	—	—	
114			—			—		—		—	
106	—			—					—		
102	—			—					—		
94			—			—		—		—	
93											—
67											—
47							—	—	—	—	
46							—	—	—	—	
20				—	—	—	—	—	—	—	

Table 4

Proportion of restriction types (I-X, new, new restriction types) of *mcvBA1* obtained through *AclI* digestion after three independent PCR amplifications from field samples, cloning of the PCR product, and PCR-reamplification of forty randomly selected white clones from the respective clone library (1-3).

Clone library	I	II	III	IV	V	VI	VII	VIII	IX	X	new
Mondsee, 9 Dec 03											
1	35	8	8	10	3	18	0	0	10	3	8
2	16	22	8	22	3	5	0	0	3	3	19
3	5	33	0	23	5	10	0	0	8	3	13
Mean ± 1SE	19±9	21±7	5±3	18±4	3±1	11±4	0	0	7±2	3±0	13±3
Wörthersee, 28 Aug 03											
1	30	5	3	20	5	13	0	0	3	8	15
2	26	31	10	7	0	17	0	0	0	0	10
3	26	5	5	14	0	19	0	0	12	2	19
Mean ± 1SE	27±1	14±9	6±2	14±4	2±2	16±2	0	0	5±4	3±2	14±3