Published in final edited form as:

Toxicol Appl Pharmacol. 2009 February 15; 235(1): 1–9. doi:10.1016/j.taap.2008.11.001.

Paraoxonase 1 (PON1) Status and Substrate Hydrolysis^a

Rebecca J. Richter, Gail P. Jarvik, and Clement E. Furlong

Departments of Medicine–Division of Medical Genetics, and Genome Sciences University of Washington, Seattle, WA 98195-7720

Abstract

Paraoxonase 1 (PON1) hydrolyzes a number of organophosphorus (OP) compounds including insecticides and nerve agents. The *in vivo* efficacy of PON1 to protect against a specific OP exposure depends on the catalytic efficiency of hydrolysis. The Q192R polymorphism affects the catalytic efficiency of hydrolysis of some substrates and not others. While PON1_{R192} hydrolyzes paraoxon approximately 9-times as efficiently as PON1_{Q192}, the efficiency is insufficient to provide *in vivo* protection against paraoxon/parathion exposure. The two PON1₁₉₂ alloforms have nearly equivalent but higher catalytic efficiencies for hydrolyzing diazoxon (DZO) and provide equivalent *in vivo* protection against DZO exposures. On the other hand, PON1_{R192} is significantly more efficient in hydrolyzing chlorpyrifos oxon (CPO) than PON1_{Q192} and provides better protection against CPO exposure. Thus, for some exposures it is only the level of plasma PON1 that is important, whereas for others it is both plasma level and the PON1₁₉₂ alloform(s) present in plasma that are important. In no case is the plasma level of PON1 unimportant, provided that the catalytic efficiency is sufficient to protect against the exposure. Two-substrate enzyme assay/analysis protocols that reveal both PON1 plasma levels and PON1₁₉₂ phenotype (QQ; QR; RR) are designed to optimize the separation of PON1₁₉₂ phenotypes; however, they have not been optimized for evaluating *in vivo* rates of OP detoxication. This study describes the adaptation of a non-OP, two-substrate determination of PON1 status to the conversion of the PON1 status data to physiologically relevant rates of DZO and CPO detoxication. Conversion factors were generated for rates of hydrolysis of different substrates.

Keywords

paraoxonase 1; PON1 polymorphism; chlorpyrifos; chlorpyrifos oxon; diazinon; diazoxon; pharmacogenetics

Introduction

Genetic variability in human plasma paraoxonase 1 (PON1) activity (POase) has been of interest since the first reports of a polymorphic distribution of POase activity in human

^aSome of the data described here was presented at two recent conferences: 1) Medical Chemical Defense Bioscience Review, Hunt Valley Inn, Hunt Valley, MD, June 5th, 2008, and 2) The Third International Conference on Paraoxonases, UCLA, Los Angeles, CA, September 7th-10th, 2008.

© 2008 Elsevier Inc. All rights reserved.

Corresponding Author Dr. Clement E. Furlong Medical Genetics Box 357720 University of Washington Seattle, WA 98195-7720 Telephone: 206-543-1193 Fax: 206-543-3050 clem@u.washington.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

plasma samples (reviewed in Diepgen and Geldmacher v. Mallinckrodt, 1986). Over the years, many different assays have been developed to explore the PON1 activity polymorphism in human populations (reviewed in Diepgen and Geldmacher-von Mallinckrodt, 1986; Ortigoza-Ferado *et al.*, 1984). The observed polymorphism of high vs. low POase activity suggested that individuals with high plasma POase activity would be more resistant to paraoxon (PO)/parathion exposures than individuals with low activity. This concept persisted for over four decades. It was not until human PON1 was purified (Gan *et al.*, 1991; Furlong *et al.*, 1991), the human PON1 cDNA cloned and characterized (Hassett *et al.*, 1991), the molecular basis of the high/low activity polymorphism identified (Humbert *et al.*, 1993; Adkins *et al.*, 1993) and mouse model systems developed, that the toxicological significance of the genetic variability of plasma POase was understood (Li *et al.*, 1993, 1995, 2000; Shih *et al.*, 1998).

Characterization of the human cDNA sequences revealed two coding region polymorphisms, M55L and Q192R (Hassett *et al.*, 1991). Further characterization of the polymorphisms demonstrated that it was the Q192R polymorphism that was responsible for high vs. low POase activity (Humbert *et al.*, 1993; Adkins *et al.*, 1993). Eckerson *et al.* (1983) developed a two-substrate assay/analysis protocol that plotted rates of PO hydrolysis against rates of phenyl acetate hydrolysis. This analysis showed that it was possible to resolve phenotypes of high vs. low POase activity and at the same time demonstrated that within a given phenotype, there was a wide variability of activity in a given population. Davies *et al.* (1996) made use of this approach in characterizing additional substrates, where they plotted the rates of hydrolysis of a number of different OP compounds and phenyl acetate against rates of PO hydrolysis. The experiments of Davies *et al.* revealed that plots of diazoxonase activity (DZOase) vs. POase for plasma samples from a population provided a complete resolution of the three PON1₁₉₂ phenotypes in the population (QQ, QR and RR). In further refining the two-substrate assay/analysis protocols, we found that the PON1_{R192} alloform was much more sensitive to salt inhibition than the PON1_{Q192} alloform, explaining the clear separation of the PON1 phenotypes (Richter and Furlong, 1999). Unfortunately, a number of investigators have interpreted the lower activity of PON1_{R192} at high salt concentration to indicate that PON1_{R192} homozygotes would be more sensitive to diazinon (DZS)/diazoxon (DZO) exposures than would be PON1_{Q192} homozygotes or PON1₁₉₂ heterozygotes. PON1_{R192} protects as well as PON1_{Q192} against DZO exposure, if not slightly better. Even though PON1_{R192} has a lower V_{max} , it has a higher affinity (lower K_m) for DZO such that the catalytic efficiencies of the two PON1₁₉₂ alloforms are nearly the same (Li *et al.*, 2000).

Experiments with *PON1* null (*PON1*^{-/-}) mice allowed for a determination of the ability of the two purified human PON1₁₉₂ alloforms to protect against OP exposures when injected into the *PON1*^{-/-} mice (Li *et al.*, 2000). The results showed that injected PON1_{R192} protected better than PON1_{Q192} against CPO exposure, and both PON1₁₉₂ alloforms protected nearly equivalently against DZO exposure, whereas, surprisingly, neither protected against PO exposure. The determination of the *in vitro* catalytic efficiencies of each PON1₁₉₂ alloform provided a clear explanation for these observations. The catalytic efficiency of PON1_{R192} for hydrolysis of CPO was better than that of PON1_{Q192}, explaining the increased protection afforded by injection of purified human PON1₁₉₂ into the *PON1*^{-/-} mice. The equivalent catalytic efficiencies of the two PON1₁₉₂ alloforms for hydrolysis of DZO were consistent with the equivalent protection provided by injecting each of these alloforms into *PON1*^{-/-} mice prior to DZO exposure. The lack of protection against PO exposure by injection of purified PON1_{R192} into the *PON1*^{-/-} mice was also explained by the catalytic efficiency data. Despite the fact that the catalytic efficiency of PON1_{R192} was approximately 9-times that of PON1_{Q192}, it was still too low to protect against *in vivo* exposures, in agreement with earlier observations in rats (Chambers *et al.*, 1994; Pond *et al.*, 1995). Experiments on the OP sensitivity of *PON1* transgenic (*PON1*-tg) mice provided

additional confirmatory evidence for the importance of both PON1 level and catalytic efficiency in protecting against OP exposure. Cole *et al.* (2005) showed that *PON1*-tg mice expressing human PON1_{R192} in place of mouse PON1 were significantly more resistant to CPO exposure than were PON1-tg mice that expressed PON1_{Q192} in place of mouse PON1.

These experiments by Li *et al.* (2000) and Cole *et al.* (2005) clearly pointed out the importance of catalytic efficiency of hydrolysis in determining whether PON1 could protect against a given OP exposure. In addition to considering the catalytic efficiency of each PON1₁₉₂ alloform, it is very important to also consider the level of plasma PON1. For chlorpyrifos (CPS)/CPO exposures, both the PON1₁₉₂ phenotype as well as PON1 level are important whereas for DZS/DZO exposures, it is primarily the level of PON1 that is important. In no case is the level of plasma PON1 unimportant. For PO exposures (Li *et al.*, 2000), and nerve agent exposures (Rochu *et al.*, 2007; Yamada *et al.*, 2001) the existing data indicate that PON1 is not effective in protecting against the exposure.

Levels of plasma PON1 are influenced by polymorphisms in the promoter region. Data from three different laboratories provided information on the promoter region polymorphisms that contribute to the large variability in PON1 levels observed among individuals with the same PON1₁₉₂ phenotypes (Brophy *et al.*, 2001a, 2001b; Leviev and James, 2000; Suehiro *et al.*, 2000). The C-108T polymorphism, that occurs in an Sp1 transcription factor binding site (Deakin *et al.*, 2003), has a significant effect on PON1 expression with the T-108 allele expressing less PON1 than the C-108 allele.

Some concern has been expressed about the significance of measuring rates of DZO hydrolysis under non-physiological pH and salt concentration (O'Leary *et al.*, 2005). The data reported here address this question and at the same time provide factors for converting data generated from PON1 status analyses either with the substrate pair DZO/PO (Davies *et al.*, 1996) or a recently reported non-toxic-OP substrate pair 4-(chloromethyl)phenyl acetate/phenyl acetate (Richter *et al.*, submitted to *Circulation: Cardiovascular Genetics*) to physiologically relevant rates of OP hydrolysis. The later assay was developed to provide laboratories not equipped to work with highly toxic OP substrates with a means of establishing an individual's PON1 status. Determining physiologically relevant rates of OP detoxication for DZO and CPO for individuals is important, since these rates can vary by 65- to 164-fold, respectively, in populations if both infants and adults are considered (Furlong *et al.*, 2006).

Interest in PON1 status has expanded well beyond the toxicology community since low PON1 status has been associated with cardiovascular (reviewed in James, 2006) and other diseases (reviewed in Mackness *et al.*, 2002; Marsallich *et al.*, 2008). PON1 is also involved in the activation (Tougou *et al.*, 1998) and inactivation (Biggadike *et al.*, 2000) of specific drugs and the inactivation of bacterial quorum-sensing factors (Ozer *et al.*, 2005; Stoltz *et al.*, 2008).

Methods

Study Population

The plasma samples used for this study came from a University of Washington IRB-approved project investigating the role of PON1 in vascular disease. Informed consent was obtained from all study participants.

Sampling

Blood samples were obtained by venipuncture and were drawn into lithium-heparin tubes. Cells were separated from plasma by centrifugation for 15 min at $1800 \times g$ and the plasma samples were stored at -80°C until assayed.

Chemicals

CMPA [4-(Chloromethyl)phenyl acetate], phenyl acetate and other reagent chemicals were obtained from Sigma Chemical Co. (St. Louis, MO). Paraoxon, chlorpyrifos oxon and diazoxon were obtained from Chem Service (West Chester, PA). The structures of the compounds used in this study are shown in Figure 1.

Precautions

Paraoxon, chlorpyrifos oxon and diazoxon are highly toxic organophosphate compounds. They are potent inhibitors of acetylcholinesterases and other important esterases and lipases, and may cause life-threatening neurologic symptoms if ingested or if they come in contact with skin. Double gloves (nitrile), a laboratory coat, and a chemical hood are used as personal protection when handling these chemicals. Organophosphate solutions are treated with alkali solution (i.e., sodium hydroxide) and broken down to non-toxic products before disposal.

Enzyme Assays

All enzyme activity assays were carried out in a SPECTRAMax[®] PLUS Microplate Spectrophotometer (Molecular Devices, Sunnyvale, CA) as described previously (Richter and Furlong, 1999). The assay values were corrected for path-length using software provided with the microplate reader. For continuous monitoring of hydrolysis products that absorb in the UV range, ultraviolet transparent 96-well microplates from Costar (Cambridge, MA) were used. For monitoring of hydrolysis products that absorb in the visible range, standard flat bottom 96-well microplates from Greiner One (Monroe, NC) were used. For efficiency of loading the microplate wells, a multi-channel pipette (Matrix, Hudson, NH) was used. All assays were run in triplicate. Only initial linear rates of hydrolysis were used for calculations, with results normalized using the path-length correction software provided by the manufacturer. Hydrolysis of paraoxon was monitored at 405 nm and the activity expressed in units/L (U/L) using the molar extinction coefficient $18 \text{ mM}^{-1}\text{cm}^{-1}$ for p-nitrophenol. Rates of DZO hydrolysis were monitored at 270 nm in UV transparent microplates and the activity was expressed as U/L based on the molar extinction coefficient of $3 \text{ mM}^{-1}\text{cm}^{-1}$ for the DZO hydrolysis product, 2-isopropyl-4-methyl-6-hydroxypyrimidine (IMHP). Arylesterase activity (hydrolysis of phenyl acetate) was monitored at 270 nm. Activities were expressed in Units/ml, based on the molar extinction coefficient of $1.31 \text{ mM}^{-1}\text{cm}^{-1}$ for phenol. Samples with replicate values that differed by more than 10% were re-assayed. The assay conditions with respect to salt concentration and pH values are noted in the figure legends.

Results

The aim of this study was to provide protocols for converting PON1 status data generated with non-OP substrates to physiologically relevant rates of DZO and CPO hydrolysis for individuals with any of the three PON1₁₉₂ phenotypes (QQ; QR; RR). A second aim was to provide conversion factors that would allow the comparison of data generated with the new PON1 status protocol with data generated in earlier studies with other substrates.

Two protocols for determining PON1 status are shown in Figure 2. Our earlier studies (Richter and Furlong, 1999) demonstrated that a two-substrate assay/activity plot with rates

of DZO hydrolysis plotted against rates of paraoxon hydrolysis revealed an individual's functional PON1 status—plasma PON1 activity level and PON1₁₉₂ alloform(s) present in their plasma. A comparison of the functional PON1 status protocol with SNP analyses revealed PON1 mutations that were not detectable by SNP analyses alone (Jarvik *et al.*, 2003a).

The PON1 two-substrate/activity plot population distribution using the substrate pair DZO/PO is shown in Figure 2A. A recently developed protocol using the non-OP substrate pair phenyl acetate (assayed at high salt) and CMPA is shown in Figure 2B (Richter *et al.*, submitted to *Circulation: Cardiovascular Genetics*). Both protocols provided a clear resolution of the three PON1₁₉₂ phenotypes (functional genotypes).

The PON1_{192Q/R} alloforms can hydrolyze substrates with widely differing catalytic efficiencies (Fig. 1; Davies *et al.*, 1996; Li *et al.*, 2000). To develop protocols for converting rates of hydrolysis of one substrate to another, it was necessary to characterize substrate activity relationships separately for each PON1₁₉₂ phenotype. Since concern has been expressed regarding the usefulness of measuring rates of DZO hydrolysis under non-physiological conditions of salt and pH value (O'Leary *et al.*, 2005), we first determined the K_m values for DZO hydrolysis by each of the PON1₁₉₂ alloforms at physiological pH, salt concentration (150 mM NaCl) and temperature (37 °C) (Fig. 3A, B). As we had observed previously when we determined K_m the values for DZO hydrolysis at high salt and pH 8.5, the PON1_{R192} alloform had a higher affinity for DZO (K_m = 0.27 mM; Fig. 3A) than the PON1_{Q192} alloform (K_m = 0.52 mM; Fig 3B). These values were determined using plasma samples phenotyped as homozygous for either PON1_{R192} or PON1_{Q192}, to determine the properties of the PON1₁₉₂ alloforms while associated with high density lipoprotein (HDL).

Since PON1 is also important in modulating exposures to CPS/CPO, we also determined the K_m values for CPO hydrolysis (CPOase) by plasma from PON1_{Q192} and PON1_{R192} homozygotes under physiological conditions. As we had observed earlier when we determined K_m values for CPO hydrolysis with the PON1₁₉₂ alloforms purified from human plasma, PON1_{Q192} had a lower affinity for CPO (K_m = 168 μ M; Fig. 3C) than did PON1_{R192} (K_m = 109 μ M; Fig. 3D).

Many laboratories determine PON1₁₉₂ variability only by DNA analysis (PCR/restriction enzyme assays) and some additionally determine PON1 levels by measuring arylesterase activity [rates of phenyl acetate hydrolysis (AREase)] at ambient temperature. To generate factors for converting rates of hydrolysis of different substrates, rates of phenyl acetate hydrolysis at low salt (traditional AREase assay) and high salt (new PON1 status AREase assay) concentrations were plotted against rates of OP hydrolysis carried out under physiological conditions and saturating substrate for each PON1₁₉₂ phenotype. Figure 4 shows these rate plots for CPO and Figure 5 for DZO. Interestingly, the plot in Figure 4E, which shows the combined rates of phenyl acetate hydrolysis at high salt concentration plotted against rates of CPO hydrolysis, resolved all three PON1₁₉₂ phenotypes into separate groups, due to the differential effects of NaCl on the two PON1₁₉₂ alloforms.

Figure 5 shows the rates of phenyl acetate hydrolysis at low and high salt concentrations plotted against the rates of DZO hydrolysis for all three PON1₁₉₂ phenotypes. A plot of the combined rates for all three PON1₁₉₂ phenotypes with AREase assayed at high salt did not show the resolution of phenotypes observed with CPO (data not shown).

The plots shown in Figures 4 and 5 allowed us to generate factors for converting AREase rates measured at either high or low salt concentrations to physiologically relevant rates of OP hydrolysis. Conversion factors for AREase at both high and low salt concentrations were determined so that data from the non-OP PON1 status assay or data from many earlier

studies could be converted into physiologically relevant rates of DZO or CPO hydrolysis. These conversion factors are presented in Table 1.

To examine the physiological relevance of our earlier PON1 status analysis, where rates of DZO hydrolysis determined at pH 8.5 and 2 M NaCl were plotted against rates of paraoxon hydrolysis under the same conditions (Richter and Furlong, 1999), we plotted rates of DZO hydrolysis measured at high pH and salt concentration vs. rates of DZO hydrolysis determined under physiological conditions for each of the PON1₁₉₂ phenotypes (Fig. 6). Interestingly, for the PON1_{Q/Q192} and PON1_{Q/R192} individuals, the rates determined under the two conditions were nearly identical and for the PON1_{R/R192} individuals the conversion factor from physiological rates to rates determined at high salt, high pH rates was only 0.84 (Table 1). Since the rates of hydrolysis of DZO under physiological conditions were nearly the same as those determined under conditions of high pH and high salt, we compared both sets of data points on the same PON1 status plot (Fig. 7). These data show that whether the assay is carried out at 2 M NaCl and the pH optimum (8.5) for PON1 hydrolysis of DZO, or under physiological conditions at saturating substrate, the results are nearly identical.

Conversion factors for rates of AREase activity measured at low and high salt concentrations were determined, since there are many earlier studies on rates of phenyl acetate hydrolysis at ambient temperature and low salt concentration (Fig. 8). These conversion factors are summarized in Table 1. Since many earlier studies also have reported rates of PO hydrolysis, rates of AREase activity at high salt, were plotted against rates of POase activity for each PON1₁₉₂ phenotype to obtain conversion factors for these data. Interestingly, the high salt conditions for the AREase assay resulted in an excellent resolution of the three PON1₁₉₂ phenotypes (Fig. 8H), compared with the initial two-dimensional plot reported by Eckerson *et al.* with the low salt AREase/PO assay (1983). If assays are carried out at ambient temperatures that vary significantly from 23 °C, it may be necessary to generate temperature correction factors.

Discussion

Over the years, many different assays have been used to measure plasma PON1 activity, (reviewed in Ortigoza-Ferado *et al.*, 1984; Diepgen and Geldmacher v. Mallinckrodt, 1986). The initial interest in plasma PON1 activity was to provide an estimate of sensitivity to PO exposure prior to the demonstration that plasma PON1 level and phenotype are not significant in modulating PO exposures (Li *et al.*, 2000). In more recent years, as more catalytic activities of PON1 have been revealed, genetic variability in PON1 as a risk factor for cardiovascular disease (Jarvik *et al.*, 2000; reviewed in James, 2006) and other diseases (reviewed in Mackness *et al.*, 2002; Marsillach *et al.*, 2008) has generated increased interest. Many studies have examined only *PON1* single nucleotide polymorphisms (SNPs) (Mackness *et al.*, 2001; Wheeler *et al.*, 2004; Lawlor *et al.*, 2006). However, since PON1 levels vary at least 15-fold in the adult population (Davies *et al.*, 2006; Roest *et al.*, 2007; Furlong *et al.*, 2007) and the catalytic efficiencies can vary significantly between the Q192R alloforms, depending on the substrate (Davies *et al.*, 1996; Li *et al.*, 2000), studies that have examined only PON1 SNPs have not been informative (Jarvik *et al.*, 2003b). Haplotype analysis is also not informative, due to the lack of significant linkage disequilibrium across the PON1 locus (Jarvik *et al.*, 2003b). Several of the most experienced investigators in PON1 research have pointed out the inadequacy of examining PON1 genotype alone as a risk factor for disease or exposure (Mackness *et al.*, 2001; Deakin and James, 2004; La Du, 2003)

The PON1 status determinations, on the other hand, provide information on both PON1 levels and the functional PON1₁₉₂ alloform(s) present in plasma. We have identified

mutations in the PON1 gene that resulted in the inactivation of one or the other PON1₁₉₂ allele, based on discrepancies between PCR analysis and the functional PON1 status analysis (Jarvik et al., 2003a). There are many things that can happen between DNA and the final protein product. The PON1 status analysis examines the end product of all of these contributors by measuring the functional PON1 alloform(s) present in plasma (Richter and Furlong, 1999; Jarvik et al. 2000, 2003a).

It is worth mentioning problems associated with the use of POase measurements as determinants of risk of exposure or disease. The primary problem is the nearly 9-fold difference in catalytic efficiencies of the two PON1₁₉₂ alloforms in hydrolyzing PO (Li *et al.*, 2000). Many studies have tried to find correlations between POase activity and risk of disease or exposure lumping subjects of all three PON1₁₉₂ phenotypes together. Useful information could be obtained comparing controls and subjects of the same PON1₁₉₂ phenotype, but the large difference in catalytic efficiency of PO hydrolysis between the two PON1₁₉₂ phenotypes coupled with different gene frequencies in different ethnic groups, make the data impossible to interpret. To date, the only substrate that is useful for comparing hydrolytic activities across PON1₁₉₂ phenotypes is phenyl acetate (AREase) when assayed at low salt concentration (Furlong, 2006). AREase activity will provide a useful surrogate measure of PON1 protein levels across PON1₁₉₂ phenotypes.

In the earlier study by O'Leary *et al.* (2005) where they looked for a correlation between DZOase activities measured at low DZO level (50 μ M) vs. DZOase measured at high DZO level (500 μ M), they did not find a good correlation between the rates measured under the two conditions. Two factors contributed to the lack of correlation. The most important issue is trying to correlate activities of the two PON1₁₉₂ alloforms with different kinetic properties on the same plot. A second issue is most likely related to the HPLC assay used from which it would be difficult to obtain accurate rate data at low substrate concentrations. The rate data presented here for each PON1₁₉₂ phenotype show excellent correlations for all of the different substrates examined and conditions used for the assays. These experiments also show that it is important to compare rates within each PON1₁₉₂ phenotype since each PON1₁₉₂ alloform has different catalytic properties for many of the substrates hydrolyzed by this protein.

One of the main aims of this study was to provide a means of standardizing PON1 activity measurements and to provide conversion factors that will allow comparison of rates determined by one research group with those obtained by other laboratories. A recent comparison of inter-laboratory variability between our laboratory and Dr. Nina Holland's laboratory at the University of California at Berkeley showed that inter-laboratory variability in PON1 activity measurements was less than 10% (Huen *et al.*, submitted to *Clin. Chim. Acta*).

A further aim of this study was to provide physiologically relevant data with non-OP substrates that can be used in the development of PBPK/PD models (Timchalk and Poet, 2008). Appropriate conversion factors were obtained by measuring PON1 hydrolytic activities at saturating substrate with the non-OP substrates and the corresponding rates of relevant OP substrates at saturating substrate concentrations, physiological pH and temperature (Table 1). The data in Figures 4 and 5 show an excellent correlation between rates of hydrolysis for each substrate, when subjects with identical PON1₁₉₂ functional genotypes (phenotypes) are compared. Knowing the rate of substrate hydrolysis at saturating substrate concentration for a given subject allows for the determination of rate of OP hydrolysis at any given substrate concentration, based on the Michaelis-Menton relationship and using the conversion factors summarized in Table 1.

$$v = \frac{V_{\max} [S]}{K_m + [S]}$$

There are several additional points to consider in evaluating PON1 status and OP exposure. It is clear from earlier work that an individual's PON1 status plays a more important role in modulating direct exposure to the oxon component of the exposure (Shih *et al.*, 1998; Li *et al.*, 2000). Huff *et al.* (1995) reported that CPO inhibits acetylcholinesterase approximately 1,000 times faster than does the parent compound CPS. At this level of difference, it may well be that minor oxon contaminants in the parent compound preparations may account for the observed inhibition reported for the parent compound. It is also important to note that most if not all exposures contain an oxon component (Yuknavage *et al.*, 1997; California EPA, 1998). Another important consideration is the developmental dependence of plasma PON1 levels. Newborns have only one-third to one-fourth the plasma levels of PON1 that they will have as adults (Augustinsson and Barr, 1963; Ecobichon and Stephens, 1973; Mueller *et al.*, 1983; Furlong *et al.*, 2006) and the time of reaching mature levels varies from individual to individual (Cole *et al.*, 2003). Taking all of these factors into consideration, it is estimated that the combination of genetic and developmental variability of PON1 status in the population can result in as much as 130- to 160-fold variability in sensitivity to CPO exposure and 65-fold variability in sensitivity to DZO exposure (Furlong *et al.*, 2006).

In summary, it is not possible at this time to predict plasma levels of PON1 simply from DNA analysis, even if all nearly 200 PON1 SNPs (Furlong *et al.*, 2008) are analyzed. Since it is the functional level of enzyme and catalytic efficiency that determines rates of detoxication of both endogenous metabolites and xenobiotics, it is important to know both the plasma level of PON1 as well as the active PON1 alloform(s) present in plasma when estimating an individual's sensitivity to a specific OP exposure. For some exposures, e.g., DZS/DZO, it is primarily the level of plasma PON1 that is important for modulating the exposure. For other exposures such as CPS/CPO, it is both the plasma level of PON1 as well as PON1₁₉₂ alloform(s) present in plasma that are important in modulating the exposure (Li *et al.*, 2000). In no case is PON1 protein level unimportant. The high throughput PON1 status determination reveals the relative plasma PON1 level within each PON1₁₉₂ phenotype and at the same time reveals the alloform(s) present in plasma. For heterozygotes, there may be some bias as to the relative levels of the two PON1₁₉₂ alloforms since expression from each of the alleles is subject to *cis*-regulation (Brophy *et al.*, 2001b; Leviev and James, 2000; Suehiro *et al.*, 2000). This can be seen from the larger scatter of data points in the two-dimensional analysis where rates of one substrate are plotted against rates of a second substrate (Fig. 1). La Du *et al.* (2001) pointed out the importance of considering the concentration of each PON1 alloform in heterozygotes. The high-throughput two-substrate PON1 status assay/analysis provides the information necessary for establishing an individual's PON1 status, the relative plasma level of PON1 and the PON1₁₉₂ alloform(s) present in their plasma. A determination of actual plasma PON1 level is obtained either by measuring AREase activity at low salt (Furlong, 2006) or using the conversion factors from Table 1 and the data from the non-OP PON1 status determination.

The same approach that we describe here can be used to establish conversion factors for rates of hydrolysis of drugs/drug precursors or physiological metabolites. Such factors will be particularly useful for determining rates of hydrolysis/metabolism of substrates for which there are no convenient spectral assays. Once the calibration factors are established for each PON1₁₉₂ phenotype, the high-throughput, convenient spectral assays can be used to generate the important pharmacokinetic parameters necessary for estimating metabolism of a specific drug or metabolite.

Acknowledgments

This work was supported by National Institute of Environmental Health Sciences (ES09883) (Dr. Furlong), (ES04696) (Dr. H. Checkoway), (ES07033) (Dr. D. Eaton) and the National Heart, Lung, and Blood Institute (RO1 HL67406 and HL074366) (Dr. G. Jarvik).

References

- Adkins S, Gan KN, Mody M, La Du BN. Molecular basis for the polymorphic forms of human serum paraoxonase/arylesterase: glutamine or arginine at position 191, for the respective A or B allozymes. *Am. J. Hum. Genet* 1993;52:598–608. [PubMed: 7916578]
- Augustinsson K, Barr BM. Age variation in plasma arylesterase activity in children. *Clin. Chim. Acta* 1963;8:568–573. [PubMed: 14063971]
- Biggadike K, Angell RM, Burgess CM, Farrell RM, Hancock AP, Harker AJ, Irving WR, Ioannou C, Procopiou PA, Shaw RE, Solanke YE, Singh OM, Snowden MA, Stubbs RJ, Walton S, Weston HE. Selective plasma hydrolysis of glucocorticoid gamma-lactones and cyclic carbonates by the enzyme paraoxonase: an ideal plasma inactivation mechanism. *J. Med. Chem* 2000;43:19–21. [PubMed: 10633034]
- Brophy VH, Hastings MD, Clendenning JB, Richter RJ, Jarvik GP, Furlong CE. Polymorphisms in the human paraoxonase (PON1) promoter. *Pharmacogenetics* 2001a;11:77–84. [PubMed: 11207034]
- Brophy VH, Jampsa RL, Clendenning JB, Jarvik GP, Furlong CE. Promoter polymorphisms affect paraoxonase (PON1) expression. *Am. J. Hum. Genet* 2001b;68:1428–1436. [PubMed: 11335891]
- California Environmental Protection Agency. Report for the Application and Ambient Air Monitoring of Chlorpyrifos (and the oxon analogue). Tulare County During Spring/Summer, 1996. 1998. <http://www.cdpr.ca.gov/docs/emon/pubs/tac/tacpdfs/chlrpfs.pdf>
- Chambers JE, Ma T, Boone JS, Chambers HW. Role of detoxication pathways in acute toxicity levels of phosphorothionate insecticides in the rat. *Life Sci* 1994;54:1357–1364. [PubMed: 7514706]
- Cole TB, Jampsa RL, Walter BJ, Arndt TL, Richter RJ, Shih DM, Tward A, Lusi AJ, Jack RM, Costa LG, Furlong CE. Expression of human paraoxonase (PON1) during development. *Pharmacogenetics* 2003;13:357–364. [PubMed: 12777966]
- Cole TB, Walter BJ, Shih DM, Tward AD, Lusi AJ, Timchalk C, Richter RJ, Costa LG, Furlong CE. Toxicity of chlorpyrifos and chlorpyrifos oxon in a transgenic mouse model of the human paraoxonase (PON1) Q192R polymorphism. *Pharmacogenet. and Genomics* 2005;15:589–598.
- Davies HG, Richter RJ, Keifer M, Broomfield CA, Sowalla J, Furlong CE. The effect of the human serum paraoxonase polymorphism is reversed with diazoxon, soman and sarin. *Nat. Genet* 1996;14:334–336. [PubMed: 8896566]
- Deakin S, Leviev I, Brulhart-Meynet MC, James RW. Paraoxonase-1 promoter haplotypes and serum paraoxonase: a predominant role for polymorphic position - 107, implicating the Sp1 transcription factor. *Biochem. J* 2003;372:643–649. [PubMed: 12639220]
- Deakin SP, James RW. Genetic and environmental factors modulating serum concentrations and activities of the antioxidant enzyme paraoxonase-1. *Clin. Sci. (Lond.)* 2004;107:435–447. [PubMed: 15265000]
- Diepgen TL, Geldmacher v. Mallinckrodt M. The human serum paraoxonase polymorphism. *Arch. Toxicol. Suppl* 1986;9:154–158. [PubMed: 3028323]
- Dowd JE, Riggs DS. A comparison of estimates of Michaelis-Menten kinetic constants from various linear transformations. *J. Biol. Chem* 1965;240:863–869. [PubMed: 14275146]
- Eckerson HW, Wyte CM, La Du BN. The human serum paraoxonase/arylesterase polymorphism. *Am. J. Hum. Genet* 1983;35:1126–1138. [PubMed: 6316781]
- Ecobichon DJ, Stephens DS. Perinatal development of human blood esterases. *Clin Pharmacol. Ther* 1973;14:41–47. [PubMed: 4734200]
- Furlong CE, Richter RJ, Chapline C, Crabb JW. Purification of rabbit and human serum paraoxonase. *Biochemistry* 1991;30:10133–10140. [PubMed: 1718413]

- Furlong CE, Holland N, Richter RJ, Bradman A, Ho A, Eskenazi B. PON1 status of farmworker mothers and children as a predictor of organophosphate sensitivity. *Pharmacogenet. Genomics* 2006;16:183–190. [PubMed: 16495777]
- Furlong, CE.; Richter, RJ.; Li, W-F.; Brophy, VH.; Carlson, C.; Meider, M.; Nickerson, D.; Costa, LG.; Ranchalis, J.; Lusi, AJ.; Shih, DM.; Tward, A.; Jarvik, GP. The functional consequences of polymorphisms in the human PON1 gene. In: Mackness, B.; Mackness, M.; Aviram, M.; Paragh, G., editors. *The Paraoxonases: Their Role in Disease, Development and Xenobiotic Metabolism*. Springer; Dordrecht, The Netherlands: 2008. p. 267-281.
- Gan K, Smolen A, Eckerson HW, La Du BN. Purification of human serum paraoxonase/arylesterase: Evidence for one esterase catalyzing both activities. *Drug Metab. Dispos* 1991;19:100–106. [PubMed: 1673382]
- Hasset C, Richter RJ, Humbert R, Chapline C, Crabb JW, Omiecinski CJ, Furlong CE. Characterization of cDNA clones encoding rabbit and human serum paraoxonase: the mature protein retains its signal sequence. *Biochemistry* 1991;30:10141–10149. [PubMed: 1657140]
- Huff R, Abou-Donia MB. In vitro effect of chlorpyrifos oxon on muscarinic receptors and adenylate cyclase. *Neurotoxicology* 1995;16:281–290. [PubMed: 7566687]
- Humbert R, Adler DA, Distech CM, Hassett C, Omiecinski CJ, Furlong CE. The molecular basis of the human serum paraoxonase activity polymorphism. *Nat. Genet* 1993;3:73–76. [PubMed: 8098250]
- James RW. A long and winding road: defining the biological role and clinical importance of paraoxonases. *Clin. Chem. Lab. Med* 2006;44:1052–1059. [PubMed: 16958594]
- Jarvik GP, Rozek LS, Brophy VH, Hatsukami TS, Richter RJ, Schellenberg GD, Furlong CE. Paraoxonase phenotype is a better predictor of vascular disease than PON1192 or PON155 genotype. *Atheroscler. Thromb. Vasc. Biol* 2000;20:2442–2447.
- Jarvik GP, Jampsa R, Richter RJ, Carlson C, Rieder M, Nickerson D, Furlong CE. Novel paraoxonase (PON1) nonsense and missense mutations predicted by functional genomic assay of PON1 status. *Pharmacogenetics* 2003a;13:291–295. [PubMed: 12724622]
- Jarvik GP, Hatsukami TS, Carlson CS, Richter RJ, Jampsa R, Brophy VH, Margolin S, Rieder MJ, Nickerson DA, Schellenberg GD, Heagerty PJ, Furlong CE. Paraoxonase activity, but not haplotype utilizing the linkage disequilibrium structure predicts vascular disease. *Arterioscler. Thromb. Vasc. Biol* 2003b;23:1465–1471. [PubMed: 12805074]
- La Du BN, Billecke S, Hsu C, Haley RW, Broomfield CA. Serum paraoxonase (PON1) isozymes: the quantitative analysis of isozymes affecting individual sensitivity to environmental chemicals. *Drug Metab. Dispos* 2001;29:566–569. [PubMed: 11259353]
- La Du BN. Future studies of low-activity PON1 phenotype subjects may reveal how PON1 protects against cardiovascular disease. *Arterioscler. Thromb. Vasc. Biol* 2003;23:1317–1318. [PubMed: 12909569]
- Lawlor DA, Gaunt TR, Hinks LJ, Davey Smith G, Timpson N, Day IN, Ebrahim S. The association of the PON1 Q192R polymorphism with complications and outcomes of pregnancy: findings from the British Women's Heart and Health cohort study. *Paediatr. Perinat. Epidemiol* 2006;20:244–250. [PubMed: 16629699]
- Levieu I, James RW. Promoter polymorphisms of human paraoxonase PON1 gene and serum paraoxonase activities and concentrations. *Arterioscler. Thromb. Vasc. Biol* 2000;20:516–521. [PubMed: 10669651]
- Li W-F, Costa LG, Furlong CE. Serum paraoxonase status: a major factor in determining resistance to organophosphates. *J. Toxicol. Environ. Health* 1993;40:337–346. [PubMed: 7693961]
- Li W-F, Furlong CE, Costa LG. Paraoxonase protects against chlorpyrifos toxicity in mice. *Toxicol. Lett* 1995;76:29–226.
- Li W-F, Costa LG, Richter RJ, Hagen T, Shih DM, Tward A, Lusi AJ, Furlong CE. Catalytic efficiency determines the in vivo efficacy of PON1 for detoxifying organophosphates. *Pharmacogenetics* 2000;10:767–780. [PubMed: 11191881]
- Mackness B, Davies GK, Turkie W, Lee E, Roberts DH, Hill E, Roberts C, Durrington PN, Mackness MI. Paraoxonase status in coronary heart disease: are activity and concentration more important than genotype? *Arterioscler. Thromb. Vasc. Biol* 2001;21:1451–1457. [PubMed: 11557671]

- Mackness, B.; Durrington, PN.; Mackness, MI. PON1 in other diseases. In: Costa, LG.; Furlong, CE., editors. Paraoxonase (PON1) in Health and Disease: Basic and Clinical Aspects. Kluwer Academic Press; Boston: 2002. p. 185-195.
- Marsillach, J.; Parra, S.; Ferre, N.; Coll, B.; Alonso-Villaverde, C.; Joven, J.; Camps, J. Paraoxonase-1 in chronic liver diseases, neurological diseases and HIV infection. In: Mackness, B.; Mackness, M.; Aviram, M.; Paragh, G., editors. The Paraoxonases: Their Role in Disease, Development and Xenobiotic Metabolism. Springer; Dordrecht, The Netherlands: 2008. p. 187-198.
- Mueller RF, Hornung S, Furlong CE, Anderson J, Giblett ER, Motulsky AG. Plasma paraoxonase polymorphism: a new enzyme assay, population, family, biochemical and linkage studies. *Am. J. Hum. Genet* 1983;35:393–408. [PubMed: 6305189]
- O'Leary KA, Edwards RJ, Town MM, Boobis AR. Genetic and other sources of variation in the activity of serum paraoxonase/diazoxonase in humans: consequences for risk from exposure to diazinon. *Pharmacogenet. Genomics* 2005;15:51–60. [PubMed: 15864126]
- Ortigoza-Ferado J, Richter RJ, Hornung SK, Motulsky AG, Furlong CE. Paraoxon hydrolysis in human serum mediated by a genetically variable arylesterase and albumin. *Am. J. Hum. Genet* 1984;36:295–305. [PubMed: 6324579]
- Ozer EA, Pezzulo A, Shih DM, Chun C, Furlong C, Lusi AJ, Greenberg EP, Zabner J. Human and murine paraoxonase 1 are host modulators of *P. aeruginosa* quorum-sensing. *FEMS Microbiol. Lett* 2005;253:29–37. [PubMed: 16260097]
- Pond AL, Chambers HW, Chambers JE. Organophosphate detoxication potential of various rat tissues via A-esterase and alsesterase activities. *Toxicol. Lett* 1995;78:245–252. [PubMed: 7542808]
- Richter RJ, Furlong CE. Determination of paraoxonase (PON1) status requires more than genotyping. *Pharmacogenetics* 1999;9:745–753. [PubMed: 10634137]
- Rochu D, Chabriere E, Masson P. Human paraoxonase: a promising approach for pre-treatment and therapy of organophosphorus poisoning. *Toxicology* 2007;233:47–59. Epub 2006 Sep 3. [PubMed: 17007987]
- Roest M, van Himbergen TM, Barendrech AB, Peeters PH, van der Schouw YT, Voorbij HA. Genetic and environmental determinants of the PON-1 phenotype. *Eur. J. Clin. Invest* 2007;37:187–196. [PubMed: 17359486]
- Shih DM, Gu L, Xia Y-R, Navab M, Li W-F, Hama S, Castellani LW, Furlong CE, Costa LG, Fogelman AM, Lusi AJ. Mice lacking serum paraoxonase are susceptible to organophosphate toxicity and atherosclerosis. *Nature* 1998;394:284–287. [PubMed: 9685159]
- Stoltz DA, Ozer EA, Taft PJ, Barry M, Liu L, Kiss PJ, Moninger TO, Parsek MR, Zabner J. *Drosophila* are protected from *Pseudomonas aeruginosa* lethality by transgenic expression of paraoxonase-1. *J. Clin. Invest.* Aug 14;2008 [Epub ahead of print] PMID: 18704198.
- Suehiro T, Nakamura T, Inoue M, Shiinoki T, Ikeda Y, Kumon Y, Shindo M, Tanaka H, Hashimoto K. A polymorphism upstream from the human paraoxonase (PON1) gene and its association with PON1 expression. *Atherosclerosis* 2000;150:295–298. [PubMed: 10856521]
- Timchalk C, Poet TS. Development of a physiologically based pharmacokinetic and pharmacodynamic model to determine dosimetry and cholinesterase inhibition for a binary mixture of chlorpyrifos and diazinon in the rat. *Neurotoxicology* 2008;29:428–43. Epub 2008 Mar 10. [PubMed: 18394709]
- Tougou K, Nakamura A, Watanabe S, Okuyama Y, Morino A. Paraoxonase has a major role in the hydrolysis of prulifloxacin (NM441), a prodrug of a new antibacterial agent. *Drug Metab. Dispos* 1998;26:355–359. [PubMed: 9531524]
- Wheeler JG, Keavney BD, Watkins H, Collins R, Danesh J. Four paraoxonase gene polymorphisms in 11212 cases of coronary heart disease and 12786 controls: meta-analysis of 43 studies. *Lancet* 2004;363:689–695. [PubMed: 15001326]
- Yamada Y, Takatori T, Nagao M, Iwase H, Kurada N, Yanagida J, Shinozuka T. Expression of paraoxonase isoform did not confer protection from acute sarin poisoning in the Tokyo subway terrorist attack. *Int. J. Leg. Med* 2001;115:82–84.
- Yuknavage KL, Fenske RA, Kalman DA, Keifer MC, Furlong CE. Simulated dermal contamination with capillary samples and field cholinesterase biomonitoring. *J. Toxicol. and Env. Health* 1997;51:35–55. [PubMed: 9169060]

Figure 1.
Structures of the compounds used in this study.

Figure 2.

Comparison of the two protocols for determining PON1 status. **(A)** Assays using the highly toxic OP substrates DZO and PO; and **(B)** assays using the non-OP substrates phenyl acetate and CMA. The 183 plasma samples included 86 PON1^{Q192} homozygotes, 79 heterozygotes and 18 PON1^{R192} homozygotes with genotypes verified by PCR. From Richter *et al.* Submitted to Circulation: Cardiovascular Genetics.

Figure 3.

Substrate dependence of DZO hydrolysis by **A**, a PON1_{Q192} plasma sample ($K_m = 0.52$ mM); **B**, a PON1_{R192} plasma sample ($K_m = 0.27$ mM) and for CPO hydrolysis by **C**, a PON1_{Q192} plasma sample ($K_m = 168$ μ M) and **D**, a PON1_{R192} plasma sample ($K_m = 109$ μ M). The plasma samples were matched for rates of AREase activity. The substrate concentrations are indicated in the graphs and only the initial linear rates of hydrolysis were used for kinetic analyses. The closed symbols indicate rates of hydrolysis (v) at the indicated substrate concentration and the open symbols are the $[S]/v$ reciprocal values for each of the rate data points (v). K_m values were determined from the x-axis intercepts of the $[S]/v$ vs. $[S]$ plots (Dowd and Riggs, 1965).

Figure 4.

Comparison of the rates of AREase hydrolysis measured at conditions of low salt vs. CPOase, at physiological conditions (ph 7.4, 150 mM NaCl, 37 °C) for samples of each PON1₁₉₂ phenotype (A-C), and AREase hydrolysis measured at conditions of high salt (2M NaCL)/high pH (pH 8.0) vs. CPOase, at physiological conditions (pH 7.4, 150 mM NaCl, 37 °C) for samples of each PON1₁₉₂ phenotype (E-G). (○-○) = PON1_{Q/Q192} plasma samples, n = 61; (■-■) = PON1_{Q/R192} plasma samples, n = 49; (Δ-Δ) = PON1_{R/R192} plasma samples, n = 18. Panel D = combined data from A-C; Panel H = Combined data from panels E-G.

Figure 5.

Comparison of the rates of AREase hydrolysis measured at conditions of low salt vs. DZOase, at physiological conditions (pH 7.4, 150 mM NaCl, 37 °C) for samples of each PON1₁₉₂ phenotype (A-C), and AREase hydrolysis measured at conditions of high salt (2 M NaCl)/high pH (pH 8.0) vs. DZOase, at physiological conditions (pH 7.4, 150 mM NaCl, 37 °C) for samples of each PON1₁₉₂ phenotype (E-G). (○-○) = PON1^Q/Q¹⁹² plasma samples, n = 61; (■-■) = PON1^Q/R¹⁹² plasma samples, n = 49, (Δ-Δ) = PON1^R/R¹⁹² plasma samples, n = 18

Figure 6.

Comparison of rates of DZO hydrolysis measured at high salt (2M NaCl)/high pH (8.5) vs. rates of DZO hydrolysis measured under physiological conditions (ph 7.4, 150 mM NaCl, 37 °C). (○—○) = PON1_{Q/Q192} plasma samples, n = 61; (■—■) = PON1_{Q/R192} plasma samples, n = 49; (Δ—Δ) = PON1_{R/R192} plasma samples, n = 18.

Figure 7.

PON1 status analysis determined with the high pH (8.5), high salt (2 M NaCl) DZOase assay ($n = 128$) and the “physiological” DZOase assay. Closed symbols = high salt/high pH DZOase data points and open symbols = the “physiological” DZOase data points. POase activity was determined at 2 M NaCl and pH 8.5, since paraoxon is not a physiologically relevant substrate (Li *et al.*, 2000).

Figure 8.

A-C, Plots of AREase activity at high vs. AREase activity at low salt for the three PON1₁₉₂ phenotypes. **D**, combined data from **A-C**. **E-G**, plots of AREase measured at high salt concentration vs. POase measured at high salt concentration. **H**, combined data from **E-G**. (○-○) = PON1_{Q/Q192} plasma samples, n = 61; (■-■) = PON1_{Q/R192} plasma samples, n = 49; (Δ-Δ) = PON1_{R/R192} plasma samples, n = 18.

Table 1

Conversion factors for rates of substrate hydrolysis

Phenotype	Conversion Factors	r^2
QQ	${}^b\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 172 = \text{DZOase}_{\text{phys}}{}^c \text{ (U/L)}$	0.93
QR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 204 = \text{DZOase}_{\text{phys}} \text{ (U/L)}$	0.82
RR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 286 = \text{DZOase}_{\text{phys}} \text{ (U/L)}$	0.87
QQ	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 69 = \text{CPOase}_{\text{phys}}{}^d \text{ (U/L)}$	0.87
QR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 103 = \text{CPOase}_{\text{phys}} \text{ (U/L)}$	0.88
RR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 189 = \text{CPOase}_{\text{phys}} \text{ (U/L)}$	0.89
QQ	${}^e\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 110 = \text{DZOase}_{\text{phys}} \text{ (U/L)}$	0.84
QR	$\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 100 = \text{DZOase}_{\text{phys}} \text{ (U/L)}$	0.72
RR	$\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 83 = \text{DZOase}_{\text{phys}} \text{ (U/L)}$	0.93
QQ	$\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 45 = \text{CPOase}_{\text{phys}} \text{ (U/L)}$	0.73
QR	$\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 50 = \text{CPOase}_{\text{phys}} \text{ (U/L)}$	0.84
RR	$\text{AREase}_{\text{LS}} \text{ (U/ml)} \times 55 = \text{CPOase}_{\text{phys}} \text{ (U/L)}$	0.92
QQ	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 3.8 = \text{POase} \text{ (U/L)}$	0.75
QR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 15.9 = \text{POase} \text{ (U/L)}$	0.50
RR	$\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 47.6 = \text{POase} \text{ (U/L)}$	0.90
QQ	${}^f\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 1.6 = \text{AREase}_{\text{LS}} \text{ (U/ml)}$	0.85
QR	${}^f\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 2.0 = \text{AREase}_{\text{LS}} \text{ (U/ml)}$	0.66
RR	${}^f\text{AREase}_{\text{HS}} \text{ (U/ml)} \times 3.5 = \text{AREase}_{\text{LS}} \text{ (U/ml)}$	0.83
QQ	$\text{DZOase}_{\text{phys}} \text{ (U/L)} \times 1.08 = \text{DZOase}_{\text{HS}}{}^g \text{ (U/L)}$	0.90
QR	$\text{DZOase}_{\text{phys}} \text{ (U/L)} \times 1.01 = \text{DZOase}_{\text{HS}} \text{ (U/L)}$	0.91
RR	$\text{DZOase}_{\text{phys}} \text{ (U/L)} \times 0.84 = \text{DZOase}_{\text{HS}} \text{ (U/L)}$	0.87

^a Correlation coefficient squared^b AREase_{HS} = arylesterase activity measured in buffer and 2M NaCl^c DZOase_{phys} = Diazoxonase activity measured under physiological conditions^d CPOase_{phys} = Chlorpyrifos oxonase activity measured under physiological conditions^e AREase_{LS} = arylesterase activity measured in buffer^f From Richter et al. (submitted to *Circulation: Cardiovascular Genetics*)^g DZOase_{HS} = Diazoxonase activity measured at 2M NaCl, pH 8.5