Published in final edited form as:

Org Lett. 2011 February 18; 13(4): 703–705. doi:10.1021/ol102936z.

Access to Functionalized Steroid Side Chains via Modified Julia Olefination

Enver Cagri Izgu, Aaron C. Burns, and Thomas R. Hoye*

Department of Chemistry, 207 Pleasant Street, SE, University of Minnesota, Minneapolis, Minnesota 55455

Abstract

Various functionalized steroidal side chains were conveniently accessed by a modified Julia olefination strategy using a common sulfone donor and an appropriate α -branched aldehyde acceptor. For the coupling of these hindered classes of reaction partners (and in contrast to typically observed trends), the benzothiazolyl(BT)-sulfone anion gave superior outcomes compared to the phenyltetrazolyl(PT)-sulfone anion.

Steroids with oxidatively modified side chains comprise an important family of compounds, largely because of their biological properties (e.g., **1-4**, Figure 1). Our interest in the preparation of analogs of sea lamprey pheromone components (e.g., **31d**) drove a need for a flexible strategy in which a common intermediate could be used in a conjunctive fashion to attach an array of structurally diverse side chains (cf. Table 1). In this regard, we have studied the use of the modified Julia olefination² involving C22-sulfonyl steroids as the anionic donor partner, and our results are described here.

Previously, steroidal aldehydes **5** have been used as the electrophilic acceptor component in Horner-Wadsworth-Emmons (HWE, **6a**),^{3a-e} modified Julia (**6b**),^{3f} and Wittig (with stabilized ylides, **6c**)^{3g-m} olefination reactions to forge the C22-C23 alkene in products **9**. Alternatively, steroidal phosphonium ylide (**7a**) and phenylsulfonyl (**7b**) donors were coupled with aldehyde acceptors **8** via Wittig³ⁿ and classical Julia^{3o} olefination reactions, respectively (Scheme 1). In this study, we have established the modified Julia coupling with donors **7c**, primarily with the benzothiazolyl (BT) sulfone [although we have also compared the use of the 1-phenyl-1*H*-tetrazol-5-yl (PT) sulfone] and a variety of acceptor aldehydes **8**.

To test feasibility, both the PT and BT sulfones **13-PT** and **13-BT**, respectively, were prepared as outlined in Scheme 2 from *i*-stigmasteryl methyl ether (**10**, two steps, 74% yield from stigmasteryl).⁴ Ozonolysis (and reductive workup with NaBH₄) of the disubstituted olefin smoothly provided the primary alcohol **11** [and (*S*)-2-ethyl-3-methylbutan-1-ol]. It is worth noting that a major problem associated with the ozonolysis of *i*-steroids, namely undesired oxidation of the methine C–H bond at the C6-ether,⁵ is avoided by using the unconventional solvent tetrahydrofuran as component of the reaction medium (THF/MeOH;

*hoye@umn.edu .

Supporting Information Available: Experimental procedures and spectroscopic characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

10:1). The resulting alcohol **11** was formed in high yield (86%), and butyrolactone was isolated as a byproduct, its amount increasing with increased reaction time. We believe that THF effectively buffers the ozonolysis reaction by acting as a sacrificial reductant and preventing over-oxidation of **10** and its derived products. The thioethers **12-PT** and **12-BT** were then prepared from **11** using the Mitsunobu protocol (PTSH, DIAD, PPh₃). Each was subsequently oxidized to the sulfone **13-PT** or **13-BT** (ammonium paramolybdate, H₂O₂).

We first attempted olefination using the *potassium* anion of **13-PT** and **13-BT**. Preforming the anion (KHMDs, THF, -78 °C) and addition of propionaldehyde provided none of the expected olefination product, nor was the starting sulfone recovered. When a solution of KHDMS was added at -78 °C to a THF solution containing both the sulfone **13-BT** and propionaldehyde, the desired propylidene coupling product was isolated in low yield with the *cis* isomer predominating.

We then turned our attention to the *sodium* anions derived from **13-PT** and **13-BT** by studying the relative stability of the sulfones upon metallation with NaHMDS in THF at -78 °C (Scheme 3). Each anion was quenched with saturated NH₄Cl solution 15 minutes after addition of base. The starting sulfones were reisolated by SiO₂ chromatography (MPLC) with 61 vs 96% recovery efficiency, respectively. This shows that the **13-BT** anion was more stable than that of **13-PT**.

We next tested the relative efficiency of each of the donor sulfones **13** to effect olefination using butyraldehyde (**8a**) as a simple acceptor substrate and NaHMDS as the base (Scheme 4). The behavior mirrored that seen in the above stability studies; namely, the yield of alkene **14a** was higher when the benzothiazolyl sulfone **13-BT** was used.

Moreover (and as described elsewhere⁶), the use of HMPA as an additive improved the **14a-E:14a-Z** product ratio.

We then studied the olefination reactions of a series of aldehydes **8** with **13-BT** as the sulfone donor (Table 1). The aldehydes contain α -methyl or α -alkoxy branching. Aldehyde (**S**)-**8b** provided **14b** both in high yield and with excellent *E/Z*-selectivity (entry 1). Use of the enantiomeric aldehyde (**R**)-**8b** gave the C24-epimer in 80% yield and with an *E/Z*-ratio of 82:18 (entry 2). Entries 3-5 demonstrate additional scope of the method. Aldehyde (**S**)-**8e** showed a high *E/Z*-selectivity. Use of the racemate *rac*-**8e** (2 equiv, entry 6) produced similar amounts of C24-epimers. While the *E/Z*-ratio of product **14e-(24R)** was essentially the same as that observed in entry 5 (i.e., 90:10), the epimeric mixture of alkenes **14e-(24S)** was formed with reduced selectivity for alkene geometry (i.e., 65:35). Thus, the degree of matching/mis-matching for the substrates in either of entries 1 vs. 2 or 5 vs. 6 is small.

In conclusion, this study demonstrates the utility of a modified Julia olefination strategy for providing easy access to steroidal products containing a variety of functionalized side chains. The steric demand of both α -branched coupling partners that participate in this transformation is noteworthy. Contrary to the general trend observed for modified Julia reaction using less hindered pairs of substrates, the olefination efficiency and alkene diastereoselectivity of the steroidal BT-sulfonyl donor was found here to be superior to the PT-sulfonyl version.⁷ This is likely a result of greater lability of the metallated PT-sulfone anion (Scheme 3). An ancillary observation of note is the use of THF as the bulk solvent to improve the ozonolysis of cyclopropyl-containing substrate **10**. This protocol may be useful for chemoselective transformation of other complex substrates bearing functionality sensitive to oxidation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

These studies were supported by the National Institute of General Medical Sciences (GM- 65597) and the National Cancer Institute (CA-76497) of the United States National Institutes of Health.

References

- (1)(a). Moore KS, Wehrli S, Roder M, Forrest JN Jr. McCrimmon D, Zasloff M. *Proc. Natl. Acad. Sci. U.S.A* 1993;90:1354–1358. [PubMed: 8433993] (b) Wehrli SL, Moore KS, Roder H, Durell S, Zasloff M. *Steroids* 1993;58:370–378. [PubMed: 8212087] (c) Rao MN, Shinnar AE, Noecker LA, Chao TL, Feibush B, Snyder B, Sharkansky I, Sarkahian A, Zhang X, Jones SR, Kinney WA, Zasloff M. *J. Nat. Prod* 2000;63:631–635. [PubMed: 10843574] (d) Sorensen PW, Fine JM, Dvornikovs V, Jeffrey CS, Shao F, Wang J, Vrieze LA, Anderson KR, Hoye TR. *Nat. Chem. Biol* 2005;1:324–328. [PubMed: 16408070] (e) Mellado GG, Zubia E, Ortega MJ, López-González PJ. *Steroids* 2004;69:291–299. [PubMed: 15183695]
- (2)(a). Baudin JB, Hareau G, Julia SA, Ruel O. *Tetrahedron Lett* 1991;32:1175–1178. (b) Blakemore PR, Cole WJ, Kocienski PJ, Morley A. *Synlett* 1998:26–28.
- (3)(a). Giroux S, Corey EJ. *J. Am. Chem. Soc* 2007;129:9866–9867. [PubMed: 17658816] (b) Yamamoto S, Watanabe B, Otsuki J, Nakagawa Y, Akamatsu M, Miyagawa H. *Bio. & Med. Chem* 2006;14:1761–1770. (c) Tochtrop GP, DeKoster GT, Cistola DP, Covey DF. *Bio. & Med. Chem. Lett* 2002;12:433–435. (d) Kinney WA, Zhang X, Williams JJ, Johnston S, Michalak RS, Deshpande M, Dostal L, Rosazza JP. *N. Org. Lett* 2000;2:2921–2922. (e) Jones SR, Selinsky BS, Rao MN, Zhang X, Kinney WA, Tham FS. *J. Org. Chem* 1998;63:3786–3789. (f) Jiang B, Shi H, Xu M, Wang W, Zhou W. *Tetrahedron* 2008;64:9738–9744. (g) Shingate BB, Hazra BG, Pore VS, Gonnade RG, Bhadbhade MM. *Tetrahedron* 2007;65:5622–5635. (h) Shu Y, Jones SR, Kinney WA, Selinsky BS. *Steroids* 2002;67:291–304. [PubMed: 11856553] (i) Rao MN, McGuigan MA, Zhang X, Shaked Z, Kinney WA, Bulliard M, Laboue B, Lee NE. *J. Org. Chem* 1997;62:4541–4545. [PubMed: 11671794] (j) Okamoto M, Tabe M, Fuji T, Tanaka T. *Tetrahedron: Asymmetry* 1995;6:767–768. (k) Wei-Shan Z, HuiQiang Z, Zhi-Qin W. *J. Chem. Soc., Perkin Trans. 1* 1990:2281–2286. (l) Wei-Shan Z, Biao J, Xin-fu P. *Tetrahedron* 1990;46:3173–3188. (m) Zheng-Wu S, Wei-Shan Z. *J. Chem. Soc., Perkin Trans. 1* 1990:1765–1767. (n) Harney DW, Macrides TA. *J. Chem. Soc., Perkin Trans. 1* 1997:1353–1356. (o) D'Ambra TE, Javitt NB, Lacy J, Srinivasan P, Warchol T. *Steroids* 2000;65:401–407. [PubMed: 10899540]
- (4). Fernholz E, Ruigh WL. *J. Am. Chem. Soc* 1940;62:3346–3348.
- (5). Spencer TA, Li D, Russel JS, Tomkinson NCO, Willson TM. *J. Org. Chem* 2000;65:1919–1923. [PubMed: 10774010]
- (6). Liu P, Jacobsen EN. *J. Am. Chem. Soc* 2001;123:10772–10773. [PubMed: 11674024]
- (7)(a). Blakemore PR. *J. Chem. Soc., Perkin Trans. 1* 2002:2563–2585. (b) Aissa C. *Eur. J. Org. Chem* 2009:1831–1844.

Figure 1.
Some of the related highly potent C24-functionalized steroids (**11a-b**, **2^{1c}**, **3^{1d}**, and **41e**).

Scheme 1.

Overview of the Coupling Strategies to Construct Oxygenated Steroidal Side Chains.^a

^a St = generic steroid nucleus; BT = benzothiazolyl; PT = 1-phenyl-1*H*-tetrazol-5-yl.

Scheme 2.
Synthesis of the Key Sulfones **13-PT** and **13-BT**.

Scheme 3. Recovery/Stability of Sulfone Anions.^a^a Stoichiometric ratio of sulfone/NaHMDS was 1:1.2.

Scheme 4. Olefination of Sulfones **13-PT** or **13-BT** with **8a**.^a^a Stoichiometric ratio of sulfone/**8a**/NaHMDS was 1:1:1.2.

Table 1

Alkenes **14** via the Modified Julia Olefination.

entry	aldehyde	R	yield (%)	E/Z ^a
1	$(S)\text{-}8b$	$\mathbf{14b}$	82	94:6
2	$(R)\text{-}8b$	$\text{epi-}14b$	80	82:18
3	$8c$	$\mathbf{14c}$	90	75:25

entry	aldehyde	R	yield (%)	E/Z ^a
4	(S)-8d	14d	60	85:15
5	(S)-8e	14e-(24R)	81	90:10
6	rac-8e	14e-(24S) + 14e-(24R)	60	ca. 65:35 from 14e-(24S)

^aE/Z ratios determined by ¹H NMR analysis of product mixtures.