Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2011 March ; 20(3): 449–453. doi:10.1158/1055-9965.EPI-10-1170.

Racial differences in the incidence of mesenchymal tumors associated with *EWSR1* translocation

Jennifer Worch, MD, Jobin Cyrus, MPH, Robert Goldsby, MD, Katherine K. Matthay, MD, John Neuhaus, PhD, and Steven G. DuBois, MD

Department of Pediatrics and Department of Epidemiology/Biostatistics, University of California, San Francisco School of Medicine, San Francisco, California

Abstract

Background—The incidence of Ewing sarcoma varies by race, with very low rates among persons of African and East Asian ancestry. The incidence by race of other mesenchymal tumors that also harbor *EWSR1* translocations has not been studied.

Methods—The SEER database was queried to find cases of mesenchymal tumors associated with *EWSR1* translocations: Ewing sarcoma; clear cell sarcoma; extraskeletal myxoid chondrosarcoma; myxoid liposarcoma; desmoplastic small round cell tumor and myoepithelial tumor. Age-adjusted incidence rates were calculated for white, African-American, and Asian/Native American populations and compared statistically.

Results—Ewing sarcoma was significantly less common in the African-American and Asian/Native American populations compared to the white population, with incidence rate ratios of 0.12 (95% confidence interval 0.08 – 0.20; $p < 0.001$) and 0.54 (95% confidence interval 0.41 – 0.69; $p < 0.001$), respectively. Desmoplastic small round cell tumor was significantly more common in the African-American population compared to the white population (incidence rate ratio 3.0; 95% confidence interval 1.62 – 5.49; $p < 0.001$). Myxoid liposarcoma was significantly less common in the Asian/Native American population compared to the white population (incidence rate ratio 0.72; 95% confidence interval 0.56 – 0.92; p -value 0.006). The incidence rates for extraskeletal myxoid chondrosarcoma, myoepithelial tumors, and clear cell sarcoma did not differ significantly by race.

Conclusions—Tumors associated with *EWSR1* translocation are not uniformly more common in people of European ancestry.

Impact—The relationship between race and *EWSR1* somatic translocation is complex. Future studies investigating the genetic epidemiology of *EWSR1* translocated tumors are required.

Keywords

Mesenchymal tumors; sarcoma translocations; *EWSR1*; race; SEER

Corresponding Author: Steven G. DuBois, MD, Department of Pediatrics, UCSF School of Medicine, 505 Parnassus Avenue, M646, San Francisco, CA 94143-0106, Telephone: 415-476-3831, Facsimile: 415-502-4327, duboiss@peds.ucsf.edu.

Disclosures: None

Previous presentation: Presented in part at the 2010 Connective Tissue Oncology Society Annual Meeting; Paris, France.

Introduction

Ewing sarcoma (ES) is a malignant tumor of bone and soft tissue characterized by genetic translocations involving the Ewing sarcoma breakpoint region (*EWSR1*) in 95% of cases (1). While *EWSR1* was initially identified in ES, a number of other mesenchymal tumors also harbor *EWSR1* gene translocations (2,3). These tumors include clear cell sarcoma (CCS) with a *EWSR1/ATF1* frequency of 87 - 94% (4-6), extraskeletal myxoid chondrosarcoma (EMC) with a *EWSR1/CHN* frequency of 83% (7), myxoid liposarcoma (MLPS) with a *EWSR1/CHOP* frequency < 5% (8,9), desmoplastic small round cell tumor (DSRCT) with a *EWSR1/WT1* frequency of 96 - 97% (10,11), and myoepithelial tumors (ME) with a *EWSR1* translocation frequency of 45% (3).

ES demonstrates differential incidence according to race. People of European ancestry are much more frequently affected compared to people of African and East Asian ancestry (12,13). In addition, differences in clinical presentation and outcome by race have been previously demonstrated in this disease (14). The impact of race on the incidence of other mesenchymal tumors associated with *EWSR1* translocations has not been studied.

Given the difference in ES incidence between people of European, East Asian, and African ancestry, we hypothesized that race might also have an impact on the incidence of other mesenchymal tumors associated with *EWSR1* translocation. If the incidence of these other *EWSR1* tumors shows a similar racial distribution to ES, then the implication would be that the propensity for *EWSR1* to undergo genetic translocation varies in a uniform manner by race. To evaluate for incidence differences among patients diagnosed with mesenchymal tumors associated with *EWSR1* translocation, we performed a secondary data analysis of all cases of these tumors registered on the Surveillance, Epidemiology, and End Results (SEER) public-access database collected from various geographic areas in the United States.

Methods

The SEER Program of the National Cancer Institute (NCI) collects and publishes information on cancer incidence and survival in the United States (15). Currently, these data from 17 population-based cancer registries cover approximately 28 percent of the US population from diverse geographic regions. The population covered by SEER is comparable to the general US population with regard to measures of poverty and education, but tends to be somewhat more urban (15,16). The proportion of foreign-born persons is higher than the general US population (17). The percentages of the white, African-American, Asian-American, and Native American populations represented in the SEER registry are 25%, 26%, 54%, and 43%, respectively (18).

ICD-O-2 and ICD-O3 diagnosis codes were used to identify patients diagnosed with ES (9260/3), CCS (9044/3), EMC (9231/3), and MLPS (8852/3) from 1973 – 2007 (19). The search for cases of DSRCT (8806/3) and ME tumors (8982/3) began in the year in which these entities were first described (1991 and 1997, respectively) and ended in 2007. While angiomatoid fibrous histiocytoma (AFH) is also reported to harbor *EWSR1* translocations (20,21), cases of AFH were only rarely reported in SEER (n=4) and therefore AFH was not further analyzed in this study. SEER classifies race into 28 mutually exclusive groups by using information from the medical record. SEER further classifies race into the major categories: “White”; “Black”; and “Others”, with the latter group including people of American Indian/Alaskan native and Asian/Pacific Islander ancestry. An additional race category includes “other unspecified” cases.

The rate session feature of SEER*Stat version 6.6.2 was used to calculate age-adjusted annual incidences rates and incidence rate ratios according to the three major race categories

(22). Age-adjusted rates are a weighted average of the crude rates, where the weights are the proportions of persons in the corresponding age groups of a standard population. Incidence rate ratios were calculated comparing age-adjusted incidence rates in either the SEER “Black” or “Other” (American Indian/Alaskan native and Asian/Pacific Islander) race categories compared to the “White” race category (reference group). These incidence rate ratios were expressed with 95% confidence intervals determined by SEER*Stat using the method described by Fay and Feuer (23).

Results

The search for mesenchymal tumors associated with *EWSR1* translocation in the SEER database identified a total of 2893 cases. The number of cases for each tumor entity and the overall age-adjusted annual incidence rates are shown in Table 1.

The age-adjusted annual incidence rates for several of these tumors differed according to race. Table 2 compares the incidence in the African-American population with the incidence in the white population. As expected, the most striking difference in incidence between white and African-American populations was noted in ES. The age-adjusted annual incidence of ES was 1.6 cases/1,000,000 in the white population compared to 0.2 cases/1,000,000 in the African-American population, yielding an incidence rate ratio of 0.12 (95% confidence interval 0.08 – 0.20; $p < 0.001$). DSRCT showed the opposite pattern, with a higher age-adjusted annual incidence rate in the African-American population (0.3 cases/1,000,000) compared to the white population (0.1 cases/1,000,000 yielding an incidence rate ratio of 3.0 (95% confidence interval 1.62 – 5.49; $p < 0.001$). MLPS was less common in the African-American population (age-adjusted annual incidence rate of 1.4 cases/1,000,000) compared to the white population (age-adjusted annual incidence rate of 1.6 cases/1,000,000), though this difference was not statistically significant (incidence rate ratio of 0.83; 95% confidence interval 0.66 – 1.03; p -value = 0.1). EMC, CCS and ME had comparable age-adjusted annual incidence rates that did not differ significantly between white and African-American populations.

Differences in age-adjusted annual incidence rates were also observed between the white population compared to the “other” race category that includes the American Indian/Alaskan native and Asian/Pacific Islander populations (Table 2). ES demonstrated clear differences in incidence between white and the “other” race category. The age-adjusted annual incidence of 0.9 cases/1,000,000 in the “other” race category was significantly lower compared to 1.6 cases/1,000,000 in the white population, yielding an incidence rate ratio of 0.54 (95% confidence interval 0.41 – 0.69; $p < 0.001$). Interestingly, an incidence rate ratio of 0.72 (95% confidence interval 0.56 – 0.92; $p = 0.006$) for MLPS indicated that this tumor was also significantly less common among the American Indian/Alaskan Native/Asian and Pacific Islander population (age-adjusted annual incidence rate of 1.2 cases/1,000,000) compared to the white population (age-adjusted annual incidence rate of 1.6 cases/1,000,000). There were no differences in age-adjusted annual incidence for DSRCT, CCS, EMC, and ME between the white and “other” race categories.

Discussion

This is the first comprehensive analysis of racial differences in the incidence of mesenchymal tumors associated with *EWSR1* gene translocations. The results confirm previous reports of strong racial differences in the incidence of Ewing sarcoma and provide the first description of significant racial differences in the incidence of both DSRCT and MLPS. Specifically, DSRCT was significantly more common in the African-American population compared to the white population. In addition, MLPS had a lower incidence

among the American Indian/Alaskan native and Asian/Pacific Islander population compared to the white population. These findings suggest that the relationship between *EWSR1* translocation and race is complex.

For various cancer histologies, race related incidence differences have been demonstrated (24-26). One of the few risk factors for ES is race, even among people that have emigrated from their continents of origin (12,13). This finding suggests a genetic component of this disease. In addition, this pattern raises the possibility that people of European ancestry have an overall higher propensity to undergo somatic translocation at the *EWSR1* locus compared to people of African or Asian ancestry. The contrasting pattern of incidence by race between ES and DSRCT argues strongly against this possibility. Our analyses indicate that ES is 8 times more likely to occur in the white population compared to African-Americans and 1.9 times more likely to occur in the white population compared to Asian-Americans and Native Americans. The opposite was demonstrated for DSRCT, which is 3 times more likely to occur among African-Americans compared to whites. *EWSR1* translocations are present in a very high percentage of both of these tumors (1,10). Nevertheless, these tumors show opposite incidence rate ratios between white and African-American populations. The current findings suggest that differential propensity to translocation at the *EWSR1* locus may not be responsible for the differential incidence of these tumors between racial groups. However, it is important to note that the specific breakpoints in the *EWSR1* gene may differ between ES and DSRCT (1,10). Even among tumors of the same histology, the specific breakpoints or fusion partners may differ by race, which could not be evaluated in the current study. As such, it therefore remains possible that genetic differences at specific sites within the *EWSR1* gene may vary by race and mediate the differences observed in the current study.

The main strength of the use of the SEER database is the high number of analyzed mesenchymal tumor cases, which are rarely diagnosed. As with other retrospective population-based studies, this analysis comes with major limitations. First, race and tumor histology could not be independently confirmed. Second, data regarding potential environmental factors that might influence the observed differences are not available in SEER. Finally, information about genetic alterations of the tumor is not available in SEER. Instead, the tumor histology provided by SEER was viewed as a proxy for the presence of an *EWSR1* translocation. This assumption may be appropriate for tumors such as ES and DSRCT that show nearly universal *EWSR1* translocation (1,10). In contrast, this assumption may be less appropriate for ME and particularly MLPS, which harbor *EWSR1* gene rearrangements in only in a minority of cases (2,3,8,9). ME and MLPS were included in the current analysis to provide a complete evaluation of the role of race in tumors reported to be associated with *EWSR1* translocation. Despite the low frequency of *EWSR1* translocation in MLPS, our findings of differences in incidence according to race should prompt further investigations into their genetic epidemiology.

Despite these limitations, the significant differences in incidence rates according to race suggest that genetic differences may underlie the pathogenesis of these tumors. However, these tumors do not appear to share common racial differences in the propensity for *EWSR1* translocation. Additional study of the genetic epidemiology of these rare tumors will provide further insight into their pathogenesis.

Acknowledgments

Support: Supported by the Campini Foundation and NIH/NCRR/OD UCSF-CTSI Grant Number KL2 RR024130. The contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH.

References

1. Delattre O, Zucman J, Melot T, et al. The Ewing family of tumors--a subgroup of small-round-cell tumors defined by specific chimeric transcripts. *N Engl J Med* 1994;331:294–9. [PubMed: 8022439]
2. Romeo S, Dei Tos AP. Soft tissue tumors associated with EWSR1 translocation. *Virchows Arch* 456:219–34. [PubMed: 19936782]
3. Antonescu CR, Zhang L, Chang NE, et al. EWSR1-POU5F1 fusion in soft tissue myoepithelial tumors. A molecular analysis of sixty-six cases, including soft tissue, bone, and visceral lesions, showing common involvement of the EWSR1 gene. *Genes Chromosomes Cancer*.
4. Coindre JM, Hostein I, Terrier P, et al. Diagnosis of clear cell sarcoma by real-time reverse transcriptase-polymerase chain reaction analysis of paraffin embedded tissues: clinicopathologic and molecular analysis of 44 patients from the French sarcoma group. *Cancer* 2006;107:1055–64. [PubMed: 16878328]
5. Hisaoka M, Ishida T, Kuo TT, et al. Clear cell sarcoma of soft tissue: a clinicopathologic, immunohistochemical, and molecular analysis of 33 cases. *Am J Surg Pathol* 2008;32:452–60. [PubMed: 18300804]
6. Wang WL, Mayordomo E, Zhang W, et al. Detection and characterization of EWSR1/ATF1 and EWSR1/CREB1 chimeric transcripts in clear cell sarcoma (melanoma of soft parts). *Mod Pathol* 2009;22:1201–9. [PubMed: 19561568]
7. Panagopoulos I, Mertens F, Isaksson M, et al. Molecular genetic characterization of the EWS/CHN and RBP56/CHN fusion genes in extraskeletal myxoid chondrosarcoma. *Genes Chromosomes Cancer* 2002;35:340–52. [PubMed: 12378528]
8. Antonescu CR, Elahi A, Healey JH, et al. Monoclonality of multifocal myxoid liposarcoma: confirmation by analysis of TLS-CHOP or EWS-CHOP rearrangements. *Clin Cancer Res* 2000;6:2788–93. [PubMed: 10914725]
9. Antonescu CR, Tschernyavsky SJ, Decuseara R, et al. Prognostic impact of P53 status, TLS-CHOP fusion transcript structure, and histological grade in myxoid liposarcoma: a molecular and clinicopathologic study of 82 cases. *Clin Cancer Res* 2001;7:3977–87. [PubMed: 11751490]
10. Gerald WL, Ladanyi M, de Alava E, et al. Clinical, pathologic, and molecular spectrum of tumors associated with t(11;22)(p13;q12): desmoplastic small round-cell tumor and its variants. *J Clin Oncol* 1998;16:3028–36. [PubMed: 9738572]
11. Lae ME, Roche PC, Jin L, Lloyd RV, Nascimento AG. Desmoplastic small round cell tumor: a clinicopathologic, immunohistochemical, and molecular study of 32 tumors. *Am J Surg Pathol* 2002;26:823–35. [PubMed: 12131150]
12. Guo W, Xu W, Huvos AG, Healey JH, Feng C. Comparative frequency of bone sarcomas among different racial groups. *Chin Med J (Engl)* 1999;112:1101–4. [PubMed: 11721448]
13. Fraumeni JF Jr, Glass AG. Rarity of Ewing's sarcoma among U.S. Negro children. *Lancet* 1970;1:366–7. [PubMed: 4189632]
14. Worch J, Matthay KK, Neuhaus J, Goldsby R, DuBois SG. Ethnic and racial differences in patients with Ewing sarcoma. *Cancer* 116:983–8. [PubMed: 20052725]
15. Hankey BF, Ries LA, Edwards BK. The surveillance, epidemiology, and end results program: a national resource. *Cancer Epidemiol Biomarkers Prev* 1999;8:1117–21. [PubMed: 10613347]
16. SEER. NIH Publication No 05-4772. 2005. Surveillance Epidemiology, and End Results Program.
17. Wara WM, Jenkin RD, Evans A, et al. Tumors of the pineal and suprasellar region: Childrens Cancer Study Group treatment results 1960–1975: a report from Childrens Cancer Study Group. *Cancer* 1979;43:698–701. [PubMed: 105801]
18. Silber JH, Radcliffe J, Peckham V, et al. Whole-brain irradiation and decline in intelligence: the influence of dose and age on IQ score. *J Clin Oncol* 1992;10:1390–6. [PubMed: 1517781]
19. Fritz, A.; Percy, C.; Jack, Andrew, et al. International Classification of Diseases for Oncology. 3rd. World Health Organization; Geneva: 2000.
20. Rossi S, Szuhai K, Ijszenga M, et al. EWSR1-CREB1 and EWSR1-ATF1 fusion genes in angiomatoid fibrous histiocytoma. *Clin Cancer Res* 2007;13:7322–8. [PubMed: 18094413]

21. Tanas MR, Rubin BP, Montgomery EA, et al. Utility of FISH in the diagnosis of angiomatoid fibrous histiocytoma: a series of 18 cases. *Mod Pathol* 23:93–7. [PubMed: 19801966]
22. SEER. www.seer.cancer.gov/seerstat.
23. Fay MP, Feuer EJ. Confidence intervals for directly standardized rates: a method based on the gamma distribution. *Stat Med* 1997;16:791–801. [PubMed: 9131766]
24. Haynes, MA.; Smedley, BD. The unequal burden of cancer: An Assessment of NIH Research and Programs for Ethnic Minorities and the Medically Underserved. Washington, DC: NATIONAL ACADEMY PRESS; 1999.
25. Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57:43–66. [PubMed: 17237035]
26. Bhatia S, Sather HN, Heerema NA, Trigg ME, Gaynon PS, Robison LL. Racial and ethnic differences in survival of children with acute lymphoblastic leukemia. *Blood* 2002;100:1957–64. [PubMed: 12200352]

Table 1

Age-adjusted annual incidence rate independent of race for mesenchymal tumors associated with *EWSR1* translocation.

Histology*	Age-Adjusted Annual Incidence Rate (cases per 1,000,000)
Ewing sarcoma (n = 1185)	1.4
Myxoid liposarcoma (n = 1236)	1.6
Extraskelatal myxoid chondrosarcoma (n = 214)	0.3
Clear cell sarcoma (n = 138)	0.2
Myoepithelial tumor (n = 59)	0.2
Desmoplastic small round cell tumor (n = 61)	0.1

* Numbers reflect incident cases reported to SEER from 1973-2007, except for desmoplastic small round cell tumor (1991-2007) and myoepithelial tumors (1997 - 2007).

Table 2

Age-adjusted annual incidence rates for tumors associated with *EWSR1* translocations in white, African American and Asian/Native American populations.

Histology**	<i>EWSR1</i> Translocation		Age-Adjusted Annual Incidence Rate (cases per 1,000,000)			Rate Ratio* (95% CI)	
	Most Common Fusion Partner	<i>EWSR1</i> Translocation Frequency from Literature	White	African-American	Asian/Native American	African American	Asian/Native American
Ewing sarcoma	<i>FLI1</i>	95%	1.6	0.2	0.9	0.12 (0.08 – 0.2)	0.54 (0.41 – 0.69)
Myxoid liposarcoma	<i>CHOP</i>	< 5%	1.6	1.4	1.2	0.83 (0.66 – 1.03)	0.72 (0.56 – 0.92)
Extraskeletal myxoid chondrosarcoma	<i>CHN</i>	85%	0.3	0.3	0.2	1.14 (0.67 – 1.82)	0.58 (0.27 – 1.1)
Clear cell sarcoma	<i>ATF1</i>	87-94%	0.2	0.2	0.2	1.12 (0.59 – 2.07)	1.02 (0.52 – 1.84)
Myoepithelial tumor	<i>POU5F1</i>	45%	0.2	0.3	0.3	1.46 (0.62 – 3.15)	1.75 (0.74 – 3.67)
Desmoplastic small round cell tumor	<i>WT1</i>	96-97%	0.1	0.3	0.1	3.0 (1.62 – 5.49)	1.04 (0.36 – 2.6)

* Ratio of incidence rate in the African American or Asian/Native American population to incidence rate in the white population (reference).