Published in final edited form as:

Invert Neurosci. 2007 March ; 7(1): 17–30. doi:10.1007/s10158-006-0036-9.

Insect sodium channels and insecticide resistance

Ke Dong

Department of Entomology, Genetics Program and Neuroscience Program, Michigan State University, East Lansing, MI 48824, USA

Ke Dong: dongk@msu.edu

Abstract

Voltage-gated sodium channels are essential for the generation and propagation of action potentials (i.e., electrical impulses) in excitable cells. Although most of our knowledge about sodium channels is derived from decades of studies of mammalian isoforms, research on insect sodium channels is revealing both common and unique aspects of sodium channel biology. In particular, our understanding of the molecular dynamics and pharmacology of insect sodium channels has advanced greatly in recent years, thanks to successful functional expression of insect sodium channels in *Xenopus* oocytes and intensive efforts to elucidate the molecular basis of insect resistance to insecticides that target sodium channels. In this review, I discuss recent literature on insect sodium channels with emphases on the prominent role of alternative splicing and RNA editing in the generation of functionally diverse sodium channels in insects and the current understanding of the interactions between insect sodium channels and insecticides.

Keywords

Para; *DSC1*; Sodium channel; Knockdown resistance; Alternative splicing; RNA editing

Introduction

Voltage-gated sodium channels are integral transmembrane proteins responsible for the rapidly rising phase of action potentials, and they are critical for electrical signaling in most excitable cells. In response to membrane depolarization, sodium channels open (activate) and allow sodium ions to flow into the cell, thereby depolarizing the membrane potential. A few milliseconds after channel activation, the channel pore is occluded, a process known as fast inactivation, which is partially responsible for the falling phase of the action potential.

Because of their crucial role in membrane excitability, sodium channels are the target site of a great variety of neurotoxins, such as tetrodotoxin, scorpion toxins, and batrachotoxin, which are produced by plants and animals for defense or predation (Cestele and Catterall 2000; Wang and Wang 2003). Insecticidal pyrethrins, found in extracts of the flowers of *Chrysanthemum* species, also act on sodium channels (Narahashi 1988). Sodium channels are also the primary target of DDT and modern synthetic pyrethroids, which are structural derivatives of the naturally occurring pyrethrins (Narahashi 1988). Furthermore, recent evidence indicates that a new class of pyrazoline- like insecticides, oxadiazines, also target sodium channels (Wing et al. 2005; Silver and Soderlund 2005a).

Due to intensive use of insecticides in arthropod control, many pest populations have developed resistance to these compounds. One major mechanism of resistance is known as knockdown resistance (*kdr*). Insects exhibiting *kdr* have reduced target-site (sodium channel) sensitivity to pyrethroids and DDT resulting from one or more point mutations in the insect sodium channel protein. The important role of sodium channels in insecticide resistance has prompted many researchers to study insect sodium channel biology. In particular, the past decade has witnessed significant accumulation of knowledge on the functional diversity and pharmacology of insect sodium channels. In this review, I will focus on recent literature that sheds light on the mechanisms of generating functionally diverse sodium channels from a single gene in insects and the current understanding of the interactions between insect sodium channels and insecticides, such as pyrethroids and indoxacarb. Readers are referred to an excellent and complementary review on this topic (Soderlund 2005).

Structure and function of sodium channels derived from mammalian systems

Extensive molecular analysis of mammalian sodium channels in the last two decades has generated comprehensive views into the structure and function of voltage-gated sodium channels (Catterall 2000; Yu and Catterall 2003; Goldin 2001, 2003). Mammalian sodium channels consist of a large pore-forming transmembrane α -subunit and several small auxiliary β -subunits. Expression of the α -subunit alone in *Xenopus* oocytes or mammalian cell lines is sufficient to obtain sodium currents, and the auxiliary β -subunits seem mainly to modulate sodium channel gating and/or protein expression. The α -subunit contains four homologous domains (named I–IV), each having six membrane spanning segments (named S1–S6) connected by intracellular or extracellular loops of amino acid sequences (Fig. 1). The selectivity filter and pore are formed by transmembrane segments S5 and S6 together with the membrane-reentrant segments that are part of the loop connecting S5 and S6 of each domain (Fig. 1). Amino acids D, E, K, and A in the loop connecting S5 and S6 of domains I, II, III, and IV, respectively, are the key determinants of ion selectivity. Each S4 segment has five to eight basic residues, either arginine or lysine, separated from each other by two neutral residues. These positively charged S4 segments function as voltage sensors, which move outward in response to membrane depolarization to initiate the opening of the sodium channel. The hydrophobic IFMT motif in the middle of the short linker connecting domains III and IV serves as an inactivation gate and is important for causing fast inactivation by binding to the inactivation gate receptor located within or near the intracellular mouth of the sodium channel pore. Several residues in the intracellular loop connecting S4 and S5 (i.e., IIIS4–S5 and IVS4–S5) are believed to be part of the docking receptor for the inactivation gate.

At least nine different sodium channel α -subunits (Nav1.1 to Nav1.9) have been identified in mammals (Goldin et al. 2000; Catterall et al. 2003). These sodium channel isoforms exhibit distinct expression patterns in the central and peripheral nervous systems and skeletal and cardiac muscle (Goldin 1999). Nav1.1, Nav1.2, Nav1.3, and Nav1.6 are predominately expressed in the central nervous system (CNS), whereas Nav1.7, Nav1.8, and Nav1.9 are expressed in the peripheral nervous system (PNS). Nav1.4 and Nav1.5 are expressed in skeletal and cardiac muscles, respectively. Mammalian sodium channel α -subunit isoforms also exhibit distinct electrophysiological properties (Goldin 2001; Dib-Hajj et al. 2002; Yu and Catterall 2003). Selective expression of different sodium channel genes (with unique gating properties) contributes to the specialized function of sodium channels in various mammalian tissues and cell types (Yu and Catterall 2003).

Structure and function of insect sodium channels

One or two insect sodium channel genes?

In the late 1980s, two putative sodium channel genes, *DSC1* and *para*, were isolated from *Drosophila melanogaster*. *DSC1* was isolated by Larry Salkoff and associates by screening a *D. melanogaster* genomic DNA library using an eel sodium channel gene probe (Salkoff et al. 1987). A *DSC1* orthologous gene was later identified from the German cockroach (*Blattella germanica*) and named *BSC1* (Liu et al. 2001). Like sodium channel α -subunits, *DSC1* and *BSC1* have four homologous domains, each having six transmembrane segments. Until very recently, *DSC1* and *BSC1* had been predicted to encode a voltage-gated sodium channel based on their overall similarity of deduced amino acid sequence and domain organization to eel and mammalian sodium channel proteins (Littleton and Ganetzky 2000). However, a recent study showed that *BSC1* and *DSC1* encode a novel family of Ca^{2+} -selective cation channels, not sodium channels (Zhou et al. 2004; W. Song and K. Dong, unpublished results). All voltage-gated sodium channels, including insect sodium channels, contain amino acids D, E, K, and A in the loops connecting S5 and S6 of domains I, II, III, and IV, respectively, whereas voltage-gated calcium channels contain acidic residues (E, E, E, and E) at the corresponding positions (See Table 2 in Zhou et al. 2004). Site-directed mutagenesis has shown that the E residue in domain III is a critical determinant of the Ca^{2+} selectivity of the *BSC1* channel (Zhou et al. 2004). Interestingly, several other putative sodium channels in invertebrates, including ApCSN1 (*Aiptasia pallida*, sea anemone), CYNA1 (*Cyanea capillata*), PpSCN1 (*Polyorchis penicillatus*, Hydrozoan jellyfish), and LbNa_v1 (*Loligo bleekeri*, squid), all possess an E in domain III (Refs. in Zhou et al. 2004). Therefore, it seems likely that the other invertebrate *BSC1/DSC1*-like channels may also exhibit Ca^{2+} selectivity, although functional expression of these channels remains to be established. The biological role of *DSC1* and *BSC1* channels in insects also remains to be determined. However, a recent study showed that partially reduced expression of the *DSC1* gene was correlated with certain olfactory defects in *D. melanogaster* (Kulkarni et al. 2002), suggesting a role of *DSC1* in olfaction.

The *para* gene was isolated in Barry Ganetzky's laboratory from mutants exhibiting a temperaturesensitive paralytic phenotype (Loughney et al. 1989). The overall structure of and the amino acid sequence of the Para sodium channel shares a high similarity with those of mammalian sodium channel α -subunits. The structural features that are critical for mammalian sodium channel function, including residues crucial for sodium selectivity, are conserved in the Para sodium channel (Loughney et al. 1989). Most importantly, subsequent functional expression and characterization in *Xenopus* oocytes conclusively demonstrated that *para* encodes a sodium channel (Feng et al. 1995; Warmke et al. 1997). Except for significant sequence similarities with the *DSC1* channel protein and limited sequence similarity with putative Ca^{2+} channel $\alpha 1$ subunits, Para is not similar to any other genes in the *Drosophila* genome (Littleton and Ganetzky 2000). Therefore, it seems that *para* is the only gene that encodes the sodium channel in *Drosophila* and presumably in other insect species as well.

Because of the intimate involvement of sodium channels in insecticide resistance (see below), *para*-orthologous genes have been isolated from several medically or agricultural important insect pest species (Soderlund and Knipple 2003). However, in most cases, only partial cDNA clones were obtained. Full-length cDNA clones are available only for three *para* orthologues: *Vssc1* from the house fly (Ingles et al. 1996), *BgNa_v* from the German cockroach (Dong 1997); and *VmNa_v* from the varroa mite (Wang et al. 2003). The availability of the full-length clones made it possible to successfully express these *para* orthologues in *Xenopus* oocytes and to demonstrate that these genes indeed encode

functional sodium channels (Ingles et al. 1996; Tan et al. 2002a; Song et al. 2004; Du et al., unpublished).

TipE: an auxiliary subunit?

Robust expression of insect sodium channels in *Xenopus* oocytes requires a small transmembrane protein, TipE, which has two transmembrane segments connected by a large extracellular loop and intracellular amino and carboxyl termini (Feng et al. 1995). Coexpression of TipE with Para in oocytes increases the amplitude of peak current and modifies the kinetics of fast inactivation (Feng et al. 1995; Warmke et al. 1997), suggesting that TipE functions as an auxiliary subunit like the β -subunits of mammalian sodium channels. A TipE ortholog cloned from the house fly appears to fulfill the same role as TipE (Lee et al. 2000). TipE also enhances the expression of BgNa_v in oocytes (Tan et al. 2002a). These results suggest the functional conservation of these transmembrane proteins in different insect species.

Recently, four TipE homologs (TEH1, 2, 3, 4) are found in *D. melanogaster* and their predicted secondary structures resemble those of TipE (Derst et al. 2006). Like TipE, TEH1, 2, and 3, but not TEH4, increase the expression of a Para variant in oocytes (Derst et al. 2006). TEH1 also alters voltage-dependent inactivation and recovery from inactivation of the Para variant. Interestingly, except for TEH1, which is only expressed in the central nervous system, TEH2, 3, 4 are widely expressed in both neuronal and nonneuronal tissues. The involvement of these proteins in regulating sodium or other ion currents in insect tissues/ cells remains to be investigated.

Extensive alternative splicing of insect sodium channel gene transcripts

Alternative splicing of the *para* transcript was first reported in the *D. melanogaster* (Thackeray and Ganetzky 1994). To date, a total of nine alternatively splice sites—a, b, c/d, i, j, e, f, h, l/k—have been identified in *para* (Fig. 1; Thackeray and Ganetzky 1994, 1995; O'Dowd et al. 1995; Lee et al. 2002). Exons, a, b, i, j, e, f, and h, are optional and are located in the intracellular linkers, whereas the two mutually exclusive exons c/d and l/k are situated in the transmembrane regions. Interestingly, most of these alternative splice sites are conserved in *D. virilis* (Thackeray and Ganetzky 1995), the house fly *Vssc1* gene (Lee et al. 2002), and the cockroach *BgNa_v* gene (Tan et al. 2002a; Song et al. 2004). Such a high degree of conservation, however, does not seem to extend to beyond insect sodium channels. For example, only two of the splice sites, corresponding to b and l/k in *Drosophila*, are found in the *VmNa_v* (varroa mite sodium channel) transcript, which has three unique splice sites (Wang et al. 2003; Y. Du et al., unpublished results). Although the splice sites are highly conserved in insect species, the frequency of exon usage can be quite different. For example, more than 60% of the *para* (both *D. melanogaster* and *D. virilis*) and *Vssc1* transcripts contain exon b, but less than 20% of BgNa_v transcripts contain exon B (Capital letters are used to denote alternative exons in BgNa_v).

At the position corresponding to the *Drosophila* l/k splice site, which is located in IIIS3–S4, there are three mutually exclusive exons, called G1, G2, and G3, in *BgNav* (in Fig. 1; Tan et al. 2002b). Exons G1 and G2 share 65% amino acid sequence homology, and correspond to exons l and k in *Drosophila*, respectively. In contrast, exon G3 has no sequence homology with either G1 or G2, and possesses a stop codon that would generate a truncated protein containing only the first two domains. Interestingly, at this conserved splicing site, *VmNa_v* contains an optional alternative exon (named exon 3) that encodes IIIS3–S4 (Wang et al. 2003). Because IIIS4 is essential for sodium channel function, exclusion of exon 3 likely also generates a non-functional channel. Furthermore, the exon/intron arrangement of this region in insects bears a striking similarity with that of the genomic region coding for IIIS3–

S4 of the vertebrate sodium channel protein, Na_v1.6 (Scn8a), of mice, humans, and fish (Plummer et al. 1997). Like exon G3 in *BgNa_v*, the mammalian counterparts possess a stop codon that would generate a truncated protein containing only the first two domains. Therefore, alternative splicing in this region appears to be of a very ancient origin and is preserved in organisms ranging from mites and insects to humans.

RNA editing of insect sodium channel transcripts

RNA editing is a well-known post-transcriptional modification that could significantly change protein function by introducing site-specific alterations in gene transcripts. RNA editing results in the conversion of one base to another, or the insertion and deletion of nucleotides (Bass 2001). Both A-to-I and U-to-C editing events have been reported for insect sodium channel genes. Ten A-to-I editing sites are identified in the *para* transcript in *D. melanogaster* by direct sequencing or restriction enzyme digestion analysis of partial cDNA clones (Palladino et al. 2000; Hanrahan et al. 2000; Reenan et al. 2000). The actual number of RNA editing sites may be much greater than ten, because the region examined in these studies represents only a portion of the complete *para* open reading frame. Eight of the ten editing events result in amino acid changes in the transmembrane segments or intracellular linkers (Fig. 2). Some of these RNA editing sites are conserved in *D. virilis* (Hanrahan et al. 2000); however, possible functional alterations caused by these amino acid changes remain to be examined. Surprisingly, none of these A-to-I editing sites are found in *BgNa_v*. Instead, two unique A-to-I RNA editing sites and three U-to-C editing sites are identified in *BgNa_v*, each resulting in amino acid changes in the transmembrane segments (Fig. 2; Song et al. 2004; Liu et al. 2004). One of the U-to-C editing sites, F/S¹⁹¹⁹, is also found in *para* (Liu et al. 2004). It is not clear whether similar RNA editing events occur in mammalian sodium channel transcripts.

Functional consequences of alternative splicing and RNA editing of insect sodium channels

As discussed above, selective expression of different sodium channel genes (with unique gating properties) contributes to the functional diversity of sodium channels in various mammalian tissues and cell types (Yu and Catterall 2003). Similarly, electrophysiological recordings in different insect neurons also revealed distinct sodium currents, indicating heterogeneity in sodium channel properties in insects (Byerly and Leung 1988; Saito and Wu 1991, 1993; Schafer et al. 1994; O'Dowd et al. 1995; Le Corronc et al. 1999; Lapied et al. 1990, 1999; Wicher et al. 2001; Grolleau and Lapied 2000; Defaix and Lapied 2005). It is still unclear how insects produce distinct sodium currents from a single sodium channel gene. One possibility is that alternative splicing and RNA editing generate sodium channels with unique gating properties. Recent functional characterization in *Xenopus* oocytes of several cockroach sodium channel variants that undergo splicing and/or RNA-editing suggests that this is indeed the case. For example, electrophysiological characterization of 20 *BgNa_v* splice variants in *Xenopus* oocytes revealed an impressive spectrum of differences in the level of sodium current expression and gating properties (e.g., the voltage-dependence of activation and/or inactivation) (Song et al. 2004). One variant, *BgNa_v11*, activates at more negative membrane potentials (e.g., -60 mV), whereas others require more membrane depolarization (e.g., -45 mV) for channel activation. Similarly, the half-maximal voltages for steady-state inactivation range from -37 to -60 mV. *BgNa_v7* has a significant overlap between the voltage dependences of activation and steady-state inactivation, with $V_{1/2}$ values of -36.7 and -37.2 mV, respectively. Consequently, this variant produces a large window current over a range of -45 to -25 mV, which could lead to membrane depolarization at subthreshold potentials associated with oscillatory activities, summation of synaptic input, or modulation of firing frequency (Crill 1996).

The role of specific alternative exons in modulating sodium channel gating has been demonstrated. Exon B in BgNa_v encodes an eight-amino acid sequence in the linker connecting domains I and II. BgNa_v variants containing exon-B expressed poorly in oocytes (Song et al. 2004), but deletion of exon-B caused the level of sodium current expression to increase dramatically (Song et al. 2004). Similarly, introduction of exon-B into variants lacking exon-B resulted in the reduction of peak sodium currents of these recombinant channels (Song et al. 2004). These results indicate exon-B plays a critical role in the regulation the level of sodium channel expression and/or the size of current in *Xenopus* oocytes. Considering that the location and sequence of this exon is highly conserved in *Drosophila para* and also in house fly *Vssc1* (Lee et al. 2002; Song et al. 2004), the role of exon b/B in regulating insect sodium current expression may be universal. Exon b/B, located in the first intracellular linker, contains a consensus sequence for phosphorylation. Down-regulation of sodium current expression by protein kinases has been documented for mammalian sodium channels (Numann et al. 1991; West et al. 1991; Smith and Goldin 1996, 1997, 2000). Exon b/B could serve as a regulatory on-or-off switch to control the neuronal excitability by responding to second messengers to meet unique physiological needs in specific tissues or cell types. In this regard, it is interesting to note that the majority of optional exons in insect sodium channels are located in intracellular linkers. These linker sequences seem ideal for interaction with other cellular proteins, such as G-proteins and kinases, and for integrating cell and tissue-specific signals to “customize” sodium channel properties.

The functional consequence of RNA editing of insect sodium channel transcripts has also been shown recently. A U-to-C RNA editing event at the F/S site at the C-terminal domain of BgNa_v generated a non-inactivating (persistent) Na⁺ current (Liu et al. 2004). Eliminating the U-to-C editing at the L/P site in IIS3 caused the BgNa_v1-1 variant to activate and inactivate at more hyperpolarizing membrane potentials. Similarly, abolishment of the A-to-I editing at the K/R site in IS2 shifted the voltage-dependence of activation of the BgNa_v1-2 variant to a more hyperpolarized potential. Clearly, these editing events are responsible for the different activation and inactivation properties of the BgNa_v1-1 and BgNa_v1-2 variants. The modulation of sodium channel gating properties by specific RNA editing events supports the hypothesis that RNA editing is involved in the fine-tuning of the neuronal activity (Seeburg 2000).

In short, recent studies suggest that the functional plasticity of the insect sodium channel is generated primarily by alternative splicing and RNA editing of a single gene transcript. Whereas mammals rely on multiple sodium channel genes to produce functionally distinct isoforms, insects seem to depend on extensive alternative splicing and RNA editing of a single sodium channel gene to presumably fulfill customized sodium channel activities in different cell types and tissues.

Sodium channels as insecticide targets

Sodium channels are targets of a variety of neurotoxins including naturally occurring toxins from animals and plants, therapeutic drugs, and synthetic insecticides pyrethroids and oxadiazines. These neurotoxins are grouped based on their distinct binding sites on the sodium channel and/or unique effects on sodium channel function. The extensive body of work on this topic is discussed in several excellent reviews (Gordon 1997; Zlotkin 1999; Cestele and Catterall 2000; Wang and Wang 2003).

Pyrethroids

Pyrethroids are synthetic structural derivatives of insecticidal pyrethrins present in the pyrethrum extract of *Chrysanthemum* species (Elliott 1977). They are grouped into two

categories (type I and type II) based on their distinct poisoning symptoms, effects on nerve preparations, and their chemical structures (Gammon et al. 1981; Lawrence and Casida 1982; for review see Narahashi 1986). Type I pyrethroids lack an α -cyano group which is present at the phenylbenzyl alcohol position of type II pyrethroids. Type I pyrethroids cause repetitive discharges in response to a single stimulus, while type II pyrethroids caused a membrane depolarization accompanied by a suppression of the action potential. Historically, studies on the mode of action of pyrethroids have been conducted using vertebrate and non-insect invertebrate nerve preparations. Collectively, these studies show that pyrethroids cause prolonged opening of sodium channels primarily by inhibiting channel deactivation and stabilizing the open configuration of the sodium channel (for review see Narahashi 1986, 1988, 1996, 2000; Soderlund and Bloomquist 1989; Raymond-Delpech et al. 2005). The prolonged channel opening is evidenced by a large tail current associated with repolarization under voltageclamp conditions. Furthermore, voltage-clamp experiments showed that type II pyrethroids inhibit the deactivation of sodium channels to a greater extent than type I pyrethroids. The decay of tail currents induced by type II pyrethroids is at least one order of magnitude slower than those induced by type I pyrethroids. These quantitative differences in tail-current decay kinetics between type I and type II pyrethroids may account for their different actions on the nervous system. However, the molecular basis of the differential effects of type I and type II pyrethroids on sodium channels remains to be determined.

Recent studies on the mechanism of action of pyrethroids on insect sodium channels expressed in oocytes and the molecular mechanism of *kdr* (see below) confirmed that sodium channels are the target of pyrethroid insecticides. Vais et al. (2000a) showed that deltamethrin, a potent type II pyrethroid, preferably bind to the activated (open) state and inhibit deactivation of Para sodium channels expressed in oocytes. Similar effects were also observed on house fly and cockroach sodium channels (Smith et al. 1997; Tan et al. 2002a, b; Tan et al. 2005). Furthermore, these studies also demonstrated that the effects of type I and type II pyrethroids on insect sodium channels are similar to those observed from electrophysiological studies using non-insect nerve preparations.

Oxadiazines

Commercialization of indoxacarb, a new insecticide, represents a success in the development of a new generation of sodium-channel-targeted insecticides (McCann et al. 2001). In insects, indoxacarb is metabolically converted to *N*-decarbomethoxylated JW062 (DCJW), a more active metabolite. Mammals convert indoxacarb into nontoxic metabolites, which contributes to its selective toxicity to insect pests.

Recent studies showed that the sodium channel is a major cellular target of indoxacarb and DCJW (Wing et al. 2005; Silver and Soderlund 2005a). For example, indoxacarb reduces the compound action potential in the *M. sexta* larval nerve cord preparation and inhibits firing of abdominal stretch receptor organs in larvae of the fall armyworm *Spodoptera frugiperda*, but the effect was slower than that of its metabolite, DCJW (Wing et al. 1998, 2005). Both indoxacarb and DCJW blocked sodium channels in patch-clamped neurons isolated from rat dorsal root ganglia (Nagata et al. 1998; Tsurubuchi and Kono 2003; Zhao et al. 2003) and from the central nervous system (CNS) of the American cockroach neurons (Lapied et al. 2001; Zhao et al. 2005). These studies suggest that the action of indoxacarb and DCJW share similarities with those of RH3421, a dihydropyrazole insecticide (Salgado 1992), and local anesthetics (LA), which block sodium currents in a state-dependent manner, by preferably binding to the inactivated state of the sodium channel (Hille 1977). Studies using the rat Na_v1.4 and cockroach sodium channel variants expressed in *Xenopus* oocytes confirmed the state-dependent block of both mammalian and insect sodium channels by DCJW (Silver and Soderlund 2005b; Song et al. 2006; Silver and Soderlund 2006).

Pyrethroid sensitivities among naturally occurring sodium channel variants

Differential sensitivities of mammalian sodium channel isoforms to pyrethroid insecticides are well known. For example, tetrodotoxin (TTX)-sensitive sodium channels in the rat dorsal root ganglion neurons are less sensitive to pyrethroids than TTX-resistant sodium channels in the same neurons (Ginsburg and Narahashi 1993; Tatebayashi and Narahashi 1994; Song and Narahashi 1996). Rat Na_v1.2 and Na_v1.4 channels are insensitive to pyrethroids (Warmke et al. 1997; Vais et al. 1997; Smith and Soderlund 1998; Wang et al. 2001), whereas rat Na_v1.8 sodium channels are more sensitive to pyrethroids (Choi and Soderlund 2006). Interestingly, substitution of an isoleucine in IIS4–S5 in the rat Na_v1.2 channel by a methionine (which is in insect sodium channels) increased channel sensitivity to pyrethroids (Vais et al. 2000b). However, the isoleucine residue is conserved in all known mammalian sodium channel protein, and is therefore cannot be responsible for the differential sensitivities of mammalian sodium channel isoforms. Clearly, an important future goal would be to determine other amino acid residues involved in modulating the pyrethroid sensitivity of various mammalian sodium channel isoforms.

The existence of sodium channel variants in insects that exhibit distinct levels of sensitivity to pyrethroids has been demonstrated recently. In cockroaches, the BgNa_v2-1 channel variant is 100-fold less sensitive to deltamethrin than the BgNa_v1-1 variant (Tan et al. 2002b). As described above, the BgNa_v2-1 channel contains a mutually exclusive exon G2, whereas the BgNa_v1-1 channel contains exon G1. The differential pyrethroid sensitivities between BgNa_v1-1 and BgNa_v2-1 channels can be partially attributed to the presence of exon G1 or G2 (Tan et al. 2002b). More recently, a V1356A change in exon G2 is identified to be responsible for the exon G2-associated low sensitivity of the BgNa_v2-1 channel to pyrethroids (Du et al. 2006). Because there are ten scattered exon-G-independent amino acid differences between BgNa_v1-1 and BgNa_v2-1, systematic site-directed mutagenesis are necessary to identify additional residues that are involved in the different sensitivity between the two variants. The BgNa_v2-1 variant appears to be a minor variant in cockroaches (Song et al. 2004; Z. Liu and K. Dong, unpublished result) and is expressed mainly in gut and ovary (Tan et al. 2002b). The insensitivity to pyrethroids of BgNa_v2-1 is likely caused by unique channel gating properties of this variant, which are presumably customized for specific roles in gut and ovary. Regardless of their precise roles in mammalian and insect neurophysiology, pyrethroid-resistant sodium channel isoforms or variants are proving to be a unique resource for the elucidation of the interaction between sodium channels and pyrethroids at the molecular level.

Pyrethroid resistance caused by naturally selected *kdr* mutations in the sodium channel

kdr mutations confer reduced neuronal sensitivity to pyrethroids and DDT in insects (Soderlund and Bloomquist 1990). In the early 1990s, several groups conducted genetic linkage analysis and showed that *kdr* mutations were linked to the *para*-orthologous genes in several insect species (Refs. in Soderlund and Knipple 2003). Subsequent molecular analyses led to the identification of multiple point mutations in the *para*-orthologous genes that are associated with *kdr* and *kdr*-type resistance to pyrethroids in these insects (Soderlund 2005). To experimentally confirm the involvement of *kdr*-associated mutations in pyrethroid resistance, a number of laboratories used the *Xenopus* oocyte expression system to examine the sensitivity of mutant sodium channels to pyrethroids. To this end, ten sodium channel mutations have been confirmed to be responsible for *kdr* and *kdr*-type resistance (Fig. 3).

The most common *kdr* mutation is a leucine (L) to phenylalanine (F), histidine (H), or serine (S) substitution in domain II segment 6 (IIS6), associated with low levels of pyrethroid resistance. The L to F mutation is present in *kdr* house flies, *Musca domestica* (Williamson et al. 1996; Miyazaki et al. 1996); German cockroaches, *Blattella germanica* (Miyazaki et al. 1996; Dong 1997; Dong et al. 1998); diamondback moths, *Plutella xylostella* (Martinez-Torres et al. 1997; Martinez-Torres et al. 1999); peach-potato aphids, *Myzus persicae* (Schuler et al. 1998; Martinez-Torres et al. 1999a); mosquitoes, *Anopheles gambiae* (Martinez-Torres et al. 1998; Ranson et al. 2000), *Culex pipiens* (Martinez-Torres et al. 1999b) and *Culex quinquefasciatus* (Xu et al. 2005); horn flies, *Haematobia irritans* (Guerrero et al. 1997); Colorado potato beetles, *Leptinotarsa decemlineata* (Lee et al. 1999a); western flower thrips, *Frankliniella occidentalis* (Forcioli et al. 2002); codling moth, *Cydia pomonella* (Brun-Barale et al. 2005); and cat fleas, *Ctenocephalides felis* (Bass et al. 2004). The L to S mutation is found in *C. pipiens* (Martinez-Torres et al. 1999b) and *A. gambiae* (Ranson et al. 2000). The L to H mutation is present in tobacco budworms, *Heliothis virescens* (Park and Taylor 1997). In *Xenopus* oocytes, the L to F/H mutation in IIS6 reduces the pyrethroid sensitivity of Para, Vssc1 and BgNa_v sodium channels by five- to ten-fold (Smith et al. 1997; Zhao et al. 2000; Vais et al. 2000a; Tan et al. 2002a; Liu et al. 2002).

In highly resistant strains, additional sodium channel mutations are often found to co-exist with the L to F mutation in IIS6. For example, in *super-kdr* house fly strains, which exhibit greater levels of resistance to DDT and pyrethroids than *kdr* houseflies, a methionine (M) to threonine (T) mutation (M918T) in the linker connecting IIS4 and IIS5 is present in addition to the L to F mutation (L1014F) in IIS6 (Williamson et al. 1996; Miyazaki et al. 1996). The same M to T mutation together with the L to F mutation is also found in a horn fly strain that exhibited high-level resistance to pyrethroids (Guerrero et al. 1997). In *Xenopus* oocytes, the double mutations drastically reduced the sensitivity of the house fly and *Drosophila* sodium channels to pyrethroids (Lee et al. 1999c; Vais et al. 2000a). Interestingly, the M918T mutation alone also reduces sodium channel sensitivity to pyrethroids to some extent (Lee et al. 1999c; Vais et al. 2001), but it has not been found alone in any population of pyrethroid-resistant insects.

In the German cockroach, additional resistance mutations have been discovered that reduce the sensitivity of sodium channels in conjunction with the L to F mutation mentioned above. Four point mutations, aspartic acid to glycine (D58G) at the N- terminus, glutamic acid to lysine (E434K) and cysteine to arginine (C764R) in the linker between IS6 and IIS1, and proline to leucine (P1888L) at the C-terminus, together with the L to F mutation (L993F) in IIS6, are associated with high levels of resistance (Liu et al. 2000). Unlike the housefly M918T mutation, E434K and C764R by themselves do not reduce the sensitivity of the cockroach sodium channel to pyrethroid in *Xenopus* oocytes. However, when either the E434K or C764R mutation is combined with the L993F mutation (i.e., KF or RF), the channel sensitivity is reduced by 100-fold (Tan et al. 2002a), and concomitant presence of all three mutations (KRF) reduces channel sensitivity to pyrethroid by 500-fold (Tan et al. 2002a). Whether the two mutations (D58G and P1888L) at the N- and C- terminus contribute to pyrethroid resistance remains to be determined.

In a highly resistant *P. xylostella* strain, a threonine to isoleucine (T929I) mutation in IIS5 is found along with the L to F mutation in IIS6 (Schuler et al. 1998). The T929I mutation alone drastically reduces the Para sodium channel to pyrethroids and is considered a *super-kdr* mutation (Vais et al. 2001). At the same position, a T to C change and a T to V change is detected in western flower thrips (Forcioli et al. 2002) and a highly pyrethroid-resistant cat flea population (Bass et al. 2004), respectively. The same T to I mutation is also found in pyrethroid-resistant head lice, *Pediculus capitis* along with two more mutations, M827I and

L932F (Lee et al. 2000, 2003; Tomita et al. 2003; Yoon 2006). The M827I + T929I + L932F triple mutant Vssc1 channel is completely insensitive to pyrethroids (Yoon 2006).

It is important to note that some *kdr*-type strains lack the L to F/H/S mutation in IIS6. Instead, a valine (V) to M change at 421 in IS6 is found in several pyrethroid-resistant *H. virescens* populations (Park et al. 1997). The V421M mutation reduces pyrethroid sensitivity by ten-fold when introduced into *Drosophila*, housefly and cockroach sodium channels (Zhao et al. 2000; Lee and Soderlund 2001; Liu et al. 2002). The L to F/H/S mutation is also not found in any pyrethroid-resistant arachnids. Instead, an F to I mutation in IIS6 is present in several pyrethroid-resistant southern cattle tick (*Boophilus microplus*) strains (He et al. 1999). This mutation completely abolishes the pyrethroid sensitivity of cockroach sodium channels to structurally diverse pyrethroids (Tan et al. 2005). Finally, four amino acid changes in the sodium channel have been identified in pyrethroid resistant varroa mites, an ectoparasite of honey bees, including an L to P mutation in the third linker connecting domains III and IV (Wang et al. 2002). Interestingly, pyrethroid-susceptible insects already possess a P at the corresponding position. This is intriguing as varroa mites are more sensitive to the pyrethroid fluvalinate (which is used in the control of mites) than honey bees (Santiago et al. 2000). Indeed, substitution of P with L makes the cockroach sodium channel more sensitive to pyrethroids (Liu et al. 2006). These results indicate that the L/P residues are at least partially responsible for the differential sensitivities of insects and mites to pyrethroids.

In summary, both common and unique mutations in sodium channel genes are found to be responsible for pyrethroid resistance in different insect and arachnid pest species. The information is valuable for monitoring the frequency of pyrethroid resistance in field pest populations as part of pest resistance management. New *kdr* mutations will likely be identified in other agricultural and medically important arthropod pests as pyrethroids continue to be used as a major pest control strategy.

Do *kdr* mutations define the pyrethroid-binding site(s)?

Studies are being conducted to understand how the *kdr* mutations alter sodium channel sensitivity to pyrethroids. These mutations could confer sodium channel resistance to pyrethroids by reducing the binding of pyrethroids to the sodium channel and/or by counteracting the action of pyrethroids via a binding-independent mechanism (e.g., by altering sodium channel gating). Although a high-affinity pyrethroid-binding site was detected on the α -subunit of rat brain sodium channel preparations ten years ago (Trainer et al. 1997), direct measurement of pyrethroid binding affinity and capacity to insect sodium channels has not been achieved, because the high lipophilicity of pyrethroids results in extremely high levels of nonspecific binding to membranes and filters and consequently masks any specific pyrethroid binding (Pauron et al. 1989; Rossignol 1988; Dong 1993). As an alternative strategy, electrophysiological analyses of *kdr* mutant channels in oocytes are employed to quantify the effects of *kdr* mutations on the channel gating and pyrethroid binding. Vais et al. (2000a) showed that *kdr* (L1014F) in IIS6 and *super-kdr* mutations M918T in the IIS4–S5 linker and T929I in IIS5 all enhanced closed-state inactivation, thereby reduced channel opening, which is required for the action of pyrethroids. These mutations also reduced the binding affinity for open channels (Vais et al. 2000a, 2003). Utilizing the competitive binding of active 1R-cis and inactive 1S-cis permethrin isomers to the sodium channel, Tan et al. (2005) showed that L993F in IIS6 and F1519W in IIS6 reduced pyrethroid binding to the cockroach sodium channel. The findings from these studies suggest that these particular residues are part of the pyrethroid binding site. However, it is possible that some of these residues do not physically interact with pyrethroids, but they are critical for the sodium channel conformation required for the

binding of pyrethroids. A recent study used the crystal structures of potassium channels to generate a housefly sodium channel model in the open conformation (O'Reilly et al. 2006). This model highlights the role of IIS4–S5 linker and the IIS5 and IIS6 helices in pyrethroid binding, supporting the involvement of M918, T929 and F1519 in pyrethroid binding. The model also predicts that several additional residues in IIS5 and IIS6 could be part of the pyrethroid binding site and might contribute to the different sensitivities between insect and mammalian sodium channels. Further experiments are needed to verify these predictions.

Conclusion

Since the seminal cloning of the *Drosophila para* gene 18 years ago, excellent progress has been made in understanding the biology of insect sodium channels. Successful functional expression in *Xenopus* oocytes and the identification of TipE as an auxiliary protein have had a major impact on many aspects of research on insect sodium channels. A recent significant development is the demonstration that alternative splicing and RNA editing are two main mechanisms by which insects generate a remarkable array of structurally and functionally diverse sodium channel variants from a single sodium channel gene. Determining how alternative splicing and RNA editing of the sodium channel gene shapes the electrical activity of specific neurons involved in specific sensory, motor, and integrative functions in vivo is an exciting next step for future research. Also, after two decades of speculation, the functional identity of the DSC1 channel has been resolved. The discovery that DSC1 is a novel calcium-selective cation channel now begs for research aimed at determining its precise function in insect development, physiology and behavior.

Without doubt, the great interest in insect sodium channels in the past decade is in part prompted by an exciting connection between pyrethroid resistance and sodium channel mutations. The intensive research efforts in this area have greatly improved our understanding of the molecular basis of pyrethroid resistance. An impressive collection of *kdr* or *kdr*-type sodium channel mutations have been identified and confirmed to be involved in reducing sodium channel sensitivity to pyrethroids. The naturally occurring pyrethroid-resistant sodium channel variants provide a complementary approach for understanding the interaction of pyrethroids with insect sodium channels at the molecular level. Collectively, these insect sodium channel mutations and variants provide an excellent foundation for further studying the interaction of sodium channels with pyrethroids. It seems certain that the next few years will witness continuing progress on the elucidation of the structural features in sodium channels that are required for the binding/action of pyrethroids.

Sodium channels are excellent target sites for a variety of neurotoxins, in addition to pyrethroids. There are at least nine known toxin binding sites on the sodium channel, which could be potential target sites of new insecticides. Indoxacarb/DCJW is an example of a new chemical which successfully exploits a binding site on the sodium channel unique from other classes of insecticides. Additionally, several scorpion toxins are selectively active on insect sodium channels, but not on their mammalian counterparts (Bosmans et al. 2005; Strugastsky et al. 2005). The unique pharmacology displayed by insect sodium channels can serve to provide leads for new insecticide discovery and development. Furthermore, study of the action of novel neurotoxins in turn should enhance our basic understanding of sodium channel gating and pharmacological properties unique to insects.

Acknowledgments

The research in the author's laboratory is supported by NSF grants (IBN 9696092 and IBN 9808156), NIH (GM57440), USDA-NRI (35607-14966) and BARD (IS-3480-03) and the Michigan State University Rackham Endowment Fund. The author thanks past and current lab members for their contributions to the research in the

author's laboratory, thank Dr. Kris Silver and other lab members for critical reading of this article, and Ms. Jung-Eun Lee for assistance with the references.

References

- Bass, BL. RNA editing. Oxford University Press; Oxford: 2001.
- Bass C, Schroeder I, Turberg A, Field LM, Williamson MS. Identification of mutations associated with pyrethroid resistance in the para-type sodium channel of the cat flea, *Ctenocephalides felis*. *Insect Biochem Mol Biol* 2004;34:1305–1313. [PubMed: 15544944]
- Bosmans F, Martin-Eauclaire MF, Tytgat J. The depressant scorpion neurotoxin LqgIT2 selectively modulates the insect voltage-gated sodium channel. *Toxicon* 2005;45:501–507. [PubMed: 15733572]
- Brun-Barale A, Bouvier JC, Pauron D, Berge JB, Sauphanor B. Involvement of a sodium channel mutation in pyrethroid resistance in *Cydia pomonella* L, and development of a diagnostic test. *Pest Manag Sci* 2005;61:549–554. [PubMed: 15657957]
- Byerly L, Leung H. Ionic currents of *Drosophila* neurons in embryonic cultures. *J Neurosci* 1988;8:4379–4393. [PubMed: 2460598]
- Catterall WA. From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron* 2000;26:13–25. [PubMed: 10798388]
- Catterall WA, Goldin AL, Waxman SG. International Union of Pharmacology International Union of Pharmacology. XXXIX Compendium of voltage-gated ion channels: sodium channels. *Pharmacol Rev* 2003;55:575–578. [PubMed: 14657413]
- Cestele S, Catterall WA. Molecular mechanisms of neurotoxin action on voltage-gated sodium channels. *Biochimie* 2000;82:883–892. [PubMed: 11086218]
- Choi JS, Soderlund DM. Structure-activity relationships for the action of 11 pyrethroid insecticides on rat Na_v 1.8 sodium channels expressed in *Xenopus* oocytes. *Toxicol Appl Pharmacol* 2006;211:233–244. [PubMed: 16051293]
- Crill WE. Persistent sodium current in mammalian central neurons. *Annu Rev Physiol* 1996;58:349–362. [PubMed: 8815799]
- Defaix A, Lapied B. Role of a novel maintained low-voltage- activated inward current permeable to sodium and calcium in pacemaking of insect neurosecretory neurons. *Invert Neurosci* 2005;5:135–146. [PubMed: 16177888]
- Derst C, Walther C, Veh RW, Wicher D, Heinemann SH. Four novel sequences in *Drosophila melanogaster* homologous to the auxiliary Para sodium channel. *Biochem Biophys Res Commun* 2006;339:938–948.
- Dib-Hajj SD, Black JA, Cummins TR, Waxman SG. Na_v1.9: a sodium channel with unique properties. *Trends Neurosci* 2002;25:253–259. [PubMed: 11972962]
- Dong, K. PhD thesis. Cornell University; 1993. Molecular mechanism of knockdown (*kdr*)-type resistance to pyrethroid insecticides in the German cockroach (*Blattella germanica* L.).
- Dong K. A single amino acid change in the Para sodium channel protein is associated with knockdown-resistance (*kdr*) to pyrethroid insecticides in German cockroach. *Insect Biochem Mol Biol* 1997;27:93–100. [PubMed: 9066120]
- Dong K, Valles SM, Scharf ME, Zeichner B, Bennett GW. The knockdown resistance (*kdr*) mutation in pyrethroidresistant German cockroaches. *Pestic Biochem Physiol* 1998;60:195–204.
- Du Y, Liu Z, Nomura Y, Khambay B, Dong K. An alanine in segment 3 of domain III (IIIS3) of the cockroach sodium channel contributes to the low pyrethroid sensitivity of an alternative splice variant. *Insect Biochem Mol Biol* 2006;36:161–168. [PubMed: 16431283]
- Elliott, M. Synthetic pyrethroids. In: Elliott, M., editor. Synthetic pyrethroids. American Chem Soc; Washington: 1977. p. 1-28.
- Feng G, Deak P, Chopra M, Hall LM. Cloning and functional analysis of TipE, a novel membrane protein that enhances *Drosophila para* sodium channel function. *Cell* 1995;82:1001–1011. [PubMed: 7553842]

- Forcioli D, Frey D, Frey JE. High nucleotide diversity in the *para*-like voltage-sensitive sodium channel gene sequence in the western flower thrips (Thysanoptera:Thripidae). *J Econ Entomol* 2002;95:838–848. [PubMed: 12216829]
- Gammon DW, Brown MA, Casida JE. Two classes of pyrethroid action in the cockroach. *Pestic Biochem Physiol* 1981;15:181–191.
- Ginsburg KS, Narahashi T. Differential sensitivity of tetrodotoxin-sensitive and tetrodotoxin-resistant sodium channels to the insecticide allethrin in rat dorsal root ganglion neurons. *Brain Res* 1993;627:239–248. [PubMed: 8298967]
- Goldin AL. Diversity of mammalian voltage-gated sodium channels. *Ann NY Acad Sci* 1999;868:38–50. [PubMed: 10414280]
- Goldin AL. Resurgence of sodium channel research. *Annu Rev Physiol* 2001;63:871–894. [PubMed: 11181979]
- Goldin AL. Mechanisms of sodium channel inactivation. *Curr Opin Neurobiol* 2003;13:284–290. [PubMed: 12850212]
- Goldin AL, Barchi RL, Caldwell JH, Hofmann F, Howe JR, Hunter JC, Kellen RG, Mandel G, Meisler MH, Netter YB, Noda M, Tamkun MM, Waxman SG, Wood JN, Catterall WA. Nomenclature of voltage-gated sodium channels. *Neuron* 2000;28:365–368. [PubMed: 11144347]
- Gordon, D. Sodium channels as targets for neurotoxins: Mode of action and interaction of neurotoxins with receptor sites on sodium channels. In: Gutman, Y.; Lazarowici, P., editors. *Toxins and signal transduction*. Harwood Academic Publishers; The Netherlands: 1997. p. 119-149.
- Grolleau F, Lapied B. Dorsal unpaired median neurons in the insect central nervous system: towards a better understanding of the ionic mechanisms underlying spontaneous electrical activity. *J Exp Biol* 2000;203:1633–1648. [PubMed: 10804154]
- Guerrero FD, Jamroz RC, Kammlah D, Kunz SE. Toxicological and molecular characterization of pyrethroid-resistant horn flies, *Haematobia irritans*: identification of *kdr* and super-*kdr* point mutations. *Insect Biochem Mol Biol* 1997;27:745–755. [PubMed: 9443375]
- Hanrahan CJ, Palladino MJ, Ganetzky B, Reenan RA. RNA editing of the *Drosophila para* Na⁺ channel transcript: evolutionary conservation and developmental regulation. *Genetics* 2000;155:1149–1160. [PubMed: 10880477]
- He H, Chen AC, Davey RB, Ivie GW, George JE. Identification of a point mutation in the *para*-type sodium channel gene from a pyrethroid-resistant cattle tick. *Biochem Biophys Res Commun* 1999;261:558–561. [PubMed: 10441465]
- Hille B. Local anesthetics: hydrophilic and hydrophobic pathways for the drug-receptor reaction. *J Gen Physiol* 1977;69:497–515. [PubMed: 300786]
- Ingles PJ, Adams PM, Knipple DC, Soderlund DM. Characterization of voltage-sensitive sodium channel gene coding sequences from insecticides-susceptible and knockdown-resistant house fly strains. *Insect Biochem Mol Biol* 1996;26:319–326. [PubMed: 8814780]
- Kulkarni NH, Yamamoto AH, Robinson KO, Mackay TFC, Anholt RRH. The DSC1 channel, encoded by the *smi60E* locus, contributes to odor-guided behavior in *Drosophila melanogaster*. *Genetics* 2002;161:1507–1516. [PubMed: 12196396]
- Lapied B, Malecot CO, Pelhate M. Patch-clamp study of the properties of the sodium current in cockroach single isolated adult aminergic neurons. *J Exp Biol* 1990;151:387–404.
- Lapied B, Stankiewicz M, Grolleau F, Rochat H, Zlotkin E, Pelhate M. Biophysical properties of scorpion alpha-toxin-sensitive background sodium channel contributing to the pace-maker activity in insect neurosecretory cells (DUM neurons). *Eur J Neurosci* 1999;11:1449–1460. [PubMed: 10103139]
- Lapied B, Grolleau F, Sattelle DB. Indoxacarb, an oxadiazine insecticide, blocks insect neuronal sodium channels. *Br J Pharmacol* 2001;132:587–595. [PubMed: 11159709]
- Lawrence LJ, Casida JE. Pyrethroid toxicology: mouse intra-cerebral structure-toxicity relationships. *Pestic Biochem Physiol* 1982;18:9–14.
- Le Corrionc H, Hue B, Pitman RM. Ionic mechanisms underlying depolarizing response of an identified insect motor neuron to short periods of hypoxia. *J Neurophysiol* 1999;81:307–318. [PubMed: 9914291]

- Lee SH, Soderlund DM. The V410M mutation associated with pyrethroid resistance in *Heliothis virescens* reduces the pyrethroid sensitivity of housefly sodium channels expressed in *Xenopus* oocytes. *Insect Biochem Mol Biol* 2001;31:19–29. [PubMed: 11102831]
- Lee SH, Dunn JB, Clark JM, Soderlund DM. Molecular analysis of *kdr*-like resistance in a permethrin-resistant strain of Colorado potato beetle. *Pestic Biochem Physiol* 1999a;63:63–75.
- Lee SH, Smith TJ, Knipple DC, Soderlund DM. Mutations in the house fly *Vssc1* sodium channel gene associated with super-*kdr* resistance abolish the pyrethroid sensitivity of *Vssc1*/tipE sodium channels expressed in *Xenopus* oocytes. *Insect Biochem Mol Biol* 1999c;29:185–194. [PubMed: 10196741]
- Lee SH, Yoon KS, Williamson MS, Goodson SJ, Takano-Lee M, Edman JD, Devonshire AL, Clark JM. Molecular analysis of *kdr*-like resistance in permethrin-resistant strains of head lice, *Pediculus capitis*. *PesticBiochemPhysiol* 2000;66:130–143.
- Lee SH, Ingles PJ, Knipple DC, Soderlund DM. Developmental regulation of alternative exon usage in the house fly *Vssc1* sodium channel gene. *Invert Neurosci* 2002;4:125–133. [PubMed: 12488972]
- Lee SH, Gao JR, Yoon KS, Mumcuoglu KY, Taplin D, Edman JD, Takano-Lee M, Clark JM. Sodium channel mutations associated with knockdown resistance in the human head louse, *Pediculus capitis* (De Geer). *Pestic Biochem Physiol* 2003;75:79–91.
- Littleton JT, Ganetzky B. Ion channels and synaptic organization: analysis of the *Drosophila* genome. *Neuron* 2000;26:35–43. [PubMed: 10798390]
- Liu Z, Valles SM, Dong K. Novel point mutations in the German cockroach *para* sodium channel gene are associated with knockdown resistance (*kdr*) to pyrethroid insecticides. *Insect Biochem Mol Biol* 2000;30:991–997. [PubMed: 10899465]
- Liu Z, Chung I, Dong K. Alternative splicing of the *BSC1* gene generates tissue-specific isoforms in the German cockroach. *Insect Biochem Mol Biol* 2001;31:703–713. [PubMed: 11267908]
- Liu Z, Tan J, Valles SM, Dong K. Synergistic interaction between two cockroach sodium channel mutations and a tobacco budworm sodium channel mutation in reducing channel sensitivity to a pyrethroid insecticide. *Insect Biochem Mol Biol* 2002;32:397–404. [PubMed: 11886774]
- Liu Z, Song W, Dong K. Persistent tetrodotoxin-sensitive sodium current resulting from U-to-C RNA editing of an insect sodium channel. *Proc Natl Acad Sci USA* 2004;101:11862–11867. [PubMed: 15280550]
- Liu Z, Tan J, Huang ZY, Dong K. Effect of a pyrethroid resistance-associated sodium channel mutation from varroa mites on cockroach sodium channel sensitivity to a pyrethroid insecticide. *Insect Biochem Mol Biol* 2006;36:885–889. [PubMed: 17046602]
- Loughney K, Kreber R, Ganetzky B. Molecular analysis of the *para* locus, a sodium channel gene in *Drosophila*. *Cell* 1989;58:1143–1154. [PubMed: 2550145]
- Martinez-Torres D, Chandre F, Williamson MS, Darriet F, Berge JB, Devonshire AL, Guillet P, Pasteur N. Molecular characterization of pyrethroid knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* s.s. *Insect Mol Bio* 1998;7:179–184. [PubMed: 9535162]
- Martinez-Torres D, Foster SP, Field LM, Devonshire AL, Williamson MS. A sodium channel point mutation is associated with resistance to DDT and pyrethroid insecticides in the peach-potato aphid *Myzus persicae* (Sulzer) (Hemiptera:Aphididae). *Insect Mol Biol* 1999a;8:339–346. [PubMed: 10469251]
- Martinez-Torres D, Chevillon C, Brun-Barale A, Berge JB, Pasteur N, Pauron D. Voltage-dependent Na⁺ channels in pyrethroid resistant *Culex pipiens* L mosquitoes. *Pest Sci* 1999b;55:1012–1020.
- McCann SF, Annis GD, Shapiro RD, Piotrowski W, Lahm GP, Long JK, Lee KC, Hughes MM, Myers BJ, Griswold SM. The discovery of indoxacarb: oxadiazines as a new class of pyrazoline-type insecticides. *Pest Manag Sci* 2001;57:153–164. [PubMed: 11455646]
- Miyazaki M, Ohyama K, Dunlap DY, Matsumura F. Cloning and sequencing of the para-type sodium channel gene from susceptible and *kdr*-resistant German cockroaches (*Blattella germanica*) and house fly (*Musca domestica*). *Mol Gen Genet* 1996;252:61–68. [PubMed: 8804404]
- Nagata K, Ikeda T, Honda H, Shono T. Suppression of voltage-gated sodium currents by the dihydropyrazole insecticide, DPX-JW062 in rat dorsal root ganglion neurons. *J Pest Sci* 1998;23:62–64.

- Narahashi, T. Toxins that modulate the sodium channel gating mechanism. In: Kao, CY.; Levinson, SR., editors. Tetrodotoxin, saxitoxin and the molecular biology of the sodium channel. Annals New York Academy of Sciences; New York: 1986. p. 133-151.
- Narahashi, T. Molecular and cellular approaches to neurotoxicology: past, present and future. In: Lunt, GG., editor. Neurotox '88: molecular basis of drug and pesticide action. Elsevier; New York: 1988. p. 563-582.
- Narahashi T. Neuronal ion channels as the target sites of insecticides. *Pharmacol Toxicol* 1996;78:1–14. [PubMed: 8821391]
- Narahashi T. Neuroreceptors and ion channels as the basis for drug action: past, present, and future. *J Pharmacol Exp Ther* 2000;294:1–26. [PubMed: 10871290]
- Numann R, Catterall WA, Scheuer T. Functional modulation of brain sodium channels by protein kinase C phosphorylation. *Science* 1991;254:115–118. [PubMed: 1656525]
- O'Dowd DK, Gee JR, Smith MA. Sodium current density correlates with expression of specific alternatively spliced sodium channel mRNAs in single neurons. *J Neurosci* 1995;15:4005–4012. [PubMed: 7751961]
- O'Reilly AQ, Khambay BPS, Williamson MS, Field LM, Wallace BA, Davies TGE. Modeling insecticide binding sites at the voltage-gated sodium channel. *Biochem J* 2006;396:255–263. [PubMed: 16475981]
- Palladino MJ, Keegan LP, O'Connell MA, Reenan RA. A-to-I pre-mRNA editing in *Drosophila* is primarily involved in adult nervous system function and integrity. *Cell* 2000;102:437–449. [PubMed: 10966106]
- Park Y, Taylor MF. A novel mutation L1029H in sodium channel hscp associated with pyrethroid resistance for *Heliothis virescens* (Lepidoptera: Noctuidae). *Insect Biochem Mol Biol* 1997;27:9–13. [PubMed: 9061924]
- Park Y, Taylor MF, Feyereisen R. A valine421 to methionine mutation in IS6 of the hscp voltage-gated sodium channel associated with pyrethroid resistance in *Heliothis virescens*. *Biochem Biophys Res Commun* 1997;239:688–691. [PubMed: 9367829]
- Pauron D, Barhanin J, Amichot M, Pralavorio M, Berge J-B, Lazdunski M. Pyrethroid receptor in the insect Na⁺ channel: alteration of its properties in pyrethroid-resistant flies. *Biochem* 1989;28:1673–1677.
- Plummer NW, McBurney MW, Meisler MH. Alternative splicing of the sodium channel *SCN8A* predicts a truncated two-domain protein in fetal brain and non-neuronal cells. *J Biol Chem* 1997;272:24008–24015. [PubMed: 9295353]
- Ranson H, Jensen B, Vulule JM, Wang X, Hemingway J, Collins FH. Identification of a point mutation in the voltagegated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids. *Insect Mol Biol* 2000;9:491–497. [PubMed: 11029667]
- Raymond-Delpech V, Matsuda K, Sattelle BM, Rauh JJ, Sattelle DB. Ion channels:molecular targets of neuroactive insecticides. *Invert Neurosci* 2005;5:119–133. [PubMed: 16172884]
- Reenan RA, Nanrahan CJ, Ganetzky B. The mlenapts RNA helicase mutation in *Drosophila* results in a splicing catastrophe of the *para* Na⁺ channel transcript in a region of RNA editing. *Neuron* 2000;25:139–149. [PubMed: 10707979]
- Rossignol DP. Reduction in number of nerve membrane sodium channels in pyrethroid resistant house flies. *Pestic Biochem Physiol* 1988;32:146–152.
- Saito M, Wu CF. Expression of ion channels and mutational effects in giant *Drosophila* neurons differentiated from cell division-arrested embryonic neuroblasts. *J Neurosci* 1991;11:2135–2150. [PubMed: 1712379]
- Saito, M.; Wu, CF. Ionic channels in cultured *Drosophila* neurons. In: Pichon, Y., editor. Comparative molecular neurobiology. Birkhauser Verlag; Basel: 1993. p. 366-389.
- Salgado VL. Slow voltage-dependent block of sodium channels in crayfish nerve by dihydropyrazole insecticides. *Mol Pharmacol* 1992;41:120–126. [PubMed: 1310138]
- Salkoff L, Butler A, Wei A, Scavarda N, Giffen K, Ifune C, Goodman R, Mandel G. Genomic organization and deduced amino acid sequence of a putative sodium channel gene in *Drosophila*. *Science* 1987;237:744–748. [PubMed: 2441469]

- Santiago GP, Otero-Colina G, Sanchez DM, Guzman MER, Vandame R. Comparing effects of three acaricides on *Varroa jacobsoni* (Acari: Varroidae) and *Apis mellifera* (Hymenoptera: Apidae) using two application techniques. *Florida Entomol* 2000;83:468–476.
- Schafer S, Rosenboom H, Menzel R. Ionic currents of Kenyon cells from the mushroom body of the honeybee. *J Neurosci* 1994;14:4600–4612. [PubMed: 7519255]
- Schuler TH, Martinez-Torres D, Thompson AJ, Denholm I, Devonshire AL, Duce IR, Williamson MS. Toxicological, electrophysiological and molecular characterization of knockdown resistance to pyrethroid insecticides in the diamond-back moth, *Plutella xylostella* (L.). *Pestic Biochem Physiol* 1998;59:169–182.
- Seeburg PH. RNA helicase participates in the editing game. *Neuron* 2000;25:261–263. [PubMed: 10719882]
- Silver K, Soderlund DM. Action of pyrethroid-type insecticides at neuronal target sites. *Pestic Biochem Physiol* 2005a;81:136–143.
- Silver K, Soderlund DM. State-dependent block of rat Nav1.4 sodium channels expressed in *Xenopus* oocytes by pyrazoline-type insecticides. *NeuroToxicol* 2005b;26:397–406.
- Silver K, Soderlund DM. Differential sensitivity of rat voltage-sensitive sodium channel isoforms to pyrazolinetype insecticides. *Toxicol Appl Pharmacol* 2006;214:209–217. [PubMed: 16487558]
- Smith RD, Goldin AL. Phosphorylation of brain sodium channels in the I–II linker modulates channel function in *Xenopus* oocytes. *J Neurosci* 1996;16:1965–1974. [PubMed: 8604040]
- Smith RD, Goldin AL. Phosphorylation at a single site in the rat brain sodium channel is necessary and sufficient for current reduction by protein kinase A. *J Neurosci* 1997;17:6086–6093. [PubMed: 9236220]
- Smith RD, Goldin AL. Potentiation of rat brain sodium channel currents by PKA in *Xenopus* oocytes involves the I–II linker. *Am J Physiol Cell Physiol* 2000;278:C638–C645. [PubMed: 10751312]
- Smith TJ, Soderlund DM. Action of the pyrethroid insecticide cypermethrin on rat brain IIa sodium channels expressed in *Xenopus* oocytes. *Neurotox* 1998;19:823–832.
- Smith TJ, Lee SH, Ingles PJ, Knipple DC, Soderlund DM. The L1014F point mutation in the house fly *Vssc1* sodium channel confers knockdown resistance to pyrethroids. *Insect Biochem Mol Biol* 1997;27:807–812. [PubMed: 9474777]
- Soderlund, D. Sodium channels. In: Gilbert, LI.; Iatrou, K.; Gill, SS., editors. *Comprehensive insect science. Pharmacology*. Vol. 5. Elsevier B.V; Amsterdam: 2005. p. 1-24.
- Soderlund DM, Bloomquist JR. Neurotoxic actions of pyrethroid insecticides. *Annu Rev Entomol* 1989;34:77–96. [PubMed: 2539040]
- Soderlund, DM.; Bloomquist, JR. Molecular mechanisms of insecticide resistance. In: Roush, RT.; Tabashnik, BE., editors. *Pesticide resistance in arthropods*. Chapman and Hall; New York: 1990. p. 58-96.
- Soderlund DM, Knipple DC. The molecular biology of knockdown resistance to pyrethroid insecticides. *Insect Biochem Mol Biol* 2003;33:563–577. [PubMed: 12770575]
- Song JH, Narahashi T. Modulation of sodium channels of rat cerebellar Purkinje neurons by the pyrethroid tetramethrin. *J Pharmacol Exp Ther* 1996;277:445–453. [PubMed: 8613953]
- Song W, Liu Z, Tan J, Nomura Y, Dong K. RNA editing generates tissue-specific sodium channels with distinct gating properties. *J Biol Chem* 2004;279:2554–2561.
- Song W, Liu Z, Dong K. Molecular basis of differential sensitivity of insect sodium channels to DCJW, a bioactive metabolite of the oxadiazine insecticide indoxacarb. *Neurotoxicol* 2006;27:237–244.
- Strugastsky D, Zilberberg N, Stankiewicz M, Ilan N, Turkov M, Cohen L, Pelhate M, Gilles N, Gordon D, Gurevitz M. Genetic polymorphism and expression of a highly potent scorpion depressant toxin enable refinement of the effects on insect Na channels and illuminate the key role of Asn-58. *Biochem* 2005;44:9179–9187. [PubMed: 15966742]
- Tan J, Liu Z, Nomura Y, Goldin AL, Dong K. Alternative splicing of an insect sodium channel gene generates pharmacologically distinct sodium channels. *J Neurosci* 2002a;22:5300–5309. [PubMed: 12097481]

- Tan J, Liu Z, Tsai TD, Valles SM, Goldin AL, Dong K. Novel sodium channel gene mutations in *Blattella germanica* reduce the sensitivity of expressed channels to deltamethrin. *Insect Biochem Mol Biol* 2002b;32:445–454. [PubMed: 11886779]
- Tan J, Liu Z, Wang R, Huang ZY, Chen AC, Gurevitz M, Dong K. Identification of amino acid residues in the insect sodium channel critical for pyrethroid binding. *Mol Pharmacol* 2005;67:513–522. [PubMed: 15525757]
- Tatebayashi H, Narahashi T. Differential mechanism of action of the pyrethroid tetramethrin on tetrodotoxin-sensitive and tetrodotoxin-resistant sodium channels. *J Pharmacol Exp Ther* 1994;270:595–603. [PubMed: 8071852]
- Thackeray JR, Ganetzky B. Developmentally regulated alternative splicing generates a complex array of *Drosophila para* sodium channel isoforms. *J Neurosci* 1994;14:2569–2578. [PubMed: 8182428]
- Thackeray JR, Ganetzky B. Conserved alternative splicing patterns and splicing signals in the *Drosophila* sodium channel gene *para*. *Genetics* 1995;141:203–214. [PubMed: 8536968]
- Tomita T, Yaguchi N, Mihara M, Takahashi M, Agui N, Kasai S. Molecular analysis of a para sodium channel gene from pyrethroid-resistant head lice, *Pediculus humanus capitis* (Anoplura: Pediculidae). *J Med Entomol* 2003;40:468–474. [PubMed: 14680113]
- Trainer VL, McPhee JC, Boutelet-Bochan H, Baker C, Scheuer T, Babin D, Demoute J-P, Guedin D, Catterall WA. High affinity binding of pyrethroids to the α -subunit of brain sodium channel. *Mol Pharmacol* 1997;51:651–657. [PubMed: 9106631]
- Tsurubuchi Y, Kono Y. Modulation of sodium channels by the oxadiazine insecticide indoxacarb and its N-decarbomethoxylated metabolite in rat dorsal root ganglion neurons. *Pest Manag Sci* 2003;59:999–1006. [PubMed: 12974351]
- Vais H, Williamson MS, Hick CA, Eldursi N, Devonshire AL, Usherwood PNR. Functional analysis of a rat sodium channel carrying a mutation for insect knockdown resistance (*kdr*) to pyrethroids. *FEBS Lett* 1997;413:327–332. [PubMed: 9280307]
- Vais H, Williamson MS, Goodson SJ, Devonshire AL, Warmke JW, Usherwood PNR, Cohen C. Activation of *Drosophila* sodium channels promotes modification by deltamethrin: reductions in affinity caused by knock-down resistance mutations. *J Gen Physiol* 2000a;115:305–318. [PubMed: 10694259]
- Vais H, Atkinson S, Eldursi N, Devonshire AL, Williamson MS, Usherwood PN. A single amino acid change makes a rat neuronal sodium channel highly sensitive to pyrethroid insecticides. *FEBS Lett* 2000b;470:135–138. [PubMed: 10734222]
- Vais H, Williamson MS, Devonshire AL, Usherwood PN. The molecular interactions of pyrethroid insecticides with insect and mammalian sodium channels. *Pest Manag Sci* 2001;57:877–888. [PubMed: 11695180]
- Vais H, Atkinson S, Plutau F, Goodson SJ, Devonshire AL, Williamson MS, Usherwood PN. Mutations of the para sodium channel of *Drosophila melanogaster* identify putative binding sites for pyrethroids. *Mol Pharmacol* 2003;64:914–922. [PubMed: 14500748]
- Wang SY, Wang GK. Voltage-gated sodium channels as primary targets of diverse lipid-soluble neurotoxins. *Cell Signal* 2003;15:151–159. [PubMed: 12464386]
- Wang SY, Barile M, Wang GK. A phenylalanine residue at segment D3–S6 in Nav1.4 voltage-gated Na⁺ channels is critical for pyrethroid action. *Mol Pharmacol* 2001;60:620–628. [PubMed: 11502895]
- Wang R, Liu Z, Dong K, Elzen P, Pettis J, Huang ZY. Novel mutations in a sodium channel gene are associated with fluvalinate resistance in the varroa mite, *Varroa destructor*. *J Apic Res* 2002;40:17–25.
- Wang R, Huang ZY, Dong K. Molecular characterization of an arachnid sodium channel gene from the varroa mite (*Varroa destructor*). *Insect Biochem Mol Biol* 2003;33:733–739. [PubMed: 12826100]
- Warmke JW, Reenan RAG, Wang P, Qian S, Arena JP, Wang J, Wunderler D, Liu K, Kaczorowski GJ, Van der Ploeg LHT, Ganetzky B, Cohen CJ. Functional expression of *Drosophila para* sodium channels: Modulation by the membrane protein tipE and toxin Pharmacology. *J Gen Physiol* 1997;110:119–133. [PubMed: 9236205]

- West JW, Numann R, Murphy BJ, Scheuer T, Catterall WA. A phosphorylation site in the Na⁺ channel required for modulation by protein kinase C. *Science* 1991;254:866–868. [PubMed: 1658937]
- Wicher D, Walther C, Wicher C. Non-synaptic ion channels in insects—basic properties of currents and their modulation in neurons and skeletal muscles. *Prog Neurobiol* 2001;64:431–525. [PubMed: 11301158]
- Williamson MS, Martinez-Torres D, Hick CA, Devonshire AL. Identification of mutations in the housefly para-type sodium channel gene associated with knockdown resistance (*kdr*) to pyrethroid insecticides. *Mol Gen Genet* 1996;252:51–60. [PubMed: 8804403]
- Wing KD, Schnee ME, Sacher M, Connair M. A novel oxadiazine insecticide is bioactivated in lepidopteran larvae. *Arch Insect Biochem Physiol* 1998;37:91–103.
- Wing, KD.; Andaloro, JT.; McCann, SF.; Salgado, VL. Indoxacarb and the sodium channel blocker insecticides: chemistry, physiology and biology in insects. In: Gilbert, LI.; Iatrou, K.; Gill, SS., editors. *Comprehensive insect science. Insect control*. Vol. 6. Elsevier B.V; Amsterdam: 2005. p. 30-53.
- Xu Q, Liu H, Zhang L, Liu N. Resistance in the mosquito, *Culex quinquefasciatus*, and possible mechanisms for resistance. *Pest Manag Sci* 2005;61:1096–1102. [PubMed: 16032654]
- Yoon, KS. PhD Thesis. University of Massachusetts; 2006. Detection and mechanism of pediculicide resistance in the human head louse, *Pediculus capitis*; p. 1-137.
- Yu FH, Catterall WA. Overview of the voltage-gated sodium channel family. *Genome Biol* 2003;4:207.1–207.7. [PubMed: 12620097]
- Zhao Y, Park Y, Adams ME. Functional and evolutionary consequences of pyrethroid resistance mutations in S6 transmembrane segments of a voltage-gated sodium channel. *Biochem Biophys Res Commun* 2000;278:516–521. [PubMed: 11095943]
- Zhao X, Ikeda T, Yeh JZ, Narahashi T. Voltagedependent block of sodium channels in mammalian neurons by the oxadiazine insecticide indoxacarb and its metabolite DCJW. *Neurotoxicol* 2003;24:83–96.
- Zhou W, Chung I, Liu Z, Goldin A, Dong K. A voltagegated calcium-selected channel encoded by a sodium channel- like gene. *Neuron* 2004;42:101–112. [PubMed: 15066268]
- Zhao X, Ikeda T, Salgado VL, Yeh JZ, Narahashi T. Block of two types of sodium channels in cockroach neurons by indoxacarb insecticides. *Neurotoxicol* 2005;26:455–465.
- Zlotkin E. The insect voltage-gated sodium channel as target of insecticides. *Annu Rev Entomol* 1999;44:429–455. [PubMed: 9990721]

Fig. 1.

The schematic diagram of the sodium channel protein indicating four homologous domains (I–IV), each having six transmembrane segments (1–6). The location of segments encoded by the alternative exons are indicated with solid blocks. Optional exons a, b, e, f, h, i, and j, and mutually exclusive exons c/d and l/k identified in *D. melanogaster para* are present in the housefly sodium channel gene *Vssc1* (Lee et al. 2002). Most of the exons are also found in *BgNa_v*. (Capital letters are used to denote alternative exons in *BgNa_v*) (Song et al. 2004). However, mutually exclusive exons c/d and optional exon h are not identified in *BgNa_v*. Corresponding to the l/k site, three mutually exclusive exons G1/G2/G3 are found in *BgNa_v* (Tan et al. 2002b). Exon K encoding IS2–S3 was found in *BgNa_v*, but is lacking in *para* and *Vssc1*. See the text for details 18 *Invert Neurosci* (2007) 7:17–30

Fig. 2.

RNA editing sites in Para and BgNa_v. Ten A-to-I editing sites have been identified in para, eight of which result in amino acid changes (*solid dots*). The amino acid positions of these editing sites are not numbered in the original publications (Palladino et al. 2000; Hanrahan et al. 2000). Here, six sites are numbered according to the amino acid sequence of the GenBank sequence (accession number: M32078). *We cannot give amino acid positions for R/K and N/S sites because there are more than one R or N residues in these regions. Therefore, only the approximate locations are indicated. Two A-to-I editing sites (K/R¹⁴⁸ and I/M¹⁶⁶⁰) and three U-to-C editing sites (L/P¹²⁸⁵, V/A¹⁶⁸⁵, and F/S¹⁹¹⁹) are identified in BgNa_v. F/S¹⁹¹⁹ is also identified in para (Liu et al. 2004) See the text for details

Fig. 3.

kdr mutations in insect sodium channels. Only those *kdr* mutations that have been confirmed to reduce the sodium channel sensitivity to pyrethroids are indicated (*solid dots*). Positions of M827I, M918T, T929I, and L932F mutations are numbered according to the amino acid sequence of the housefly Vssc1 sodium channel protein (Ingles et al. 1996; Williamson et al. 1996). Positions of E434K, C764K, and F1519I mutation are numbered according to the amino acid sequence of the cockroach sodium channel protein (Dong 1997). V421M is numbered according to the amino acid sequence of the tobacco budworm sodium channel protein (Park et al. 1997). L1770P is numbered according to the amino acid sequence of the varroa mite sodium channel protein (Wang et al. 2003). See the text for details