Published in final edited form as:

J Am Chem Soc. 2011 March 16; 133(10): 3308–3311. doi:10.1021/ja1113936.

Assessment of the Intermediacy of Arylpalladium Carboxylate Complexes in the Direct Arylation of Benzene: Evidence for C-H Bond Cleavage by "Ligandless" Species

Yichen Tan and John F. Hartwig*

Department of Chemistry, University of Illinois at Urbana-Champaign, 600 South Mathews Avenue, Urbana, 61801

Abstract

Palladium-catalyzed direct arylations of benzene have been proposed to occur by the generation of a phosphine-ligated arylpalladium pivalate complex $\text{LPd}(\text{Ar})(\text{OPiv})$ and reaction of this complex with benzene. We have isolated an example of the proposed intermediate and evaluated whether this complex reacts with benzene to form the biaryl products of the catalytic process. In contrast to the proposed mechanism, no biaryl product was formed from cleavage of the benzene C-H bond by $\text{LPd}(\text{Ar})(\text{OPiv})$. However, reactions of $\text{LPd}(\text{Ar})\text{OPiv}$ with benzene and additives that displace or consume the phosphine ligand formed the arylated products in good yield, suggesting that a "ligandless" arylpalladium(II) carboxylate complexes undergoes the C-H cleavage step. Consistent with this conclusion, we found that reactions catalyzed by $\text{Pd}(\text{OAc})_2$ without ligand occur faster than, and with comparable selectivities to, reactions catalyzed by $\text{Pd}(\text{OAc})_2$ and phosphine ligand.

Direct arylation, the reaction of aryl halides with arenes or heteroarenes to form biaryl or aryl-heteroaryl products, is an attractive alternative to traditional cross-coupling because it occurs without the need to prepare organometallic or main-group reagents.¹ Early efforts to develop this process focused on the direct arylation of heteroarenes,² arenes with directing groups³ and electron-deficient arenes.⁴ In 2006, the direct arylation of benzene was reported by Lafrance and Fagnou.⁵ Substoichiometric amounts of pivalic acid were used in the reported catalyst system, and this carboxylic acid was proposed to function as a proton shuttle during the aryl C-H cleavage step.⁶

Despite the improvements in the reaction scope, little experimental information on the mechanism of this reaction is available, especially on the mechanism of the cleavage of the C-H bonds in arenes. Phosphine-ligated arylpalladium carboxylates $\text{LPd}(\text{Ar})(\text{OCOR})$ are typically proposed to react with heteroarenes or arenes to form biaryl complexes through a concerted metallation-deprotonation (CMD) pathway.⁷ DFT calculations of this pathway have been conducted, and these studies suggested that arylpalladium acetates are competent to undergo C-H cleavage with various heteroarenes.⁸ Indeed, a recent report from Fagnou's group showed that the reaction of $\text{Pd}(\text{P}t\text{Bu}_3)(\text{Ph})(\text{OAc})$ with 4-nitropyridine *N*-oxide formed the arylated product.⁹ However, arylpalladium pivalate complexes have not been studied either experimentally or theoretically, and the reactions of isolated arylpalladium carboxylate complexes with arenes have not been conducted.

jhartwig@illinois.edu.

Supporting Information Available: Experimental procedures, spectra for all new compounds, and crystallographic data for **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>

We report the synthesis and characterization of the phosphine-ligated arylpalladium pivalate complexes $\text{Pd}(\text{P}t\text{Bu}_3)(\text{Ar})(\text{OPiv})$. Inconsistent with previous proposals, the reaction of isolated $\text{Pd}(\text{P}t\text{Bu}_3)(\text{Ar})(\text{OPiv})$ with benzene does not form Ar-Ph products. However, reactions conducted with additives that displace or consume the phosphine ligand promote the formation of the arylated products, and these results and others reported herein imply the involvement of “ligandless” arylpalladium(II) species in the C-H cleavage step of direct arylation of benzene.

The synthesis of $\text{P}t\text{Bu}_3\text{Pd}(\text{Ar})(\text{OPiv})$ is outlined in Scheme 1. $\text{P}t\text{Bu}_3$ -ligated arylpalladium iodide **1** was prepared by the reaction of $\text{Pd}(\text{dba})_2$ with $\text{P}t\text{Bu}_3$ in neat 2-iodotoluene. Reaction of **1** with AgOPiv formed the $\text{P}t\text{Bu}_3$ -ligated arylpalladium pivalate **2** in good yield. Complex **2** was characterized by elemental analysis and NMR spectroscopy. The solid-state structure of complex **2** was determined by single-crystal X-ray diffraction (Figure 1). The pivalate ligand binds to the Pd center in a η^2 -mode with Pd-O bond distances of 2.124(4) Å and 2.301(4) Å. The longer Pd-O bond is trans to the aryl group.

The competency of complex **2** to be an intermediate in the catalytic process was then assessed, and the results of these studies are summarized in Table 1 (entry 1–3) and eq 1. To provide a benchmark to compare the reactivity of **2**, we conducted the reaction of 2-bromotoluene with benzene catalyzed by $\text{Pd}(\text{OAc})_2$ and $\text{P}t\text{Bu}_3$. This reaction formed 2-phenyl-toluene in 71% yield (eq 1). In contrast, the stoichiometric reaction of complex **2** with benzene in the presence of K_2CO_3 in DMA at 110 °C formed only trace amounts of 2-phenyl-toluene. The major organic products were toluene and 2,2'-dimethyl-biphenyl. The major organometallic product (55% NMR yield) was palladacycle **6** from intramolecular C-H bond cleavage of the phosphine ligand. Attempts to suppress the formation of 2,2'-dimethyl-biphenyl by conducting the reaction with lower concentration of **2** that would be present in the catalytic system led to similar results. Reaction of **2** in benzene in the presence of PivOH also failed to form 2-phenyl-toluene. Thus, the $\text{P}t\text{Bu}_3$ -ligated arylpalladium pivalate complex **2** is not competent to be the intermediate that cleaves the arene C-H bond in the catalytic direct arylation process.

(1)

The absence of the product of the catalytic reaction in the stoichiometric reactions of arylpalladium pivalate complex **2** led us to study the effects of additives that could be present in the catalytic system on the distribution of products. The catalytic system forms halide and pivalate anion, and it could contain adventitious oxygen. The results of reactions containing such additives are summarized in Table 1 (entry 4–7). Reactions of **2** with benzene in the presence of a soluble NR_4Br salt, the soluble NR_4OPiv salt, or O_2 formed 2-phenyl-toluene (**3**) in higher yields than in the absence of these additives. The reaction with 0.10 M added NHep_4Br formed **3**, but in just 30% yield with a ratio of organic products **3/5** of only 1.3:1. However, the same reaction at a concentration of complex **2** (0.003 M) that would be closer to that in the catalytic system formed **3** in 51% yield. Moreover, the ratio of **3/5** was 7.3, and this value matched that of the catalytic system. Modest yields of **3** (49% and 63%, respectively) and good ratios of **3/5** were also obtained from reaction with added NHep_4OPiv or O_2 . We propose that displacement of the phosphine ligand by the halide¹⁰ or pivalate anion, or oxidation of the phosphine ligand by the oxygen generates a “ligandless” arylpalladium species that undergoes the C-H cleavage step.

To test our hypothesis that an intermediate lacking a phosphine ligand undergoes the C-H cleavage step, reaction of benzene with $[\text{Pd}(\text{o-Tol})\text{Br}_2]_2^{2-}$ (**7**) lacking a phosphine ligand was conducted (eq 2) with base and pivalic acid. The reaction of **7** with benzene in the presence of PivOH formed 57% yield of biaryl **3**, and the ratio of **3/5** was again similar to that of the catalytic process (7.0).¹¹ These data support our hypothesis that the C-H cleavage step occurs through an arylpalladium species lacking phosphine ligand.

(2)

If our proposal that the C-H bond cleavage step occurs by an intermediate lacking a phosphine ligand is valid, and the C-H bond cleavage step is turnover-limiting as proposed previously,^{5,9} the reactions of bromoarenes with benzene should form the arylated product with faster rates when catalyzed by palladium in the absence of phosphine. Indeed, the reactions of 4-bromotoluene with benzene catalyzed by Pd(OAc)₂ alone occurred faster than, and with comparable selectivities to, reactions catalyzed by Pd(OAc)₂ and a phosphine ligand. Under published conditions with added Davephos,⁵ 76% yield of 4-phenyl-toluene was obtained, and the reaction reached full conversion after 10 h. In comparison, the same reaction catalyzed by Pd(OAc)₂ alone formed 4-phenyl-toluene in 80% yield after only 4 h. These results further support the proposal that C-H bond cleavage of arenes occurs by arylpalladium species lacking a phosphine ligand.¹²

(3)

To assess further whether the C-H cleavage step occurs through such “ligandless” arylpalladium species, we studied the product distribution from the reaction of bromobenzene with two distinct arenes catalyzed by Pd(OAc)₂ with and without phosphine ligand. The results are provided in eq 4 and 5. The reaction of 4-bromotoluene with benzene and benzene-*d*₆ catalyzed by Pd(OAc)₂ and Davephos formed a 5.4(1):1 ratio of 4-MeC₆H₄-C₆H₅ to 4-MeC₆H₄-C₆D₅. The same reaction without added ligand formed an indistinguishable 5.6(2):1 ratio of these products (eq 4). Likewise, the selectivities from reactions of 4-bromotoluene with benzene and fluorobenzene with and without ligand were an indistinguishable 14(2):1 and 13(1):1 (eq 5), and the ratio of ortho, meta, and para isomers of the fluorobiphenyl products were indistinguishable (23:5:1 vs 24:4:1).

(4)

(5)

Finally, to assess the differences in barriers for cleavage of the C-H bond in benzene by arylpalladium species ligated by PtBu_3 and lacking a phosphine ligand in DMA solvent, we used density functional theory (DFT) to obtain computed barriers for the reaction of benzene with $\text{PtBu}_3\text{Pd}(\text{Ph})(\text{OPiv})$ and $(\text{DMA})\text{Pd}(\text{Ph})(\text{OPiv})$. The calculated free energy of activation ($\Delta G^\ddagger$) for the reaction of $(\text{DMA})\text{Pd}(\text{Ph})(\text{OPiv})$ with benzene was found to be 31 kcal/mol at 25 °C. This barrier is much lower than the 42 kcal/mol barrier computed for the reaction of $\text{PtBu}_3\text{Pd}(\text{Ph})(\text{OPiv})$ with benzene (see the Supporting Information for details). The relative magnitude of these barriers is consistent with our proposal that the arylpalladium species lacking phosphine ligand in DMA is more reactive towards C-H bond cleavage than is the phosphine-ligated **2**.

In summary, studies of the reactions of the isolated arylpalladium pivalate complex **2** with benzene have shown that this complex is not chemically competent to be an intermediate in the direct arylation of benzene. However, reactions of **2** conducted in the presence of additives that displace or consume the phosphine ligand in **2** did form the arylated products, suggesting that a “ligandless” arylpalladium species is involved in the C-H cleavage step. This conclusion is consistent with comparisons of selectivities of catalytic reactions conducted with and without added phosphine and with DFT calculations of the barrier of the reactions of benzene with phosphine-ligated and solvent-ligated arylpalladium pivalate complexes.

We propose that the higher reactivity of the ligandless species results in large part from differences in the steric properties of **2** and the species formed by replacement of PtBu_3 with DMA or halide. All isolated arylpalladium complexes ligated by PtBu_3 contain only three ligands.¹³ Thus, the higher coordinate species along the pathway to C-H bond cleavage would be expected to lie at high energy. These studies also suggest that anionic arylpalladium species undergo cleavage of the arene C-H bond with barriers that are lower than those for complexes containing hindered phosphines. This observation suggests that catalysts for direct arylation of unactivated arenes could be developed from systems containing small anionic ligands or additives. Future studies will focus on designing new methods for direct arylation of unactivated arenes based on the principles revealed by these mechanistic studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the NIH (GM-58108) for financial support and Dale Pahls for helpful guidance and suggestions on the DFT calculation. We thank Keith Fagnou for discussions at the inception of these studies. We dedicate this paper to him.

References

1. For recent reviews on direct arylations, see: (a) Alberico D, Scott ME, Lautens M. *Chem. Rev* 2007;107:174. [PubMed: 17212475] (b) Campeau L-C, Stuart DR, Fagnou K. *Aldrichimica Acta* 2007;40:35. (c) Ackermann L, Vicente R, Kapdi AR. *Angew. Chem., Int. Ed* 2009;48:9792.
2. (a) Akita Y, Itagaki Y, Takizawa S, Ohta A. *Chem. Pharm. Bull* 1989;37:1477. (b) Pivsa-Art S, Satoh T, Kawamura Y, Miura M, Nomura M. *Bull. Chem. Soc. Jpn* 1998;71:467. (c) McClure MS, Glover B, McSorley E, Millar A, Osterhout MH, Roschangar F. *Org. Lett* 2001;3:1677. [PubMed: 11405684] (d) Campeau L-C, Rousseaux S, Fagnou K. *J. Am. Chem. Soc* 2005;127:18020. [PubMed: 16366550] (e) Kobayashi K, Sugie A, Takahashi M, Masui K, Mori A. *Org. Lett* 2005;7:5083. [PubMed: 16235963] (f) Lane BS, Brown MA, Sames D. *J. Am. Chem. Soc* 2005;127:8050. [PubMed: 15926829]
3. (a) Satoh T, Kawamura Y, Miura M, Nomura M. *Angew. Chem., Int. Ed. Engl* 1997;36:1740. (b) Satoh T, Kametani Y, Terao Y, Miura M, Nomura M. *Tetrahedron Lett* 1999;40:5345. (c) Daugulis O, Zaitsev VG. *Angew. Chem., Int. Ed* 2005;44:4046. (d) Lazareva A, Daugulis O. *Org. Lett* 2006;8:5211. [PubMed: 17078680]
4. (a) Lafrance M, Rowley CN, Woo TK, Fagnou K. *J. Am. Chem. Soc* 2006;128:8754. [PubMed: 16819868] (b) Lafrance M, Shore D, Fagnou K. *Org. Lett* 2006;8:5097. [PubMed: 17048852]
5. Lafrance M, Fagnou K. *J. Am. Chem. Soc* 2006;128:16496. [PubMed: 17177387]
6. For recent reports on direct arylation of benzene under metal-free conditions, see: (a) Liu W, Cao H, Zhang H, Zhang H, Chung KH, He C, Wang H, Kwong FY, Lei A. *J. Am. Chem. Soc* 2010;132:16737. [PubMed: 20677824] (b) Shirakawa E, Itoh K-I, Higashino T, Hayashi T. *J. Am. Chem. Soc* 2010;132:15537. [PubMed: 20961045] (c) Sun C-L, Li H, Yu D-G, Yu M, Zhou X, Lu X-Y, Huang K, Zheng S-F, Li B-J, Shi Z-J. *Nat. Chem* 2010;2:1044. [PubMed: 21107368]
7. Davies DL, Donald SMA, Macgregor SA. *J. Am. Chem. Soc* 2005;127:13754. [PubMed: 16201772]
8. Gorelsky SI, Lapointe D, Fagnou K. *J. Am. Chem. Soc* 2008;130:10848. [PubMed: 18661978]
9. Sun H-Y, Gorelsky SI, Stuart DR, Campeau L-C, Fagnou K. *J. Org. Chem* 75:8180. [PubMed: 21053903]
10. For recent report on displacement of phosphine ligand by halide, see: Carrow BP, Hartwig JF. *J. Am. Chem. Soc* 2010;132:79. [PubMed: 20014842]
11. The remainder of the mass balance is toluene. At this point, we have not been able to identify the source of the proton. Reactions in DMA-*d*₉ or with added D₂O formed unlabeled toluene, and reactions with tetraheptylammonium pivalate form more toluene than those with pivalic acid and carbonate.
12. For recent reports on direct arylation under ligand-free conditions, see: (a) Glover B, Harvey KA, Liu B, Sharp MJ, Tymoschenko MF. *Org. Lett* 2003;5:301. [PubMed: 12556177] (b) Fournier D, Chabert J, Joucla L, David E, Lemaire M. *Tetrahedron* 2004;60:3221. (c) Parisien M, Valette D, Fagnou K. *J. Org. Chem* 2005;70:7578. [PubMed: 16149786] (d) Wang X, Gribkov DV, Sames D. *J. Org. Chem* 2007;72:1476. [PubMed: 17288392]
13. For recent reports on isolated arylpalladium complexes ligated by PrBu₃, see: (a) Stambuli JP, Incarvito CD, Buehl M, Hartwig JF. *J. Am. Chem. Soc* 2004;126:1184. [PubMed: 14746489] (b) Yamashita M, Hartwig JF. *J. Am. Chem. Soc* 2004;126:5344. [PubMed: 15113190] (c) Sergeev AG, Spannenberg A, Beller M. *J. Am. Chem. Soc* 2008;130:15549. [PubMed: 18956867] (d) Denmark SE, Smith RC. *J. Am. Chem. Soc* 2010;132:1243. [PubMed: 20058920]

Figure 1. ORTEP drawing of $\text{PtBu}_3\text{Pd}(2\text{-MeC}_6\text{H}_4)(\text{OPiv})$ with 50% ellipsoids. Hydrogen atoms have been omitted for clarity.

Scheme 1.
Synthesis of $\text{PtBu}_3\text{Pd}(\text{Ar})(\text{OPiv})$

Table 1

Reactivity of complex **2** with benzene^a

entry	conc. of 2	additives	yield of 3	yield of 4	yield of 5	3/5
1	0.04 M	-	trace	27%	32%	-
2	0.003 M	-	trace	33%	20%	-
3	0.003 M	PivOH (0.03 M)	trace	45%	25%	-
4	0.04 M	NHep ₄ Br (0.1 M)	30%	17%	23%	1.3
5	0.003 M	NHep ₄ Br (0.1 M)	51%	31%	7%	7.3
6	0.003 M	NHep ₄ OPiv (0.1 M)	49%	25%	5%	9.8
7	0.003 M	O ₂ (0.01 M)	63%	26%	<5%	>12

^aYield by GC/MS with an internal standard.