Acardiac Twins

— An Analysis of 10 cases —

Je G. Chi, M.D.

*Department of Pathology, Seoul National University Children's Hospital
Seoul National University College of Medicine*

The pathological characteristics of the acardiac fetus were studied based on 10 autopsy cases. These cases were collected during a 13-year period at Seoul National University Hospital.

All 10 fetuses were monochorionic twins, and six of them were male. Externally normal co-twins survived in five cases and died perinatally in three cases. The gestational period of these acardiacs ranged from 20 to 33 weeks. All of them showed a growth arrest of a fairly wide spectrum. Four cases belonged to acardius anceps, five were acardius acephalus, and one was acardius amorphus. Nine out of 10 cases were holoacardius, whereas one was hemiacardius with a vestigial heart tube present.

Characteristic artery-to-artery anastomoses were demonstrated in all cases where examination was possible. The umbilical cords of the acardiacs often consisted of only two vessels, i.e., one umbilical artery and one umbilical vein, and these vessels were directly attached to the arteries and veins of the normal co-twins on the placental surface. The vascular system of the acardiac fetuses was simplified, providing only inflow and outflow pathways through common iliac vessels and vitelline vessels.

When the head part was preserved (acardius anceps), the facial features were indistinguishable, particularly in the midfacial region, which resulted in a characteristic holoprosencephalic malformation of the brain in two out of four cases. Normal eyeball structure was not noted in any of these cases. The oral cavity and tongue were rarely recognized. Once the head part was absent (acardius acephalus) there was a wide variation of thoracic organ development. Hypoplastic lungs were seen in three cases, and they were connected to the trachea. A tracheo-esophageal fistula was seen twice.

The gastrointestinal tract was the most common feature of these acardiac monsters. However, it often lacked some portions, such as a stomach, midgut or part of a large intestine. The kidneys, testes and other parts of the urinary system were other common constituents of the acardiacs.

Key Words: Holoacardius, Acardiac fetus, Twin, Parasitic twin.

INTRODUCTION

The acardiac monster has often aroused the curiosity of embryologists, pathologists and obstetricians. Since this grotesque malformation, character-

Address for Correspondence: Je G. Chi M.D. Department of Pathology, College of Medicine, Seoul National University, 28 Yeonkun-dong, Chongno-gu, Seoul 110-744, Korea

ized by the absence of a functioning heart, occurs only in pair of twin pregnancies, vascular placental anastomosis between the acardius and its twin litter was particularly emphasized in order to understand its pathogenesis (Napolitani and Schreiber, 1960; Wilson, 1972; Benirschke and Harper, 1977).

The incidence of acardius is estimated to be one in 35,600 deliveries (Gillim and Hendricks, 1953). This statistics is based on their occurrence in about 1 percent of monozygotic twins, which, according to Guttmacher (1956), account for 25 percent of all twin births. In Korea at least five cases of acardius, including that of Kim et al (1971) and Kim et al., (1984) were reported in the literature.

Because this anomaly is so rare and the etiopathogenesis has not been fully elucidated, every single case of acardiac fetus should be thoroughly studied, including cytogenetic examination. Therefore it is necessary that pathologists and obstetricians acquire a reasonable knowledge on how to handle the specimens properly, particularly when examining their cardiovascular systems in regard to feto-feto-placental circulation.

As we examined these anomalies we were impressed by the variety of the associated malformations and also by several consistent findings among them. We found it necessary to classify the holocardius into various groups to give a more clear idea on its pathogenesis.

MATERIALS AND METHODS

During a period of 13 years, from January 1977 to December 1989, we examined 10 cases of acardiac fetuses at the Departments of Pathology of Seoul National University Hospital and Seoul National University Children's Hospital.

These 10 cases included five that were sent from other hospitals or private obstetric clinics in Seoul. In most cases the placentas were sent together with the acardiac fetuses. Some placentas, however, were sent separately without proper orientation to the fetus. Two acardius anceps and one acardius amorphus did not include any placental tissue.

The specimens were photographed and radiographs were taken. Vascular anastomosis was examined carefully. If possible, radio-opaque dye was injected to confirm any anastomosis. After dissection, the tissue was prepared for histological examination by routine procedure.

RESULTS

Overall summaries of the 10 acardiaks on their clinical features and anatomical findings are shown in Table 1. The findings of the individual case are as follows.

Case 1 (RCM #40)

The specimen consisted of a mature placenta and an acardiac twin (Fig. 1). A radiograph of the specimen (Fig. 2) showed well-formed femurs and tibias and fibulae and phalangeal bones. The pelvic bones were partly formed, particularly in the ischium and pubis. The sacrum was incompletely formed. The overlying skin was slightly edematous, and relatively abundant subcutaneous fat was included. The knee joint areas were swollen, and the lower limbs were curved inwardly. The right and left limbs measured 15cm in length, and the maximum thickness was 7cm. The digits were irregularly covered with

Fig. 1. Anterior view of acardiac fetus consisting of lower extremities and pelvic structure. The fetus looks edematous and the limbs are curved inwardly. Instead of genital protuberance, a small nubbin of redundant skin is seen in lower midline region. Case 1.

Table. 1. Clinicopathological summary of the cases

Case no.	Type	Height/weight (Gest/Age)	Sex	Umbilical cord	Placenta	Head	Thorax	Abdomen	Extremities	Other Colwin	Remark
1. RCM #40	Acardius acephalus	15cm (?)	?	1A 1V	Single (Central & marginal cords)	Absent	Absent	Small & large intestine	Lower ext.	Grossly normal	Anasarca
2. RCM #260	Acardius acephalus	13cm (?)	M	1A 1V Furcate cord	Single (Central & marginal cords)	Absent	Attenuated rib	Kidney & adrenals Lower bowel loop Testis Urinary bladder	Lower ext.	Dead several days after birth	Anasarca
3. A82-23	Acardius acephalus	21cm/120gm (?)	M	1A 1V	Single	Absent	Ribs	Intestine, testis Kidneys & bladder Adrenals	Upper & lower ext	Still born Grossly normal	Anasarca
4. A83-24	Acardius acephalus	1620gm (?)	M	1A 1V	Membranous insertion Cotyleton defect	Absent	Absent	Small & large intestine Kidneys Testis	Pelvis Lower ext.	Grossly normal	Anasarca Omphalocele
5. CHA89-57	Acardius acephalus	38cm/2190gm (33 wks)	F	1A 1V	Not examined	Rudimentary skull No brain	Ribs	GI tract & adrenal Kidney Ovary	Lower Ext.	Grossly normal	Anasarca Omphalocele
6. A82-109	Acardius aniceps	23cm/2175gm (33 wks)	M	1A 1V	Not examined	Holotelen- cephal	Enterogenous cyst Fused lung	Kidney with hamartoma Intestine	Oligosyndactyly	Still born Grossly normal	Cystic hygroma Anasarca Omphalocele
7. A86-32	Acardius aniceps	11.3cm/45.2gm (20 wks)	F	(?)	Not examined	Brain tissue cleft palate	Fused lung Trachea	Fused adrenal Intestine Liver Left kidney Asplenia Ovary	Arthrogryposis Oligosyndactyly	Unknown	Anasarca Cystic hygroma
8. A86-23	Acardius aniceps	20cm (?)	M	1A 1V	Single (marginal cords) velamentous	Microphth- almia No brain	Lungs Thymus	GI tract Adrenals Kidneys Testis	Rt ext. agenesis	Grossly normal	Anasarca
9. CHA87-54	Hemicardius aniceps	33cm (?)	M	1A 1V	Single	Holotelen- cephal Lt man- dible	Rudimentary heart Fused lung Esophagus	Midline liver, Tailipes calcaneovarus Kidneys, adrenal GI tract	Lower ext. Oligodactyly	Grossly normal	Anasarca
10. A82-75	Acardius amorphus	10cm (?)	?	Rudimentary	Not examined	Absent	Absent	Fat & cartilage	Absent	Unknown	Anasarca

Fig. 2. Specimen radiograph shows well-formed femurs, tibias and digital bone. Note also partly formed pelvic bones and sacrum, in which bowel loops were found. Case 1.

redundant skin, but their numbers were normal although the lengths were irregular. The pelvic rim area was the actual cut-off point of this twin. There was no other tissue above this level. Anteriorly there was a round defect (1.5×1.5cm) through which two umbilical vessels were inserted into the chorionic membrane of the placenta. Posteriorly the dysplastic sacral bone was exposed together with a skin defect. Numerous black hairs were seen around the defect.

Ventrally along the midline there was no genitalia-like structure. There was only a small nubbin of redundant skin which had no opening. No anal opening was seen posteriorly. Dissection of the specimen revealed a 20cm long small intestine, an appendix and a 7.5cm long large intestine (Fig. 3). Both small and large bowel loops ended blindly. The ileo-cecal region was prominent. Their diameters varied and was filled with whitish meconium. No other abdominal viscera were present. The urinary system, including a bladder or urethra, was not seen. No spinal cord structure could be seen in the sacral bone.

Fig. 3. The entire bowel loop of the fetus. The appendix (arrow) and a short segment of the large intestine are seen. Case 1.

One umbilical vein and one umbilical artery were identified in the stalk region and could be traced to the inguinal region to be connected to the internal iliac vessels. Wharton's jelly was poorly formed in the region where it was inserted into the co-twin at a length of 2cm. The placenta measured 22×20×1.0cm and was disc-shaped. A large cord was inserted into the middle of the disc and was unremarkable. There was one area where the extra-vessels were inserted and anastomosed to the placental vessels at the margin. This area was assumed to be connected to the membrane seen in the acardiac twin.

Case 2 (RCM #260)

The placenta and an irregularly-shaped acephalic monster were submitted. The monster presented a lower trunk and lower extremities (Fig. 4). The trunk measured 5.5cm and the limbs were 7.5cm in length. An umbilical cord was inserted into the midpoint of the remaining trunk. The cord, relatively well-formed, was 0.5cm in diameter. Two vessels were surrounded by Wharton's jelly. Another two vessels were connected to a vessel that was continuous to the common iliac vessels. The other end of the umbilical cord led into the placenta 3cm from the margin. Before it entered into the chorionic plate, it lost most of the Wharton's jelly and thinned. Only two vessels,

Fig. 4. This acardiac fetus consists of lower trunk and lower extremities. Unlike Case 1, a considerable amount of thorax is preserved in which several pairs of ribs are included (Fig. 8). Note that the umbilical cord (arrow) is well-formed and is inserted into the midpoint of the specimen. Case 2.

one larger than the other, were connected. On dissection of the fetus two hypoplastic kidneys and adrenals were seen. Both organs were juxtaposed in the midline. A radiograph of the fetus revealed several attenuated ribs and 15 vertebral bodies (Fig. 5). Lower bowel loops were seen. Testes were found in the pelvic cavity. The urinary bladder and atretic urethra were also found.

Case 3 (A82-23)

The acardiac twin weighed 120gms and measured 21cm. It was a globular body without a head and consisted of a well-formed trunk and lower extremities. The angle between the twisted trunk and lower limbs was about 90 degrees (Fig. 6). A rudimentary upper extremity was seen in the upper portion of the truncal mass. It appeared to be like a phocomelia, i.e., a small portion of a forearm and hand with five fingers. However, a radiograph disclosed a short humerus and one ulnar that were embedded in the edematous truncal mass. A small deficient scapula-like bone was also seen. At least 14 pairs of slender ribs and 20 vertebral bodies were noted. The pelvic bones were well-formed. The lower extremity bones were normal.

Fig. 5. Specimen radiograph showing at least six pairs of ribs and 15 vertebrae, as well as well-formed pelvis and lower extremity bones. Case 2.

Dissection of the trunk disclosed a central disc-shaped adrenal. It appeared as if two adrenals were fused together at the midline (Fig. 7). Both kidneys were located laterally to the adrenal gland. A considerable length of bowel loop was identified. It started blindly but reached the anal opening without stenosis. No meconium was noted grossly. The ureters and urinary bladder were normally formed. There were two testes in both inguinal canals. Microscopic examination of the organs and tissues of the acardiac showed a normal developing fetus corresponding to the co-twin. The umbilical cord contained one artery and one vein.

Case 4 (A83-24)

The specimen consisted of a roughly triangular placenta and two attached fetuses, one of which was an acephalic monster. The placenta measured 20×15×1.0cm. At one point, 5cm from the nearest margin of the placenta, there were two umbilical cords from two fetuses of very short length and with apparent inter-anastomosis (Fig. 8). Externally, the

Fig. 6. A well-formed trunk with rudimentary upper extremities and lower extremities are seen. An umbilical cord is seen in midpoint of this edematous fetus. Case 3.

normal fetus (Fetus A) had two umbilical arteries and one umbilical vein in the umbilical cord, while the acephalic twin (Fetus B) had a fairly slender umbilical cord that had only two umbilical vessels. To confirm the connection of the vessels barium was infused through one of the umbilical arteries of Fetus A, which opacified one of two vessels of Fetus B as well as radiating chorionic vessels on the surface of the placenta. At one point, therefore, three vessels converged, i.e., the umbilical artery of fetus A, the umbilical vein of fetus B and the chorionic vessel. The other umbilical artery of fetus A was also connected to the chorionic vessels in this region. The umbilical vein of the fetus A was also anastomosed to the umbilical artery of the fetus B on the placental surface to the chorionic vessels.

Fetus B was a monster showing generalized edema and the absence of a head and both shoulder girdles. It measured 21 cm and weighed 620gm. The lower two-thirds of the trunk and lower extremities were relatively well-developed. The genitalia were of a male, with a small phallus and urethral opening. The scrotum was hypoplastic and edematous and lacked tes-

Fig. 7. Internal view of the abdominal viscera after dissection. It shows a midline fused adrenals, kidneys and testes. Case 3.

tis. There was a round protrusion (omphalocele) just above the umbilicus. The umbilical cord lacked Wharton's jelly almost completely and was represented by a slender cord comprised of two vessels. One vessel of the umbilical cord was connected to the descending aorta just above the opening of the renal artery. Both renal arteries entered into the kidneys that were slightly displaced medially. The upper poles of the two kidneys were very close but not fused. No adrenal glands were noted. The midtrunk was represented by a thoracic spine and surrounding soft tissue which showed a marked edema. A diaphragm, lung or other thoracic structure was totally absent. The heart was non-existent and represented by only a few vascular channels that supplied and drained the soft edematous tissue in this region. Both ureters ran into the urinary bladder. Both testes could be detected in the internal inguinal ring region. They were histologically normal. Bowel loops were seen. The bowel started as a blind end and extended 13cm in total length. The large intestine was complete as well as the ileocecal valve. However, a small bowel loop was hypoplastic.

Fig. 8. Twins that are interconnected by umbilical vessels on the placental surface. The twin B is headless and edematous. Note the discrepancy of the size of the umbilical cords. Barium injected into one of two umbilical vessels causes opacity of the umbilical vessels of Twin A as well as vessels on chorionic surface. Case 4.

Case 5 (CHA89-57)

This severely edematous fetus weighed 2,190gm and measured 38cm (Fig. 9). On the left side of the midbody, a structure thought to be a face was present. A tongue was prominent. The upper extremities and head were absent, but vertebrae and lower extremities were relatively well-developed. An omphalocele was noted. In the pubic area, labia and anus were present. On radiograph, the vertebral column was severely bent, convex to the left side. Corresponding ribs were also found. A rudimentary skull base was present. On section of the truncal mass, a large amount of orange-yellow fluid escaped from many loculated cavities. On opening the abdominal cavity, a large right kidney and adrenal were seen. The left kidney was absent. Two ureters arose from the right kidney. The liver, pancreas and spleen were absent. The intestinal loop was 40cm in length and started as a blind loop. In the thoracic cage there were no lungs, heart or esophagus. On the top of the spine a small glial mass, measuring 0.5×0.3×0.3cm, was also found. On the cervical spinal cord a fluid-filled arachnoid cyst was recognized.

Case 6 (A82-109)

This acardiac twin was delivered as the second twin weighing 2,175gm fetus and was grossly abnormal. It measured 23cm in length. Generalized edema was apparent. A large head part, a medium-sized truncal part and extremities were well-recognizable (Fig. 10). The facial structure was not individually identifiable. However, several skin tags with depressed areas were noted in the anterior midportion. Rudimentary ear lobes were present. But an external auditory canal was atretic bilaterally (Fig. 11). Posteriorly in the neck multiloculated cystic lymphangioma could be seen (Fig. 10). A whole-body radiograph (Fig. 12) demonstrated a definite skull with an orbit and maxilla and an entire vertebral system with rib bones. At the region of the umbilicus there was a small skin defect through which intestinal loops protruded. No covering membrane was noted, and this area corresponded with the umbilicus. The upper and lower extremities were well-formed.

Dissection of the fetus showed no heart. A small pinkish-gray nodule that turned out to be lung tissue was found connected to one artery and a vein (Fig. 13). This extremely small lung was microscopi-

Fig. 9. External view of the acardiac fetus. Inside the edematous upper portion there were multiloculated lymphangioma, rudimentary skull and thorax. Case 5.

cally normal in architecture and maturity for its gestational age (33 weeks). This lung tissue was also connected to the trachea. In the region of the heart there was a round cyst which was histologically an enterogenous cyst. The kidneys showed multiple cortical hamartomas, but no cystic change was seen. The abdominal cavity contained no traces of a liver, spleen or pancreas. The brain weighed 60.2gm and showed a typical holotelencephaly. There was no pineal gland. The olfactory bulbs were bilaterally missing. There was oligodactyly. Present were a rudimentary right little finger, a hypoplastic left little finger, and four digits on the left and right feet.

The umbilical cord contained one artery and one vein. The aorta lacked an aortic arch and was connected to both common carotid arteries. The intestine showed multiple patches of atresia where the mesenteric root was defective. The subclavian, renal and mesenteric arteries showed normal branching from the aorta. One umbilical artery was connected to the left common iliac artery. The superior and inferior venae cavae were anastomosed directly

Fig. 10. Posterior view of the acardiac twin, showing a large edematous head as well as well-developed trunk and extremities. Posterior neck shows cystic hygroma. Case 6.

without the heart, and other venous branching was normally formed (Fig. 14).

Case 7 (A86-32)

The submitted specimen was a severely macerated fetus which was allegedly expelled after the first baby. No further history was available. It was a grossly abnormal female fetus with an abnormally formed small head, broad neck with cystic hygroma and arthrogryptic lower extremities with multiple digital anomalies. The fetus weighed 45.2gm and measured 7.8cm in crown-rump length and 11.3cm in crown-heel length. The chest and abdominal circumferences were 7.5cm and 7cm, respectively. The head was small, measuring 2.0x2.6cm. The biparietal diameter was 2.1cm. The left foot length was 1.5cm. The right foot showed a fourth to fifth syndactyly, and the left foot had only four toes. The right hand had three fingers with a deficient thumb. The left hand showed a fourth to fifth syndactyly.

Dissection was difficult because of marked autol-

Fig. 11. Anterior view of the fetus showing ear-like structures (E) and depressed nostril-like structure (N) in the head. Also note an omphalocele (arrow) in midabdomen. Case 6.

Fig. 13. Dissection of the thorax shows a round mass (lung, L) attached to the trachea (T). There were two vessels connected to this tissue. Case 6.

ysis. However, it was quite clear that this fetus had no heart. Instead, a round mass (lung) was seen in the right hemithorax (Fig. 15) that was connected to the trachea and eventually to a well-developed ton-

Fig. 12. Specimen radiograph of the fetus. the entire skeletal structure, including a well-formed cranial vault, is seen embedded in edematous subcutis and loculated cystic hygroma. Case 6.

gue. The vascular connection could not be traced. However, definitive vascular channels comparable to Case 6 could be seen. Two parallel vessels could be dissected. The liver was in the midline, and there was no definite spleen. The right kidney was absent, and the adrenals were fused in midline.

The head part showed a cleft palate and lip in midline. There was a definite mouth opening. The nose and eyeballs could not be found. A skin tag in the ear region was seen on the right side, but no opening was noted. There was a mass of brain tissue amounting to 7.3gms. A more detailed observation was impossible because of maceration. The intestinal loops were unremarkable. The fallopian tubes, uterus and ovaries were seen.

Case 8 (CHA86-23)

The specimen consisted of two fetuses and a placenta. One fetus (Fetus A) was an externally normal male baby. The other fetus (Fetus B) was a grossly monstrous male baby who had an amorphous

Fig. 14. Schematic drawing of the circulatory system is dependent on the twin litter through the single umbilical artery (UA) and an umbilical vein (UV) that are in reverse direction of flow of twin litter. ccc: common carotid artery, cia: common iliac artery, ivc: Inf. vena cava, PA: Pul. artery, sca: subclavian artery, svc: Sup. vena cava, tra: trachea.

edematous head portion, rudimentary upper extremities and a relatively preserved trunk and lower extremities (Fig. 16). The right upper extremity was missing. The fetus measured 20cm in length. The head portion was generally edematous, and the scalp was partly covered by dysplastic black hair. The hairline was poorly formed. The anterior aspect showed a mid-line hole in the center, and to the lateral of this hole were skin corrugations that mimicked auricles. However, an ear lobe or ear canal was not noted. There was no nose or mouth structure. The genitalia were ambiguous externally. Dissection of the edematous head showed a microphthalmic eyeball under the hole seen externally. A vestigial skull bone was seen, in which no identifiable brain tissue was included. The chest was dissected to show no cardiac structure. The lungs and thymus were seen. The stomach, and small and large intestines were well-developed. Both adrenals and kidneys were of normal size and

Fig. 15. Two parallel vessels are seen in prevertebral region representing inflow and outflow tracts of the fetal circulation. Case 7.

in their normal locations. Both ureters and the urinary bladder and urethra were all normal. The testes were in the abdominal cavity.

The placenta was partly fragmented, particularly where the two umbilical cords were emerging from it. Both cords were inserted marginally in a velamentous fashion (Fig. 17). The umbilicus of the fetus was of normal shape and size and contained two arteries and one vein. Fetus B had two small attenuated vessels that were connected to the vessels of Fetus A, as well as chorionic vessels on the placental surface. The arterial system of Fetus B consisted of an aorta and its branches and extended cranially into the edematous head portion without forming heart. Parallel-running venous channels were also seen that appeared to drain into the venous system connected to Fetus A.

Case 9 (CHA87-54)

The fetus was grossly abnormal, showing a markedly edematous head and upper trunk, absent

Fig. 16. Gross appearance of the normal and acardiac fetus. The acardiac twin is stunted, edematous and lacks a right upper extremity. Case 8.

Fig. 17. The placenta shows velamentous insertion of the umbilical cord connected to the vessels of Fetus A (normal) on placental surface.

upper extremities and a relatively preserved lower trunk and lower extremities. The fetus measured 33cm in crown-heel length. The head portion, 13.5x10cm, was particularly edematous and showed a nose-like protrusion (proboscis) at a point on the

Fig. 18. Gross appearance of the specimen showing recognizable head part, trunk and lower extremities. A proboscis (arrow) is apparent in mid upper face. Case 9.

upper one-third of the anterior head (Fig. 18). Eyes and ear were not found. Dissection failed to reveal any vestige of an eyeball or petrous bone. However, a small rudimentary oral cavity was seen, in which a cleft tongue was included in the left lateral side. While the left mandible was present, the right mandible was absent. The trunk contained poorly-formed cardiac and lung structures. There was a small heart measuring 14x12x10mm and weighing 0.8 gms. The right atrium was relatively large in contrast to the left atrium which was smaller. There was a round defect, 3mm in diameter, in the atrial septum (Fig. 19). The tricuspid valve was poorly formed, but the right ventricular chamber was definite with traceculae carnae. No ventricular which did not directly lead to the fused lung (Fig. 20). The mitral valve was vestigial and connected to the extremely hypoplastic left ventricle where the aorta arose. The superior and inferior vena cavae were prominent and connected to the right atrium. These two vessels, which were connected to the two umbilical vessels of the umbilical cord, were the supplier and drainer of the vascular structure to the fetus. However, their connections to the other fe-

Fig. 19. Rudimentary cardiac tube, showing a large round atrial septal defect and tricuspid valve. Case 9.

Fig. 20. Trachea and single-lobed lung and tracheoesophageal fistula. T: trachea, L: lung, S: stomach. Case 9.

tus or to the placenta could not be traced.

An esophagus was connected to the rudimentary

Fig. 21. Dorsal view of the brain, showing a typical holoprosencephaly. No interhemispheric fissure is present. Case 9.

Fig. 22. External view of the amorphous twin in Case 4. It is covered by granular skin and is edematous. Hairs are seen on the right resembling a scalp.

oral cavity. It became atretic at the midpoint where it was attached to the trachea forming a tracheoesophageal fistula. The stomach started as a tracheoesophageal fistula end (Fig. 20). Wrapped around the second portion of the duodenum an annular pancreas was seen. A midgut was absent. A distal small bowel, ileocecal valve and complete large

intestine were present. The liver was small and located in midline without right/left asymmetry. A small gallbladder was present, but there was no spleen. There was a small lung attached to a trachea. No lobation could be seen. Abdominal viscera were generally well-developed, including a stomach, kidneys, adrenals and urinary bladder. There was a cranial bone. When it was dissected a characteristic holoprosencephalic malformation could be observed. No interhemispheric fissure could be seen. Instead, there was a large posterior cyst (Fig. 21). No septum pellucidum was seen. The midbrain, cerebellum and pons were relatively well-formed compared to the cerebral hemispheres. No olfactory tracts were seen. On X-ray, a cranial vault, vertebral column and rib and pelvic bones were well-demonstrated. Both feet showed calcaneovarus deformity and had only four digits on each.

Case 10 (A82-75)

When received this amorphous mass was completely covered with whitish skin (Fig. 22). It measured 10x7x5cm. A rudimentary slender umbilical cord was attached to the midpoint of the mass. The mass was expelled "without any connection." Externally there was no recognizable extremity structure. At one end numerous black hair shafts were seen in one area resembling a scalp. Besides, no evidence of head tissue was present in cut sections and on microscopic examination. Cut sections showed edematous fat and scattered cartilage and a bone structure without any parenchymal organs. Microscopically, the skin, subcutis, bone and cartilage were all normal.

DISCUSSION

According to the severity of developmental failure in the acardiac fetus, various names have been coined (Amatuzio & Gorlin, 1981). Of all the acardi-acs, *acardius anceps* (or *paracephalus* or *aceps*) appears to be the most differentiated. Usually a partly-developed head is present, with remnants of cranial bones and brain tissue. The body and extremities are developed. There may or may not be a definite thoracic cavity. Cases 6, 7, 8 and 9 belong to this type. The faces were imperfect, with various clefts in places of ears, eyes or mouth. Accordingly, when a brain was present it was holoprosencephalic probably secondary to prechordal mesodermal defect. It is well known that a midfacial anomaly often results in the failure of the cleavage of the telen-cephalon (Chi & Kim, 1984).

Acardius acephalus is the most common of the acardi-acs that have been reported. They are head-less and lack thoracic organs. The ribs are usually developed, but thoracic vertebrae are either lacking or reduced in number. Pelvic bones and the lower extremities are well-developed, but the upper extremities may or may not be present. Of the 10 cases in this series, five were of acephalic acardi-acs. There was a wide spectrum of truncal agenesis. The lower extremities were present in all cases, although they were often deformed. The liver was usually missing. When present, it was in midline without right/left asymmetry. A rudimentary lung was seen in two cases. The gastrointestinal tract of various length was always present. The kidney and adrenal were commonly seen.

Acardius amorphus is the least-developed monster that is not recognizable as a human form. Its interior contains bone, cartilage, fat, fibrous and muscular tissue and blood vessels. It is different from a dermoid cyst in that it is attached to an umbilical cord. In this series one case (Case 10) belonged to this category. This is the type which can often be mistaken as a tumor. We agree with others that radiographic study of such monsters should be made in every instance until more detailed radiographic and anatomic studies reveal external and internal constituents sufficient to propose a reliable detailed classification on this rarity. *Acardius acormus* is the rarest form of acardia, characterized by a head without a body. The head is attached to the placenta either directly or via an umbilical cord. No such case was encountered in this series.

It was interesting to note in this series that most cases disclosed direct large artery-to-artery and vein-to-vein anastomoses between the two fetal circulations. Most authors (Richard & Benirschke, 1963; Severn & Holyoke, 1973; Lachman & McNobb, 1980; Van Allen & Smith, 1983) believed that the development of acardiac fetuses occurred only in the presence of a large artery-to-artery and vein-to-vein anastomoses, and they hypothesized that a restriction to a venous return of the acardiac is the primary event for the reversal of circulation. This obstruction was held to be the most frequently encountered om-phalocele among acardi-acs. The omphalocele was encountered in three out of 10 cases in this series. However, there were also various anomalies in the body stalk formation that could have potentially hindered the smooth circulation through the umbilical cord. It is agreed that the acardiac is a parasite dependent upon the driving force of the normal twin's heart and that blood reaches it by a reversal of the

normal flow. Oxygen and nutrient-depleted umbilical artery blood leaves the normal twin and is driven into the acardiac by way of anastomoses with its umbilical artery. Originally it was felt that with this reversal in cardiac dynamics there was a secondary diffuse atrophy of the monster's heart (hemiacardius or acardius) and secondary atrophy of other parts of the body (Van Allen & Smith, 1983). However, Potter (1961) feels that there is a primary developmental failure rather than atrophy of any parts. It is thought that the primitive cardiac tube is molded by the streaming of blood through it, determining the subsequent development of the atrial and ventricular septal sites. Such data suggest that the reversal of the blood flow in the acardiac fetus during the crucial period of cardiogenesis may cause a number of cardiac anomalies, including the cessation of development or complete cardiac agenesis.

After reviewing these cases it seems true that there is anomalous development of the fetal placental circulation to begin with. These hypoxic and nutrient-deficient conditions variously derange the primary organizing and induction pattern in such a way that subsequent development may involve innumerable combinations of arrests with a wide spectrum of rudimentary, often bizarre, growth and development. In this context Case 9 in this series is of special concern in that a primitive tubular heart was found. It suggests that the heart was arrested during its development rather than atrophied after it was once formed. The sparseness of building materials for the second fetus might have caused failure of further development. As for the development of certain particulate tissue, this can be explained by the fact that anlage may proceed to form portions of an intestine or a length of femur. All of the cases in this series appeared like a fetus that had been arrested in its development. When the head portion was preserved, it was a large edematous head with undeveloped facial features. Most acephalic monsters exhibited well-developed hind limbs. It indicates that the circulation of unsaturated venous blood is delivered to the umbilical artery of the abnormal fetus, and hypoxic blood is circulated to the lower limbs and then

cephalad.

Acknowledgements: Dr. Jin-Hee Sohn (Case 3), Dr. Hye-Kyung Lee (Cases 6 and 10), Dr. Eun-Sil Yu (Case 4), Dr. Hyo-Min Kim (Cases 7 and 8), Dr. Yeon-Lim Suh (Case 9) and Dr. Tae-Jin Kim (Case 5) performed postmortem examinations.

REFERENCES

- Amatuzio JC, Gorlin RJ: *Conjoined acardiac monsters*. *Arch Pathol Lab Med* 105:253-255, 1981.
- Benirschke K, Harper VDR: *The acardiac anomalies*. *Teratol* 15:311-316, 1977.
- Chi JG, Kim SB: *Neuroanatomic Study on Holoprosencephaly*. *Seoul J Med* 25:359-374, 1984.
- Gillim DL, Handricks GH: *Holoacardius*. *Obst & Gynec* 2:647-653, 1953.
- Gutmacher AF: *Factors affecting normal expectancy of conception*. *JAMA* 161:855-860, 1956.
- Kim Sh, Song HS, Kim YC: *A case of acardiac fetus*. *Korean J Path* 5:53-65, 1971. (in Korean).
- Kim KH, Park JS, Lee M, Lee HK, Shin SS, Chi JG: *A case of acardiac twin*. *J Korean Ped Ass* 27:84-88, 1984 (in Korean).
- Lachmann R, McNobb M, Furmansk M: *The acardiac monster*. *Eur J Pediatr* 134:195-200, 1980.
- Napolitani FD, Schreiber I: *The acardiac monster*. *Am J Obst & Gynec* 80:582-589, 1960.
- Potter EL: *Pathology of the fetus and the infant*, 2nd ed. Year Book Publishers inc, Chicago, 1961.
- Richard R, Benirschke K: *Holoacardius amorphus*. *Am J Obst & Gynec* 86:329-332, 1963.
- Schwalbe E.: *Die Morphologie der missbildungen des Menschen unter Tiere*. Fischer, Jena, Germany, 1907.
- Severn CB, Holyoke EA: *Human acardiac anomalies*. *Am J Obstet Gynec* 116:358-365, 1973.
- Van Allen MI, Smith DW, Shepard TH: *Twin reversed arterial perfusion (TRAP) sequence: A study of 14 twin pregnancies with acardius*. *Semin. Perinatal* 7:285-293, 1983.
- Wilson EA: *Reviews of holoacardius*. *Obst & Gynec* 40:740-748, 1972.