New Antiepileptic Drugs

II. Clinical Use

Byung-In Lee, M. D.

Department of Neurology, Yonsei University College of Medicine

All new AEDs underwent extensive phase three clinical trials before their approval for marketing, which provided reliable references for their clinical utility. The clinical trials were conducted most commonly in either double blind placebo controlled(DBPL) crossover or parallel group designs of add-on therapy. For FBM, innovative monotherapy designs such as monotherapy using attenuated control drug(15mg/kg of VPA) or placebo controlled monotherapy in surgical candidates have been used to document its antiepileptic efficacy. The primary target of all clinical trials was partial seizures in adulthood, which was usually followed by open-label or single-blind trials in childhood epilepsies.

1. Efficacy

A. Vigabatrin(VGB)

1) add-on therapy in adults

Clinical trials of VGB were initiated in 1982 and 6 small scales of DBPC-crossover add-on trials were conducted(table 1). Five trials demonstrated significant improvement of seizure control with a mean/median seizure frequency reduction of 37% to 54% and a responder rate($\geq 50\%$ seizure frequency reduction) of 33% to 67%. The meta-analysis of these trials indicated a 46% responder rate in a total of 98 patients having only partial seizures(Mumford and Dam, 1989). Increase of seizure frequencies after VGB add-on therapy was also observed in 0% to 20% of study patients. Altogether 6% of study patients discontinued VGB due to an exacerbation of seizures(Grant and Heel, 1991).

A recently published US multicenter DBPC-parallel group study(French et al., 1996) recruited 182 patients with partial epilepsy. Median seizure frequency reduced

by 37% in VGB group compared to 9.6% in PLC group($P=0.002$). Responder rate was 43% in VGB group compared to 19% in PLC group($P<0.001$). Seizure free outcome was observed in 5 of 92 patients (4.5%) who received VGB but none in PLC group.

The long-term efficacy of VGB was observed in several open-label follow-up studies. Tartara et al.(1989) followed 25 patients who responded favorably to VGB in an initial DBPC-crossover study for up to 47 months (median 22 months). Two patients discontinued because of adverse effects and 3 patients did so because of seizure breakthrough. In a US multicenter trial(Browne et al., 1991), 14 of 66 patients(21%) shown an initial favorable response to VGB discontinued VGB for seizure breakthrough with or without adverse effects in long-term follow up(median 43 months). Pitkainen et al. (1993) reported that 5 of 15 patients(33%) who were the responders to an initial 3 months of VGB therapy lost that response during a 5-year follow-up period. In a literature review, Grant and Heel(1991) suggested that approximately 20% of initial responders to VGB discontinued VGB because of insufficient efficacy rather than the appearance of adverse effects. In a data base of about 2,000 patients, loss of efficacy in long-term therapy occurred in 4.4% overall.

In Korea, three clinical trials of open-label VGB add-on therapy were conducted. Lee et al.(1993) studied 16 patients with various seizure types and reported 56% of responder rate. Hwang et al.(1994) reported 57% of responder rate in 14 patients with partial epilepsy taking VPA. In a study by Lee et al.(1993), 12 of 28 patients (43%) with partial epilepsy achieved greater than 50% seizure reduction compared to the baseline and 2 patients(7%) became seizure free. In a long-term treatment for up to 21 months, 70% of patients maintained its beneficial effects.

2) monotherapy in adults

Kälviäinen et al.(1995) conducted a comparative

Address for correspondence : Department of Neurology,
Yonsei University College of Medicine, 134 Shinchon-
dong Seodaemun-gu, Seoul 120-752, Korea.

Table 1. Summary of double-blind placebo controlled trials of vigabatrin as add-on therapy in partial epilepsy

	Baseline Sz. freq. (mean or median)	Study design	No. of pts	Dosage (g/day)	Treatment period (weeks)	Sz frequency		
						Overall decrease (mean/median%)	Decrease $\geq 50\%$ (% pts)	Increase (% pts)
Rimmer & Richens (1984)	5.8/wk	DBPC-crossover	24	3	9	42 (P<0.001)	67	0
Gram et al (1985)	> 4/month	DBPC-crossover	21	3	12	44 (P<0.05)	44	17
Loiseau et al (1986)	10.7/wk	DBPC-crossover	27	3	10	41 (P<0.01)	58	10
Tartara et al (1986)	4.8/wk	DBPC-crossover	21	2-3	7	54 (P<0.01)	60	20
Remy et al (1987)	7.3/wk	DBPC-crossover	19	3	12	37(CPS) (P<0.05)	6	67 absences
Tassinari et al (1987)	> 6/8wk	DBPC-crossover	39	2-3	12		33	17
French et al (1996)	$\geq 6/2m$	DBPC-parallel	182	3	16	36-VGB 9.6-PLC (P=0.0002)	43%-VGB 19%-PLC	

monotherapy trial in one hundred newly diagnosed patients with partial and/or generalized tonic-clonic seizures. VGB and CBZ both achieved satisfactory seizure control in 60% of patients, however, the seizure free rate was higher in the CBZ treatment group than the VGB treatment group (52% vs 32%; $P<0.01$). Drop out rate was 26% for VGB with all due to lack of efficacy and 30% for CBZ treatment group with 24% due to adverse effects and 6% due to lack of effect. Another study by Tanganelli and Regestra (1994) reported that, in 39 patients with partial epilepsy, seizure free was achieved in 42% of VGB and 55% CBZ monotherapy. However, the incidence of AE was lower in VGB monotherapy than CBZ monotherapy (21% vs 39%).

These monotherapy trials suggested that the anti-epileptic potency of VGB in monotherapy was slightly less than CBZ, however, VGB was more easily tolerable.

3) children.

A few controlled trials of VGB in children have been published. Multicenter European open-label add-on pilot study in 133 patients with various seizure types (Livingstone et al., 1989), 34% of patients responded favorably by achieving greater than 50% seizure frequency reduction. However, 53% of patients discontinued VGB due to lack of efficacy. In a single blind study of VGB add-on therapy in 66 patients with various seizure types (Dulac et al., 1991), responder rate was 46% but 15% of their patients experienced deterioration of

seizure control. The meta-analysis of 360 pediatric patients (Appleton, 1993) showed a responder rate of 50% in refractory childhood epilepsy.

The efficacy of VGB in children with partial seizures was recently reported by Bernadina et al. (1995). In this single-blind multicenter trial, PLC was administered for 1 month and was followed by VGB treatment for 6 months. The initial dose of VGB was 40mg/kg/day and increased to 60mg/kg/day and 80mg/kg/day at 2 month intervals. Of 46 children enrolled in the study, 7 patients dropped out prematurely due to lack of efficacy and exacerbation of seizure in one. The mean number of seizures per month decreased from 97 during PLC period to 21, 12, and 9, after 2, 4, and 6 months of VGB treatment respectively ($P<0.0001$ at each time). The number of patients who had greater than 50% seizure frequency reduction after 2, 4, and 6 months was 28, 33, and 35 respectively. Eight patients also became seizure free during the last 2 months of VGB treatment.

The remarkable efficacy of VGB in infantile spasm was reported by Chiron et al. (1991). Among 70 children with refractory infantile spasms, 68% achieved greater than 50% seizure frequency reduction with seizure free in 43%. Efficacy was particularly impressive in patients with symptomatic infantile spasm due to tuberous sclerosis.

4) efficacy in relation to seizure types

Subgroup analysis of clinical trials clearly demon-

strated the best efficacy of VGB in partial seizures and the efficacy in secondary GTC seizures was less remarkable (Rimmer and Richens, 1984; McKee et al., 1993; French et al., 1996). Tassinari et al. (1987) also reported no significant improvement in patients with mixed seizure types, multifocal EEG abnormalities, and high seizure frequency. In childhood epilepsy, Lunar et al. (1989) reported best responses to VGB in cryptogenic partial epilepsy and frequent exacerbation of non-progressive myoclonic epilepsy, Dulac et al. (1991) and Remy et al. (1986) reported frequent exacerbation of absence seizures, Bernadina et al. (1995) reported remarkable efficacy in both cryptogenic and symptomatic partial epilepsy. Infantile spasm, especially caused by tuberous sclerosis, is another epileptic syndrome responding favorably to VGB.

5) efficacy by dose

The usual daily dose of VGB in clinical trials has been 2 to 4 gm/day for adults. Mumford and Dam (1989) reported that only 25% of patients had better seizure control after increase of VGB beyond 2 gm/day. In a double-blind dose-comparison study (Sivenius et al., 1987), 3 gm/day appeared more effective than 1.5 gm/day, however, McKee et al. (1993) reported even deterioration of seizure control after the increase of VGB from 2 gm/day to 3 gm/day and 6 gm/day. Rothman et al. (1994) reported no increase of cerebral GABA concentration measured by MR spectroscopy when the dosage of VGB was increased from 3 gm/day to 6 gm/day.

These findings clearly suggest that there is an optimally effective dosage for individual patients and a ceiling to the effective dosage. In adults, a dosage greater than 3 gm/day may not improve its efficacy. In children, 80 mg/kg/day seem to be the usual maximal dose of VGB.

B. Zonisamide (ZNS)

1) add-on therapy in adults

Two DBPC-parallel group add-on therapy trials and one DB comparative study were conducted in patients with partial epilepsy (table 2).

The US multicenter trial (Wilder et al., 1986) recruited 152 patients with refractory partial epilepsy (>4 seizures/month) and the median seizure frequency decreased by 30.1% in the ZNS group compared to 0.3% increase in the PLC group ($P < 0.01$). The responder rate was 28.6% in the ZNS group and 13.2% in the PLC group ($P = 0.03$). In the European trial (Schmidt et al., 1993a), the median seizure frequency decreased by 27.7% in the ZNS group and increased by 3.9% in the PLC group. Responder rate was 30.3% and 12.7% in the ZNS and PLC groups respectively ($P < 0.02$).

Seino et al. (1988) conducted a controlled comparative study of ZNS with CBZ in 123 patients with partial seizures either untreated or refractory to conventional AED therapy. The mean seizure frequency reductions of partial seizures and secondary GTC seizures were 68.4% and 69.7% in the ZNS group and 46.6% and 70.2% in the CBZ group respectively, which suggested at least comparable efficacy of ZNS and CBZ.

In an US open-label controlled trial by Leppik et al. (1993), the median seizure frequency decreased by 40.6% in CPS, 75.3% in GTC, and 41.4% overall. In a Japanese open-label trial (Ono et al., 1988), 59% of patients with partial seizures and 53.2% of the patients with generalized seizures improved.

2) monotherapy in adults

No double-blind controlled monotherapy trials have been reported yet. A multicenter double-blind active drug controlled comparative trial in previously untreated

Table 2. Double-blind controlled trials of zonisamide in adult patients with partial epilepsy

Study	Number of patients	Seizure type	Duration of treatment	Baseline seizure frequency	Seizure improvement		Study design
					Mean/median Sz frequency reduction	$\geq 50\%$ Sz frequency reduction	
Wilder et al (1986)	152	partial	12 wk	$\geq 4/m$	ZNS : 30.1% PLC : +0.3% ($P < 0.01$)	ZNS : 26.8% PLC : 13.2% ($P = 0.03$)	DBPC-parallel
Schmidt et al (1993)	139	partial	12 wk	$\geq 4/m$	ZNS : 27.7% PLC : +3.9% ($P < 0.01$)	ZNS : 30.3% PLC : 12.7% ($P < 0.02$)	DBPC-parallel
Seino et al (1988)	123	partial	16 wk	$\geq 2/m$	partial Sz : 68.4% in ZNS 46.6% in CBZ GTC Sz : 69.7% in ZNS 70.2% in CBZ	—	DB-active drug controlled

Study patients were either untreated or refractory to conventional AED therapy.

patients is currently underway in Korea. Yagi and Seino (1992), in their data analysis, identified 55 patients treated with ZNS as the sole drug. The responder rate in these patients were 73% for partial seizures and 71% for generalized seizures.

3) children

Oguni et al.(1988) compared ZNS with VPA in 34 children with generalized epilepsy either previously untreated or refractory to conventional AED therapy. The frequency of GTC seizures decreased by 81.2% in the ZNS group and 66.7% in the VPA group. In global evaluation, 50.0% of ZNS group and 43.8% of VPA group were judged as improved. In an open-label Japanese study of 393 patients with partial or generalized seizures(Ono et al ; 1988), the improvement of seizure control was observed in 57.9% of patients with partial seizures and in 38.3% of patients with generalized tonic seizures.

4) efficacy in relation with seizure types

According to the data of a cohort of 1008 patients (Yagi and Seino, 1992), responder rate was 50% to 60% in partial seizures including secondary GTC seizures, 59% in GTC seizures, 26% for generalized tonic seizures, 67% for atypical absence seizures, 50% for atonic seizures, 43% for myoclonic seizures, and 41% for combined seizures. ZNS was equally effective in partial and secondarily GTC seizures in a double-blind study(Seino et al., 1995) and was more effective in secondarily GTC seizures in a US open-label trial(Leppik et al., 1993), supporting the assertion that ZNS is a broad spectrum AED.

Anecdotal reports of the significant efficacy of ZNS on progressive myoclonic epilepsies(PME) were reported by Henry et al.(1988) and Yagi and Seino(1992). However, in the authors limited experiences, ZNS had only modest effect on PME. The efficacy of ZNS in severe symptomatic generalized epilepsies such as Lennox-Gastaut syndrome and infantile spasm have been reported as good in more than 50% of these patients(Yamatogi and Ohtahara, 1991), however, further controlled data are required.

5) Korean Experiences

A total of 7 open-label clinical trials were conducted in Korea. The study patients were quite heterogenous and included both adults and children with various types of seizures. Overall responder rate ranged from 48% to 83%. In a study by Huh et al.(unpublished data),

35 patients were divided into 14 patients with epileptic encephalo-pathy, 6 patients with childhood onset symptomatic partial epilepsy, 12 patients with adult onset partial epilepsy, and 3 patients with other epileptic syndromes. The overall responder rate was 51% with 50% in epileptic encephalopathy, 33% in childhood onset symptomatic epilepsy, 32% in adult onset partial epilepsy. Two patients with PME of the Unverricht-Lundborg type responded well to ZNS and one patient with atypical absence seizure deteriorated after ZNS add-on therapy.

Therefore, the Korean experiences of ZNS were comparable to the Japanese experiences and supported that ZNS is a broad spectrum AED.

6) the dosage of ZNS

In the European controlled trials, the final dosage of ZNS was 7.8mg/kg in responders and 6.8mg/kg in non-responders with plasma levels of 17.0 mg/ml and 15.2mg/ml respectively. In the Japanese trial by Seino et al.(1988), the final mean daily dose was 330mg/day in the ZNS group and 600mg/day in the CBZ group. The recommended initial daily dose is 100 to 200mg for adults, and 2 to 4mg/kg for children. The usual maintenance daily dose is 200 to 400mg for adults and 4 to 8mg/kg for children and the recommended maximal dose is 600 mg for adults and 12mg/kg for children.

The therapeutic plasma level is around 20mg/ml.

C. Lamotrigine(LTG)

1) add-on therapy in adults

A total of 10 DBPC add-on trials have been conducted in patients with refractory partial seizures(table 3). All trials except one study by Sander et al.(1990) showed the significant efficacy of LTG with mean or median seizure frequency reduction of 17% to 59% and responder rate of 7% to 67%. The meta-analysis of 4 DBPC-crossover trials(Richens and Yuen, 1992) showed responder rate of 27%.

In a large scale US DBPC-parallel group study (Matsuo et al., 1994), the median seizure frequency reductions were 8% in PLC, 20% in 300mg/day LTG, and 36% in 500mg/day LTG. The responder rate in these group were 18%, 20%, and 34% respectively with statistical significance being apparent in the 500mg/day LTG group.

The data on long-term efficacy of LTG is scanty but the long-term efficacy of LTG in a few long-term follow-up studies(Sander et al., 1990) was comparable with the

Table 3. Controlled trials of lamotrigine as add-on therapy in adult patient with partial epilepsy

References	No. of patients	Dosage mg/day	Overall decrease (mean or median %)	Decrease $\geq 50\%$ (% of patients)	Study design
Binne et al (1989)	30	50-400	17*	7	DBPC-crossover
Jawad et al (1989)	21	57-400	59*	68	DBPC-crossover
Loiseau et al (1990)	23	75-300	23*	30	DBPC-crossover
Risner et al (1990)	88	300-400	25*	20	DBPC-crossover
Sander et al (1990)	18 ^a	100-300	18	NA	DBPC-crossover
Schapel et al (1993)	41	150-300	24*	20	DBPC-crossover
Smith et al (1993)	62	200 or 400	30*	18	DBPC-crossover
Matsuo et al (1993)	191	LTG 500 LTG 300 Placebo	36* 20 8	34 20 18	DBPC-parallel
Schmidt et al (1993)	21	300	22*	29	DBPC-crossover
Messenheimer et al (1994)	98	400	25*	25	DBPC-crossover

* Statistically significant improvement

NA : not available

results of short-term studies suggesting no evidence of tolerance development in long-term use.

A supra-additive effect of LTG and VGB or VPA combination therapy has been suggested. Stolarek et al.(1994) reported that the addition of LTG in patients taking VGB resulted in a reduction of overall seizure count by 37%. Panayiotopoulos et al.(1993) reported significant improvement of seizure control by LTG and VPA combination therapy in severely refractory patients, which was in agreement with the experience of another investigator(Brodie : *personal communication*).

In Korea, two open-label add-on studies were conducted. Lee et al.(1995) added LTG in 24 patients with partial epilepsy under CBZ monotherapy. Responder rate was 21% and adverse effects observed in 75% of the patients probably due to interactions between the two drugs. Park et al.(1995) investigated the effect of LTG add-on therapy in 18 patients with partial epilepsy under polytherapy which included VGB. Five of 17 patients(29%) achieved a greater than 50% seizure frequency reduction, which was comparable to the results of other add-on studies.

2) monotherapy in newly diagnosed patients

LTG is an attractive candidate for monotherapy in patients with newly diagnosed epilepsy for its wide range of efficacy across different seizure types and relatively benign side effect profile.

In a comparative study with CBZ conducted by Brodie et al.(1995), 151 of 260 recruited patients(131 LTG, 129 CBZ) completed the 48 week trial. No difference in efficacy was found between the two drugs ;

39% of LTG group and 38% of CBZ group became seizure free(partial seizures : LTG 35%, CBZ 37%, primary GTC : LTG 47% , CBZ 47%). Overall, more patients taking CBZ withdrew due to adverse events, 27% compared to 15% on LTG. In a comparative study with PHT conducted by Steiner et al.(1994) a total of 181 patients(LTG 86, PHT 95) completed the trial. Efficacy against partial and GTC seizures was comparable between the two drugs with no seizures during the last 24 weeks being achieved in 43% of LTG and 36% of PHT groups. However, more patients on PHT withdrew from the study due to adverse effects(LTG : 15%, PHT : 19%). These monotherapy trials suggested at least comparable efficacy with CBZ or PHT but better tolerability of LTG.

3) children

Although available efficacy data in childhood epilepsies is limited, encouraging results in open-label studies of LTG have been reported. Schlumberger et al.(1994) reported a responder rate of 38% in 120 children with various types of resistant seizures. LTG was quite effective in absence seizures as well as Lennox-Gastaut syndrome. Timmings and Richens(1992) found at least a 50% reduction of seizures in 10 of 11 patients with Lennox-Gastaut syndrome.

4) efficacy in relation to seizure types

In controlled add-on trials, the efficacy of LTG was mainly focused on partial seizures. In monotherapy trials, the efficacy of LTG on partial and GTC seizures was comparable with CBZ or PHT. The data on file at Well-

come Laboratory indicated comparable or superior efficacy in secondarily GTC seizures as well as other generalized seizures. Especially LTG showed a higher than 50% responder rate in absence seizures. The remarkable efficacy in patients with Lennox-Gastaut syndrome also suggested that LTG is a broad spectrum AED. It is quite likely that combination therapy of LTG and VPA will greatly improve seizure control in patients with severe symptomatic generalized epilepsies in the future.

5) dosage

The dosage of LTG in add-on therapy is depend upon the concomitant AEDs. The usual maintenance dose of LTG is 100-200mg/kg in patients not taking enzyme inducing AEDs and 500mg/day in patients taking inducers. It has been suggested that slow titration of LTG may decrease the incidence of skin rash, especially in patients taking VPA(Yuen, 1992). The plasma concentration of LTG was 1 to 4mg/ml in most clinical trials but the relationship between the plasma level and clinical efficacy have not been observed yet.

In the US trials most patients were able to tolerate LTG 500mg/day, however, only 30% of Korean patients (not taking VPA) were able to tolerate a dose higher than 300mg/day LTG, which suggested a racial differences in the tolerance of LTG.

D. Felbamate(FBM)

Controlled Trials of FBM are summarized in table 4.

1) add-on therapy trials in adults

Two DBPC-crossover trials were conducted in patients with partial epilepsy(table 4), Leppik et al.(1991) reported the statistically significant efficacy of FBM over

PLC($P=0.046$) in various efficacy criteria. However, Theodore et al.(1991) did not show any significant efficacy of FBM. In this study the plasma level of CBZ was decreased by 24%, which might be the cause of negative results.

2) monotherapy trials in adults

Two different unique designs of monotherapy trials were undertaken to demonstrate the antiepileptic efficacy of FBM. In the study conducted by Bourgeois et al.(1993), AEDs were completely discontinued for pre-surgical EEG-telemetry monitoring and immediately after the completion of study FBM and PLC were administered until the end points were reached. A significant anticonvulsant action was clearly demonstrated.

The other unique design was to compare FBM with an active attenuated control, FBM at a dosage of 3,600mg/day was compared with VPA at a dosage of 15mg/kg/day as monotherapy(Sachdeo et al., 1992; Faught et al., 1993). By all measures, a high dose of FBM was found to be superior to a beginning dose of VPA($P<0.001$).

3) children

FBM is the only AED to have undergone a DBPC-parallel add-on trial in Lennox-Gastaut syndrome(Felbamate study Group in LGS, 1993). Dosages were increased from 15mg to 45mg/kg/day over a 2-wk period in 74 patients with LGS. Outcome measures were i) number of seizures recorded on 4-hr EEG telemetry, ii) caregiver's seizure count, and iii) caregiver's global evaluation. In the caregiver's seizure count, the seizure frequency decreased by 19% in the FBM group compared to 7% in PLC group($P=0.01$). The efficacy of

Table 4. Controlled clinical trials in partial seizures with felbamate

Reference	No. of patients	Maximum dosage (mg)	Protocol type	Efficacy of FBM compared with placebo (p value)
Leppik et al., 1991	56	2,600	Double-blind, placebo-controlled, crossover	$p < 0.05$
Theodore et al., 1991	28	3,000	Double-blind, placebo-controlled, three-phase, crossover	$p < 0.06$ $p < 0.002$
Bourgeois et al., 1991	64	3,600	Presurgical evaluation: double-blind, parallel, placebo-controlled	$p < 0.04$
Faught et al., 1993	111	3,600	Double-blind, randomized, parallel : high-dose FBM vs initial dose of VPA	$p < 0.001$
Sachdeo et al., 1992	44	3,600	Double-blind, randomized, parallel : high-dose FBM vs initial dose of VPA	$p < 0.001$

FBM : felbamate, VPA : valproate
Modified from Ramsay(1993)

Table 5. Summary of three large scale controlled add-on clinical trials of gabapentin in partial epilepsy

Reference	No. patients. total (evaluatable)	Mean adjusted response ratio*	Responder rate (%)	Median decrease in seizure frequency (%)
UK Study, 1990	127(113)			
GBP 1,200 mg	61(52)	-0.192 ⁺	25.0 (P=0.049)	-29.2
Placebo	66(61)	-0.060	9.8	-12.5
US Study, 1993	306(288)			
GBP 600 mg	53(49)	-0.151	18.4 (P=0.103)	-24.3
GBP 1,200 mg	101(91)	-0.118 ⁺	17.6 (P=0.111)	-20.0
GBP 1,800 mg	54(53)	-0.233 ⁺	26.4 (P=0.007)	-31.9
Placebo	98(95)	-0.025	8.4	-5.9
Int'l Study, 1994	272(245)			
GBP 900 mg	111(96)	-0.136 ⁺	22.9 (P=0.008)	-21.8
GBP 1,200 mg	52(50)	-0.157 ⁺	28.0 (P=0.02)	-17.8
Placebo	109(99)	-0.025	10.1	-0.3

* Response ratio = $(T-B)/(T+B)$ where T is the number of seizures per 28 days during treatment and B is the number of seizures per 28 days during the baseline period.

⁺p<0.05

Modified from Ramsay(1994)

FBM was most pronounced in atonic seizure(P=0.002). Global evaluation scores were significantly better with FBM but the number of seizures counted on 4-hr EEG-telemetry was not significantly different.

It has been suggested that FBM may be effective in other generalized seizures such as absence(Devinsky *et al.*, 1992) or juvenile myoclonic epilepsy(Sachdeo *et al.*, 1992) when other drugs failed but they have not been confirmed by controlled studies. The long-term efficacy of FBM has not been well investigated(Theodore *et al.*, 1995).

E. Gabapentin(GBP)

1) add-on trials in adults

Earlier studies of GBP were conducted as dose ranging studies without regard to seizure types. In a controlled trial of GBP add-on therapy by Sivenius *et al.*(1991), the decrease of seizure frequency from baseline was 13.0% in patients receiving 1200mg/day (P=0.01), 6.5% in patients receiving 900mg/l(P=0.42) and 6.0% in patients receiving PLC(P=0.26). The pivotal clinical trials evaluated the efficacy of GBP were three large, randomized, multicenter, double-blind, placebo-controlled parallel-group clinical trials in patients with refractory partial seizures(table 5).

Patients with at least four partial seizures per month for 3 months prior to study entry were enrolled and randomized to receive either GBP or PLC for 12 weeks. All three large trials clearly demonstrated the efficacy of GBP in refractory partial seizures with a responder rate of 17.6% to 28% in patients who received 600 to

1800mg/day. In the US trial, there was a clear linear dose-response effect with maximum benefit observed at 1800mg/day GBP. Analysis by seizure type revealed that GBP reduced the frequency of all types of partial seizures and its efficacy was more pronounced in secondary GTC seizures.

A comparative pilot trial of GBP(1200mg/day) with CBZ(1200mg/day) was undertaken in a double-blind three way crossover study in active epilepsy(Wilensky *et al.*, 1992). Median seizure frequency was significantly lower in the CBZ group but side effects were more common compared to the GBP group. The dosage of GBP was probably inadequate to attain seizure control. A higher dose of GBP is currently under evaluation by using the same protocol.

The maintenance of the efficacy in long-term GBP therapy has been well demonstrated in the follow-up observations(18 months) of both international and US GBP trials as well as in several open-label follow-up studies(Ramsay, 1994).

F. Oxcarbazepine(OXC)

The clinical trials of OXC were mainly focused on the comparison with its parent compound, CBZ.

A large scale Scandinavian multicenter DB-comparative controlled trial with CBZ in 235 newly diagnosed epilepsy patients showed no differences in the seizure control rate(Seizure free : 60% OXC vs 52% CBZ) and global evaluation scores(good or excellent : 96% OXC vs 97% CBZ) between the two drugs(Dam *et al.*, 1989).

A smaller scale DB-crossover study comparing OXC and CBZ in 48 patients with refractory epilepsy showed

slightly better seizure control with OXC, which was not significant statistically (Houtkopper *et al.*, 1987). It seems quite likely that OXC has a similar degree of anticonvulsant activity as CBZ with the equipotent dose being about 50% higher than that of CBZ.

G. Topiramate (TPM)

Five DBPC-parallel group multicenter add-on trials were conducted in a total of 534 adult patients with refractory partial epilepsy (table 6). Two dose-ranging studies included target daily dosages of 200, 400, and 600mg/day (protocol YD) and 600, 800, and 1000mg/day of TPM (protocol YE) respectively. The remaining 3 studies evaluated single target dosages of 400 (protocol Y1), 600 (protocol Y2), and 800mg/day (protocol Y3) of TPM (Ben-Menachem, 1995).

Improvement over placebo was statistically significant in the average monthly seizure rate and responder rate at target doses of 400mg/day and higher. The 200mg/day group showed a strong trend favoring TPM over PLC ($P=0.051$) and is considered to be the minimally effective dose (RWJPR clinical database). Reductions in average monthly seizure frequency were most marked at the 400mg/day (45%) and 600mg/day (48%) dosage level with the responder rate of 46% and 47% respectively. The difference between TPM and PLC in

median percent seizure reduction varied from 16.5% for the 200mg group to 58.6% for the 600mg/day group, which suggested a therapeutic dosage range of 200 to 600mg/day in add-on therapy.

The efficacy of TPM over PLC was apparent in both partial and secondarily generalized tonic-clonic seizures. No development of tolerance was evident in 469 patients who received TPM for over 6 months in open-label, long-term studies and study extension (Reife, 1994).

2. Adverse Effects

All new AEDs passed through well designed placebo-controlled add-on trials to be found well tolerable and safe without causing serious systemic side effects. However, the duration of these clinical trials was too short to investigate the long-term adverse effects (AE) and their effects on the teratogenesis have not been investigated yet. Another point to be addressed is that a significant proportion of AE found in these trials might be related to the pharmacodynamic interactions between conventional and new AEDs.

A. Vigabatrin (VGB)

As adjunctive therapy in over 2,000 patients studied VGB has been well tolerated with AE mainly related to CNS symptoms. The incidence of AE varies greatly in different studies but usually occurred in one third of the study patients with most frequent AE being somnolence. The data on file of MMD shows that the AE occurring more often in VGB group (3g/day) than PLC group with the difference in percentages of incidence between the two groups were drowsiness (10.1%), dizziness (9.5%), paresthesia (7.4%), vision abnormality (7.3%), fatigue (7.2%), depression (6.6%), tremor (5.9%), amnesia (4.4%), confusion (3.6%), and ataxia (2.9%).

In a long-term study by Tartara *et al.* (1992), most patients did not report any A.E. but one patients dropped out for weight gain. In pediatric studies, the most common AE was behavioral hyperactivity and aggression with incidence of 9-15% (Fisher and Kerrigan III, 1995).

Earlier studies reported induced psychosis in only a few patients, however, Sander *et al.* (1991) reported 14 cases of psychosis in patients with intractable seizures taking VGB. Five of them had a history of previous psychosis and psychosis was resolved in all cases with withdrawal of the drug. A review by Sabers and Gram (1992) indicated the incidence of psychosis as 3-6%. It

Table 6. Efficacy of topiramate in controlled trials

Protocol /Target topiramate dosage	N	% Reduction in seizure rate	% Treatment responders
Protocol YD			
Placebo	45	13.1	18%
200 mg/day	45	29.6 ($P=0.051$)	27%
400 mg/day	45	47.8**	47%*
600 mg/day	46	44.7***	46%*
Protocol YE			
Placebo	47	1.2	9%
600 mg/day	48	40.7***	44%***
800 mg/day	48	41.0***	40%***
1000 mg/day	47	37.5***	38%***
Protocol Y1			
Placebo	24	1.1	8%
400 mg/day	23	40.7 ($P=0.065$)	35%***
Protocol Y2			
Placebo	30	-12.2	10%
600 mg/day	30	46.4**	47%***
Protocol Y3			
Placebo	28	-17.8	0%
800 mg/day	28	38.8**	43%***

* $\leq P 0.05$ (PLC compared)

** $\leq P 0.01$ (PLC compared)

*** $\leq P 0.001$ (PLC compared)

- Negative number denotes an increase in seizure rate.

is generally recommended to be cautious in administering VGB in patients with prior history of psychosis. The mechanism of VGB induced psychosis is unclear, however, a SPECT study with depamine D2 receptor ligand(Ring et al., 1992) showed decreased binding in the left basal ganglia after VGB treatment, which might be relevant to the development of psychosis in vulnerable patients.

Effect of VGB on various laboratory tests did not show any significant abnormalities compared with PLC. Studies on the effect on cognition(Dodrill et al., 1992; McGuire et al., 1992) did not suggest any cognitive adverse effects by VGB.

The drug withdrawal rate due to AE in controlled trials varied from 1.4% to 9% and in a recent large DBPC-parallel group study(French et al., 1996): 9 of 92 patients(9.8%) receiving VGB 3g/day withdrew due to AE compared to 2 of 90 patients(2.2%) with PLC. In this study Physicians' Overall Assessment of Tolerability suggested that 90% of the VGB group and 98% of the PLC group had an assessment of fairly well tolerated or better.

In a comparative monotherapy trial of VGB with CBZ (Kälviäinen et al., 1995), the incidence of AE was comparable(72% in VGB vs 86% in CBZ), however, none in the VGB group withdrew due to AE alone compared to 12 of 45 patients(26.7%) taking CBZ having withdrawn due to intolerable AE.

All these studies clearly suggest that VGB is well tolerable and safe new AED.

B. Zonisamide(ZNS)

The incidence of AE in two controlled trials(Wilder et al., 1986; Schmidt et al., 1993a), was 92.3% and 59.2% in ZNS group and 58.1% and 27.9% in PLC group respectively. Thus the differences in the incidence of AE between ZNS and PLC group in these studies were 33% and 31%, which was quite consistent. In the European study, the most frequent AE was fatigue seen in 22.5% of the ZNS group compared to 11.8% in the PLC group. The differences in incidence of AE between ZNS and PLC were highest in dizziness(12.5%), which was followed by ataxia(11.3%), anorexia(11.2%), fatigue (10.7%), abnormal thinking(9.8), somnolence(9.7%), nervousness(7.0%) and confusion(4.0%). Two of 71 patients withdrew from ZNS due to AE and no significant abnormalities in the Lab. tests were observed.

In the comparative trial of ZNS with CBZ by Seino et al.(1988), AE occurred in 51.7% of the ZNS group and 56.9% of the CBZ group. Anorexia was observed more

frequently in the ZNS group(12% vs 0%, $P=0.02$) and ataxia was more frequent in CBZ group(35% vs 9%, $P<0.01$).

The Japanese data of 1,008 patients analyzed by Seino et al.(1991) showed that the incidence of AE was 51.3% and 185 patients(18%) discontinued ZNS accordingly. The most commonly observed AE were somnolence(24%), ataxia(13%), anorexia(11%), G-I symptoms(7%) and decrease of spontaneity(6%). In patients under ZNS monotherapy, the incidence of AE was significantly less; drowsiness(9%), anorexia(7%), G-I symptoms(7%) and decrease of spontaneity(6%), headache(6%), skin rash/itching(6%) and weight loss(6%). With respect to laboratory abnormalities, 17 patients (2%) discontinued ZNS treatment due to an elevation of hepatic enzymes or leukopenia. The analysis of Korean open-label trials showed that AE developed in 107 of 178 patients(60.1%) with the most frequent AE being G-I symptoms(22%) and anorexia(22%); Other AE were weight loss(11.8%), headache(10.1%), dizziness(10.1%), somnolence(8.4%), sleep disturbances(6.2%), skin rashes(5.6%), memory disturbances(4.5%), fatigue(4.5%). Sixteen of 178 patients(9%) discontinued ZNS treatment due to AE during the trial.

During the US and European trials, the development of nephrolithiasis became a safety concern. In US multicenter open trials, symptomatic nephrolithiasis occurred in 13 of 700 patients(1.86%)(Henry and Sackellares, 1991). In another open trial, Leppik et al.(1993) reported that 4 of 93 patients(3.7%) developed renal calculi within 250-477 days after starting ZNS. In contrast, only 2 of 1,008 patients receiving ZNS during phase two and three trials in Japan developed renal stones and in the postmarket survey of 2½ years, only one of 2,444 patients had developed renal stones(Seino et al., 1995). In Korea, development of symptomatic renal calculi due to ZNS has not been reported since its introduction in 1992. These differences between western and oriental experiences are difficult to explain. Yagi et al.(1992) measured 24hr urine volume and the excreted amount of calcium, magnesium, citrate, and phosphate in epileptic patients, which showed a significantly decreased excretion of citrate only. Schmidt et al.(1993) suggested possible influences of genetic, environmental, and dietary factors. It may be interesting to observe the results of future clinical trials of TPM in the Orient, which is another sulfonamide compound to be shown to cause increased incidence of renal calculi in European trials.

C. Lamotrigine(LTG)

Among 711 patients receiving LTG in controlled trials the most frequent AE was dizziness(38.4% LTG, 13.4% PLC ; difference 24%). Other common AE(with percentage difference in incidence between LTG and PLC groups) were diplopia(20.1%), ataxia(16.2%), amblyopia(11%), headache(10%), nausea(9.1%), somnolence(7.3%) and rash(5.7%). Drug withdrawal rate due to AE was 6.9% in LTG group compared with 2.9% in PLC group.

In combined data of 2,601 patients receiving LTG in open and controlled trials, 235 patients(9%) withdrew LTG due to various AE. The most frequent reason for withdrawal of LTG was skin rash(2.1%). In controlled trials, rash of any type occurred in 10% of subjects but only 1.1% withdrew for this reason. The rate of rash was higher for subjects receiving concomitant VPA(2.5%) than for those receiving enzyme-inducing agents(1%). The reason for this has not been established, however, slower titration of LTG has been associated with lower incidence of skin rash suggesting a possible contribution of a dose related phenomenon.

A disproportionately high incidence of dizziness, ataxia, and diplopia were related to combining LTG with CBZ(Graves *et al.*, 1991 ; Schapel *et al.*, 1991 ; Warner *et al.*, 1992 ; Besag *et al.*, 1994 ; Lee *et al.*, 1995). These cerebellar toxicities are often eliminated by decreasing the dose of CBZ or offsetting the dosing of two AEDs by as little as 1 hour(Messenheimer *et al.*, 1994). Whether these AE are related to an increased level of CBZ-epoxide is unclear due to conflicting results. However, a pharmacodynamic rather than pharmacokinetic interaction seems more likely because similar toxicities also occurred in a patient taking OXC which does not produce CBZ-epoxide(Aving, 1994).

In a controlled monotherapy trial of LTG(n=131) compared with CBZ(n=129)(Brodie *et al.*, 1995), more patients taking CBZ withdrew due to AE, 27% compared to 15% on LTG. The commonest AE resulting in withdrawal with either drug was rash(LTG 9%, CBZ 13%). Somnolence was more common in the CBZ group(LTG 12%, CBZ 22%), $P<0.05$. In another monotherapy trial which compared LTG(n=86) with PHT(n=95)(Steiner *et al.*, 1994), more patients on PHT withdrew from the study due to AE(LTG 15%, PHT 19%). Significantly more patients taking PHT complained of asthenia(LTG 16%, PHT 29%), somnolence(LTG 7%, PHT 28%) and ataxia (LTG 0%, PHT 12%). However, skin rash was more common in the LTG group and was the reason for

withdrawal in 10 patients taking LTG and 5 patients on PHT.

These comparative trials indicate that LTG is more easily tolerable than conventional AEDs with significantly less CNS side effects. Skin rash seems to be a problem however, slower titration of LTG has been shown to decrease its incidence significantly. LTG has not been shown to cause any significant abnormalities on laboratory measures.

D. Felbamate

FBM was generally well tolerated in both poly-and monotherapy with only 7% of patients receiving FBM in controlled trials had to stop treatment because of AE.

AE are more common in polytherapy than monotherapy. The most common AE of FBM are GI and nonspecific CNS symptoms such as insomnia, anorexia, nausea, dizziness, fatigue, ataxia, lethargy, and weight loss. In the DBPC-crossover add-on trial by Leppik *et al.*(1991), 3 of 59 patients(5.1%) discontinued FBM because AE and the common AE(with differences in the incidence between FBM and PLC phases) were diplopia (34%), headache(32%), dizziness(32%), ataxia(31%), nausea(32%), vomiting(22%) and blurred vision(17%).

The pooled data of Carter Wallace showed that 120 of 977 adults(12%) in clinical trials discontinued FBM because of AE. These were anorexia(1.2%), nausea (1.4%), skin rash(1.2%), weight loss(1.1%), and various neurological symptoms(1.4%) or psychological symptoms(2.2%). In studies with children, 22 of 357 patients (6%) discontinued FBM due to AE similar to those seen in adults. In a postmarketing study by Wolff *et al.*(1994), FBM was discontinued in 24 of 132 patients(18%) with chronic refractory epilepsy because of AE with GI symptoms being most common.

In preclinical and clinical studies, FBM did not show any significant hepatic or hematologic abnormalities. However, since its first marketing in September 1993, 33 cases of aplastic anemia have been reported with its incidence being approximately 1 in 2,000 persons treated per year. Development of severe hepatitis was also reported in 14 patients. On August, 1994, FDA recommended the suspension of FBM. Clinical trials of FBM has not been conducted in Korea.

E. Gabapentin(GBP)

Overall 81.4% of patients who took GBP reported at least one AE compared with 56.7% in patients taking PLC. The most common AE reported were non-specific CNS symptoms.

In the US GBP trial(1993), somnolence(24.5% GBP, 12.2% PLC), dizziness(23.1% GBP, 9.2% PLC) and ataxia(20.2% GBP, 11.2% PLC) were the only AE higher in GBP group compared with PLC group at the cut-off level of 5%. In addition the incidence of AE was not different between patients taking a low and high dose of GBP.

In controlled trials, 7% of patients receiving GBP were withdrawn due to AE compared to 3% in those taking PLC. In a long-term extension study of US GBP trial, which followed 244 patients taking 600 to 2,400mg GBP for up to 2 years, 30% of the patients had withdrawn due to lack of efficacy and 4% due to AE. The effects of GBP on various laboratory measures have not been significant.

F. Oxcarbazepine(OXC)

In the Scandinavian multicenter trial in patients with newly diagnosed epilepsy(CBZ : 98 patients, OXC : 92 patients), the number of patients with AE(74% CBZ : 68% in OXC) and the mean number of AE per patient(3.5 in CBZ : 2.8 in OXC) tended to be lower with OXC treatment. Thirty eight patients experienced severe AE, mostly skin rashes, which lead to the discontinuation of the drug in 25 patients on CBZ and 13 patients on OXC($P=0.04$). Therefore, the potential of causing allergic skin reactions seems to be lower with OXC than CBZ. The Danish Oxcarbazepine Study Group reported that 14 of 51 patients(27%) developed skin rash to CBZ cross-reacted to OXC.

OXC may increase the incidence of hyponatremia. In a cross-sectional study by Nielsen et al.(1988), 21 of 41 patients on OXC were hyponatremic(serum sodium levels < 135 m mol/liter). Patients treated with doses higher than 30mg/kg/day had a significantly higher risk of becoming hyponatremic.

G. Topiramate(TPM)

In randomized trials, a total of 14% of patients receiving TPM withdrew-prematurely due to an AE compared to 3% for PLC. Most withdrawals were due to CNS related AE(Reife ; 1994). The most common AE of TPM compared to PLC(percentage difference between TPM 600mg of and PLC) were thinking abnormal(27%), dizziness(19%), fatigue(16%), confusion(13%), impaired concentration(11%), agitation(11%), headache(10%) and speech disturbances(10%). During the clinical trial, mean decrease in body weight have been observed and its incidence coded in the data on file was 11%. The weight decrease was greater in patients with

greater baseline weight and it seemed partially reversible during an extended period.

The incidence of AE was higher in patients taking a higher dose of TPM(≥ 600 mg/day) and occurred within the first two months of drug introduction coinciding with the drug titration period of the trials. Therefore some of AE seen in the higher dose patient groups may be related to the rapid titration. A controlled trial of TPM add-on therapy using slower titration of TPM is at the beginning stage in Korea.

There has been no evidence of clinically significant systemic toxicity including laboratory measures, however, a total of 32 patients out of more than 2,000 patients treated with TPM had developed renal calculi yielding an incidence of approximately 1.5%. This was comparable with the experiences with ZNS, which is another sulfonamide related drug having a weak inhibitory action on carbonic anhydrase.

3. Analysis and Conclusion of Clinical Trials of New AEDs

As reviewed, the premarketing clinical trials of new AEDs have been quite extensive. However, they were primarily concerned with providing evidence of efficacy and safety, and were often compared with placebo. Comparisons between new AEDs have not been undertaken yet. Therefore a firm assessment of their relative clinical value is not at hand. For epilepsy caring physicians who have a much wider choice than ever before, this kind of information is not greatly helpful in choosing the best drug for their patients in daily practice.

Brodie(1994) compared AEDs in various aspects including mechanisms of action, pharmacokinetics, efficacy, side effects and drug interactions. After a thorough analysis of advantages and drawbacks of each AEDs, he ranked VPA, LTG and OXC as the highest, CBZ, VGB, and GBP in as the next, which was followed by PHT, FBM, and then PB. His analysis was certainly based on the logical considerations of available data, however, it may not be entirely practical because choosing a drug varies upon the cases given. It seems worthwhile to try to correlate the available clinical data to real situations in daily practice.

A. Agents for Add-on Therapy in Partial Epilepsy

The efficacy of FBM in controlled add-on trials in patients with partial epilepsy was found to be equivocal and its anticonvulsant efficacy was subsequently proven

Table 7. Summary of double-blind placebo-controlled parallel study as add-on therapy of AEDs in partial epilepsy

AEDs	Efficacy					Adverse Effects				
	Responder rate (%)		Free (%)		2 ^o GTC	Withdrawal rate (%)		Most common	Special A.E.	Overall grade of toxicity
	AED	PLC	AED	PLC		AED	PLC			
^a VPA	38	19	8	1.5	effective	6.0	1.4	G-I symptoms CNS symptoms	alopecia tremor	modest
^b VGB	43	19	4.5	0	less effective	9.8	2.2	Somnolence	psychosis	mild
^c ZNS	30.3	12.7	NA	NA	effective	2.8	0	dizziness anorexia	renal stone	modest
^d LTG	34 (500mg/day)	18	NA	NA	effective	6.9	2.9	dizziness	rash	mild to modest
^e GBP	26.4 (1800mg/day)	8.4	N/A	N/A	effective	7	3	somnolence	none	mild
^f TPM	47 (600mg/day)	10	N/A	N/A	effective	14	3	CNS symptoms	renal stone	modest

Efficacy data from a : Willmore et al (1996) b : French et al (1996)
 c : Schmidt et al (1993) d : Matsuo et al (1993)
 e : US multicenter trial (1993) f : Reife (1994)

by monotherapy trials. The clinical trials of OXC were mainly compared with CBZ with comparable efficacy and slightly better tolerability.

All other AEDs have been clearly demonstrated to be effective compared to PLC in controlled add-on trials. The responder rate reported in these trials varied widely. Table 7 summarized the results of DBPC-parallel add-on trials on new AEDs. The responder rate was highest with TPM. The differences in responder rate between trial drugs and PLC were 37% for TPM(600mg/day), 24% for VGB(38mg/day), 18% for GBP(1800mg/day), 17.6% for ZNS and 16% for LTG(500mg/day). Therefore, with the exception of TPM, other new AEDs seemed to have a comparable efficacy to VPA(19%). The overall toxicity of TPM and ZNS seemed modest and comparable to other conventional AEDs. GBP and VGB are clearly less toxic with LTG in between. However, the toxicity of LTG may increase substantially in patients taking CBZ concomitantly for their drug interactions.

Other characteristics to be considered for choosing an add-on drug are mechanisms of action, drug interactions, daily dosing schedules and patients characteristics such as presence of secondarily generalized tonic-clonic seizures, age, occupation, concomitant medications and economic status.

1) patients under monotherapy

a) TPM is expected to provide the best protection against both partial and secondarily GTC seizures, however, its modest degree of CNS toxicity may interfere with learning in patients of school age.

b) VGB and GBP are the safest and do not have

cognitive side effects. VGB may be better in patients with partial seizures without secondarily GTC seizures but GBP protects against both type of seizures.

c) LTG provides a reasonable efficacy against both partial and secondarily GTC seizures. It may be a better choice in patients taking VPA for their synergistic effects and the requirement of a lower dose of LTG. LTG is quite expensive and its combination with VPA may improve the cost-effectiveness. Attention should be given to slower titration of LTG. LTG may cause higher incidence of cerebellar toxicities in patients taking CBZ.

d) ZNS may be effective, however, the data does not provide any significant advantages over VPA.

2) patients under polytherapy

In patients taking two or more conventional AEDs, drug interactions, cognitive AE and different mechanisms of action may be important considerations. In general, VGB may be useful in patients with partial seizures only and GBP in patients with both partial and GTC seizures.

B. Add-on Therapy in Generalized Epilepsy(GE)

No new AEDs had controlled trials in patients with refractory GE except VGB in infantile spasm and FBM in Lennox-Gastaut syndrome. However, preliminary clinical experiences of LTG and ZNS provided some clues about their efficacies in GE.

1) idiopathic GE

Because VPA is quite effective in idiopathic GE, the number of patients requiring add-on therapy may not be

so large. In absence seizures refractory to VPA, the add-on of ESM has been effective. Clinical trials of LTG and ZNS suggested that they were broad-spectrum AEDs. In cases of idiopathic GE resistant to VPA (except absence seizure), LTG is an appropriate drug to try. ZNS may be more appropriate in myoclonic epilepsy.

2) cryptogenic or symptomatic generalized epilepsy

a) West syndrome: In patients refractory to ACTH treatment, VGB would be the choice of drug to try. If it is refractory, ZNS is the second drug to add (Yamatogi and Ohtahara, 1991).

b) Lennox-Gastaut syndrome: FBM was proven to be effective in controlled trial. However, rare but serious toxicities like aplastic anemia or severe hepatitis are a major concern. LTG was reported as quite effective in this epileptic syndrome (Timings and Richens, 1992). Especially combination therapy of LTG and VPA provided excellent efficacies in the author's experience. ZNS is also effective in this syndrome and seems worth trying in refractory cases.

c) myoclonic epilepsy: Anecdotal experiences of ZNS in Japan were quite encouraging. Henry et al. (1988) and Seino et al. (1995) reported significant efficacy of ZNS in PME, which has not been confirmed yet by other investigators. In the author's limited experiences, the effect of ZNS was somewhat modest and unpredictable. It may be worth trying ZNS when VPA and clonazepam fail.

C. Monotherapy

Controlled clinical trials of new AEDs with conven-

tional AEDs are summarized in table 8. Although no new AEDs are recommended as the initial drug of monotherapy at present, the data provided a strong possibility.

1) VGB was compared with CBZ in two studies. The results suggested slightly less efficacy but significantly better safety than CBZ. VGB may be worth trying as the drug of initial monotherapy in school aged patients with mainly partial seizures.

2) In controlled trials, LTG provided an advantage compared with CBZ and PHT; which showed comparable efficacy and better tolerability. Therefore, LTG has a lot of potential to become the initial drug in near future. The problem with LTG is its high cost.

3) In a comparative trial of ZNS with CBZ (Seino et al., 1988) and VPA (Oguni et al., 1988) ZNS was similar to conventional AEDs in both efficacy and tolerability. Less drug interaction of ZNS may be an advantage, however, further clinical trials in newly diagnosed epilepsy are required for its use as the initial drug.

4) In a controlled comparative trial with CBZ (Dam et al., 1989), OXC was comparable in efficacy and same or slightly better in tolerability. The incidence of skin rash was lower but hyponatremia was commoner with OXC. Less drug interaction of OXC is an advantage over CBZ. However, the equivalent dose of OXC is 50% higher than CBZ and OXC is more costly. It is difficult to predict the future of OXC at present and further clinical experiences are required.

At present, comparative clinical trials of GBP and ZNS are ongoing as the initial drug in newly diagnosed

Table 8. Summary of controlled comparative study of new AEDs

Study AEDs	Type of study	Efficacy		Adverse effects		Comments
		Satisfactory control (%)	Sz. free (%)	Incidence (%)	Withdrawal rate (%)	
VGB vs CBZ (Kälviäinen et al : 1995)	open, newly diagnosed VGB (n=50), CBZ (n=50)	VGB : 60 CBZ : 60	VGB : 32 CBZ : 52	VGB : 72 CBZ : 87	VGB : 0 CBZ : 27	efficacy is comparable or slightly less. tolerability is better.
ZNS vs CBZ (Seino et al:1988)	DB, mixed* ZNS (n=59), CBZ (n=64)	ZNS : 66.1 CBZ : 65.1	NA	ZNS : 51.7 CBZ : 56.9	NA	efficacy and tolerability are comparable.
ZNS vs VPA (Oguni et al:1988)	DB, mixed* ZNS (n=18), VPA (n=16)	ZNS : 50.0% VPA : 43.8%	NA	NA	NA	efficacy and tolerability are comparable.
LTG vs CBZ (Brodie et al:1995)	DB, newly diagnosed LTG (n=131), CBZ (n=129)	LTG : 65% CBZ : 51%	LTG : 39 CBZ : 38	LTG : 28 CBZ : 50	LTG : 15% CBZ : 19%	efficacy is comparable and tolerability is better.
LTG vs PHT (Steiner et al:1994)	DB, newly diagnosed LTG (n=86), PHT (n=95)	NA	LTG : 43 PHT : 36	NA	LTG : 7% PHT : 28%	efficacy is comparable or slightly better, tolerability is comparable.
OXC vs CBZ (Dam et al:1989)	DB, newly diagnosed OXC (n=83), CBZ (n=82)	OXC : 96% CBZ : 97%	OXC : 52 CBZ : 97	OXC : 68 CBZ : 60	OXC : 14.1 CBZ : 25.1 (P=0.04)	efficacy and tolerability are comparable. severe AE was less common in OXC.

* Either untreated or refractory patients

epileptic patients which will provide valuable information in assessing their clinical value relative to conventional AEDs.

4. Future Prospects of New AEDs

The overview of clinical data on new AEDs clearly suggests that currently available new AEDs are better in tolerability and safety, and their drug interactions are less compared with conventional AEDs. On the other hand, their effects on seizure control were either comparable or slightly less. In fact, the seizure free rate of randomized add-on clinical trials was less than 10%, which was certainly far from being satisfactory. Also, the postmarketing clinical surveillances disclosed the occurrence of serious AE in the case of FBM, which raised a concern for the use of new AEDs.

It is likely that future clinical trials will be focused on the comparison of new AEDs with conventional AEDs, which will be followed by comparison between survivors. At the same time, the clinical efficacy of new AEDs on specific type of seizures or epileptic syndromes will be pursued. In fact some of the new AEDs have already been shown to be quite effective in certain epileptic syndromes: VGB in infantile spasms; FBM and probably LTG and ZNS in Lennox-Gastaut syndrome, which might be related to their different mechanisms of action. With less drug interactions, better safety, and different action mechanisms of new AEDs, it is quite likely that rational polytherapy designs will be developed rapidly for the management of various types of refractory epilepsy.

At present AED therapy in epilepsy belongs to the category of symptomatic therapy. Further investigations are necessary to find whether these new AEDs and/or newcomers would exert any influences on the natural course of epilepsy or epileptogenicity.

REFERENCES

- Alving J. Case of severe acute intoxication with oxcarbazepine combined with lamotrigine. *Epilepsia* 1994; 35(suppl 7): S72.
- Anhut H, Ashman P, Feuerstein TJ, et al. Gabapentin(neurontin) as add-on therapy in patients with partial seizures: a double-blind, placebo-controlled study. *Epilepsia* 1994; 35: 795-801.
- Appelton R. The role of vigabatrin in the management of infantile epileptic syndromes. *Neurology* 1993; 43(suppl 5): S21-3.
- Ben-Menachem E. Potential antiepileptic drugs: topiramate. In: Levy RH, Mattson RH, Meldrum BS, eds. *Antiepileptic drugs*, 4th ed. New York: Raven Press Ltd, 1995; 1063-70.
- Bernardina BD, Fontana E, Vigevano F, et al. Efficacy and tolerability of vigabatrin in children with refractory partial seizures: a single-blind dose increasing study. *Epilepsia* 1995; 36: 687-91.
- Besag FMC, Subel B, Pool F, et al. Carbamazepine toxicity with lamotrigine: a pharmacokinetic or pharmacodynamic interaction? *Epilepsia* 1994; 35(suppl 7): S73.
- Binnie CD, Debets RMC, Engelsman M, et al. Double-blind crossover trial of lamotrigine as add-on therapy in intractable epilepsy. *Epilepsy Res* 1989; 4: 222-9.
- Bourgeois BFD, Leppik IE, Sackellares JC, et al. Felbamate double blind efficacy trial following presurgical monitoring. *Epilepsia* 1991; 32(suppl 3): S18.
- Brodie MJ. Lamotrigine versus other antiepileptic drugs: a star rating system is born. *Epilepsia* 1994; 35(suppl 5): S41-6.
- Brodie MJ, Richens A, UK Lamotrigine/Carbamazepine Monotherapy Trial Group. Lamotrigine versus carbamazepine: a double-blind comparative study in newly diagnosed epilepsy. *Lancet* 1995; 345: 476-9.
- Browne TR, Mattson RH, Penry JK, et al. Multicenter long-term safety and efficacy study of vigabatrin for refractory complex partial seizures: an update. *Neurology* 1991; 41: 363-4.
- Chiron C, Dulac O, Beaumont D, et al. Therapeutic trial of vigabatrin in refractory infantile spasms. *J Child Neurol* 1991; 6(suppl 2): S52-9.
- Dam M, Ekberg R, Løyning Y, et al. A double-blind study comparing oxcarbazepine and carbamazepine in patients with newly diagnosed, previously untreated epilepsy. *Epilepsy Res* 1989; 3: 70-6.
- Dam M, Østergaard LH. Other antiepileptic drug: Oxcarbazepine. In: Levy RH, Mattson RH, Meldrum BS, eds. *Antiepileptic Drugs*, 4th ed. New York: Raven Press Ltd, 1995: 987-95.
- Danish Oxcarbazepine Study Group. Oxcarbazepine in patients hypersensitive to carbamazepine. *Acta Neurol Scand* 1984; 70: 223.
- Devinsky O, Kothari M, Rubin R, et al. Felbamate for absence seizures. *Epilepsia* 1992; 33(suppl 3): S84.
- Dodrill CB, Arnett JL, Sommerville KW, Sussman NM. Evaluation of the effects of vigabatrin on cognitive abilities and quality of life in epilepsy. *Neurology* 1992; 43: 2501-7.
- Dulac O, Chiron C, Luna D, et al. Vigabatrin in childhood epilepsy. *J Child Neurol* 1991; 6: 30-7.
- Faught RE, Sachdeo RC, Remier MP, et al. Felbamate monotherapy for partial onset seizures: an active-control trial. *Neurology* 1993; 43: 688-92.
- Felbamate Study Group in Lennox-Gastaut Syndrome. Efficacy of felbamate in childhood epileptic encephalopathy (Lennox-Gastaut syndrome). *N Engl J Med* 1993; 328: 29-33.
- Fisher RS, Kurrigan JF III. Toxicity. In: Levy RH, Mattson RH, Meldrum BS, eds. *Antiepileptic Drugs*, 4th ed. New York: Raven Press Ltd, 1995; 931-40.
- French JA, Mosier M, Walker S, et al. A double-blind, placebo-controlled study of vigabatrin three g/day in patients with uncontrolled complex partial seizures. *Neurology* 1996;

- 46: 54-61.
- Grant SM, Heel RC. Vigabatrin: a review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in epilepsy and disorders of motor control. *Drugs* 1991; 41: 889-926.
- Gram L, Klostervskov P, Dam M. γ -Vinyl GABA: a double-blind placebo-controlled trial in partial epilepsy. *Ann Neurol* 1985; 17: 262-6.
- Graves NM, Ritter FJ, Wagner ML, et al. Effect of lamotrigine on carbamazepine epoxide concentrations. *Epilepsia* 1991; 32(suppl 3): S13.
- Henry TR, Leppik IE, Gumnit RJ, Jacobs M. Progressive myoclonus epilepsy treated with zonisamide. *Neurology* 1988; 38: 928-31.
- Henry TR, Sackellares JC. Zonisamide. In: Resor Jr. St, Kutt H, eds. *The Medical Treatment of Epilepsy*. New York: Dekker, 1991; 432-27.
- Houtkooper MA, Lammertsma A, Meyer JWA, et al. Oxcarbazepine(GO 47680): a possible alternative to carbamazepine? *Epilepsia* 1987; 28: 693-8.
- Hwang YM, Kim HK, Kang JK, Lee MC. Clinical efficacy of vigabatrin as add-on therapy to valproic acid in intractable partial seizures. *J Korean Neurol Assoc* 1994; 12: 666-74.
- Jawad S, Richens A, Goodwin G, Yuen AWC. Controlled trial of lamotrigine for refractory seizures. *Epilepsia* 1989; 30: 356-63.
- Kälviäinen R, Äälikä M, Saukkonen AM, et al. Vigabatrin vs carbamazepine monotherapy in patients with newly diagnosed epilepsy. *Arch Neurol* 1995; 52: 989-96.
- Lee BI, Huh K, Kim JS, et al. Open clinical trial of vigabatrin as add-on therapy in severely intractable partial seizures. *J Korean Neurol Assoc* 1993; 11: 85-98.
- Lee BI, Park SC, Park HK, et al. Lamotrigine add-on therapy in patients with complex partial seizures receiving carbamazepine monotherapy. *Epilepsia* 1995; (suppl 3): S115.
- Lee KS, Kim BS, Chung JB, et al. Efficacy and safety of vigabatrin in refractory epilepsy. *J Korean Neurol Assoc* 1993; 11: 99-104.
- Leppik IE, Dreifuss FE, Pledger GW, et al. Felbamate for partial seizures: results of a controlled clinical trial. *Neurology* 1991; 41: 1785-89.
- Leppik IE, Willmore LJ, Homan RW, et al. Efficacy and safety of zonisamide: results of a multicenter study. *Epilepsia Res* 1993; 14: 165-73.
- Livingston JH, Beaumont D, Arzimanoglou A, Alcardi J. Vigabatrin in the treatment of epilepsy in children. *Br J Clin Pharmacol* 1989; 27: 109-12.
- Loiseau P, Hardenberg J, Pestre M, et al. Double-blind, placebo-controlled study of vigabatrin (gamma-vinyl GABA) in drug resistant epilepsy. *Epilepsia* 1986; 27: 115-20.
- Loiseau P, Yuen AWC, Duche B, et al. Randomized double-blind placebo-controlled crossover add-on trial of lamotrigine in patients with treatment resistant partial seizures. *Epilepsia Res* 1990; 7: 136-45.
- Luna D, Dulac O, Pajot N, Beaumont D. Vigabatrin in the treatment of childhood epilepsies: a single-blind placebo-controlled study. *Epilepsia* 1989; 30: 430-7.
- Matsuo F, Bergen D, Faught E, et al. Placebo-controlled study of the efficacy and safety of lamotrigine in patients with partial seizures. *Neurology* 1993; 43: 2284-91.
- McGuire AM, Duncan JS, Trimble MR. Effects of vigabatrin on cognitive function and mood when used as add-on therapy in patients with intractable epilepsy. *Epilepsia* 1992; 33: 128-34.
- McKee PJW, Blacklaw J, Friel E, et al. Adjuvant vigabatrin in refractory epilepsy: a ceiling to effective dosage in individual patients? *Epilepsia* 1993; 34: 937-43.
- Messenheimer J, Ramsay RE, Willmore LJ, et al. Lamotrigine therapy for partial seizures: a multicenter, placebo-controlled, double-blind crossover trial. *Epilepsia* 1994; 35: 113-21.
- Mumford JP, Dam M. Meta-analysis of European placebo-controlled studies of vigabatrin in drug resistant epilepsy. *Br J Clin Pharmacol* 1989; 27: 101-7.
- Nielsen OA, Johannessen AC, Bardrum B. Oxcarbazepine-induced hyponatremia, a cross-sectional study. *Epilepsia Res* 1988; 2: 269-71.
- Oguni H, Hayashi K, Fukuyama Y, et al. Phase III study of a new antiepileptic AD-810, zonisamide in childhood epilepsy. *Jpn J Pediatr* 1988; 41: 439-50.
- Ono T, Yagi K, Seino M. Clinical efficacy and safety of a new antiepileptic drug, zonisamide. A multi-institutional phase three study. *Clin Psychiat* 1988; 30: 471-82.
- Panayiotopoulos CP, Ferrie CD, Knott C, Robinson RO. Interaction of Lamotrigine with sodium valproate. *Lancet* 1993; 341: 445.
- Park HK, Lee BI, Park SC, et al. Combination therapy of lamotrigine and vigabatrin in patients with intractable partial epilepsy. *Epilepsia* 1995; (suppl 3): S110.
- Pitkäinen A, Ylinen A, Mattilainen R, et al. Long-term anti-epileptic efficacy of vigabatrin in drug-refractory epilepsy in mentally retarded patients: a 5-year follow-up study. *Arch Neurol* 1993; 50: 24-9.
- Ramsay RE. Advances in the pharmacotherapy of epilepsy. *Epilepsia* 1993; 34(suppl 5): S9-16.
- Ramsay RE. Clinical efficacy and safety of gabapentin. *Neurology* 1994; 44(suppl 5): S23-30.
- Reife RA. Topiramate-a novel antiepileptic agent. Internal report of RWJPRJ, 1994.
- Remy C, Favel P, Tell G, et al. Etude en double aveugle contre placebo en permutations croisées du vigabatrin dans l'épilepsie de l'adulte résistant à la thérapeutique. *Boll Lega Ital Epil* 1986; 54/55: 241-3.
- Richens A, Yuen AWC. Overview of the clinical efficacy of lamotrigine. *Epilepsia* 1992; 32(suppl 2): S13-6.
- Rimmer EM, Richens A. Double-blind study of γ -Vinyl GABA in patients with refractory epilepsy. *Lancet* 1984; 1: 189-90.
- Ring HA, Trimble MR, Costa DC, et al. Effect of vigabatrin on striatal dopamine receptors: evidence in humans for interactions of GABA and dopamine systems. *J Neurol Neurosurg Psychiatry* 1992; 55: 758-61.
- Risner ME. Multicenter, double-blind, placebo-controlled add-on crossover study of lamotrigine in epileptic outpatients with partial seizures. *Epilepsia* 1990; 31: 619-20.
- Rothman DL, Petroff OAC, Behar KL, Mattson RH. Brain GABA levels fail to rise as vigabatrin dose is doubled. *Epilepsia* 1994; 35(suppl 8): S143.
- Sabers A, Gram L. Pharmacology of vigabatrin. *Pharmacol Toxicol* 1992; 70: 237-43.
- Sachdeo RC, Kramer LD, Rosenberg A, Sachdeo S. Felbamate monotherapy: controlled trial in patients with

- partial onset seizures. *Ann Neurol* 1992; 32: 386-92.
- Sachdeo RC, Murphy JV, Kamin M. Felbamate in juvenile myoclonic epilepsy. *Epilepsia* 1992; 33(suppl 3): S118.
- Sander JWA, Patsalos PN, Oxley JR, et al. Randomized double-blind placebo-controlled add-on trial of lamotrigine in patients with severe epilepsy. *Epilepsy Res* 1990; 6: 221-6.
- Sander JWA, Hart YM, Trimble MR, Shorvon SD. Vigabatrin and psychosis. *J Neurol Neurosurg Psychiatry* 1991; 54: 435-9.
- Schapel GJ, Beran RG, Vajda FJE, et al. Double-blind, placebo-controlled, crossover study of lamotrigine in treatment resistant partial seizures. *J Neurol Neurosurg Psychiatr* 1993; 54: 448-53.
- Schapel GJ, Dollman W, Beran RG, Dunagan FM. No effect of lamotrigine on carbamazepine and carbamazepine-epoxide concentrations. *Epilepsia* 1991; 32(suppl 1): S58.
- Schlumberger E, Chavez F, Palacios L, et al. Lamotrigine in treatment of 120 children with epilepsy. *Epilepsia* 1994; 35: 359-67.
- Schmidt D, Jacob R, Loiseau P, et al. Zonisamide for add-on treatment of refractory partial epilepsy: a European double-blind trial. *Epilepsy Res* 1993a; 15: 67-73.
- Schmidt D, Ried S, Rapp P. Add-on treatment with lamotrigine for intractable epilepsy: a placebo-controlled cross-over trial. *Epilepsia* 1993b; 34(suppl 2): S66.
- Seino M, Miyazaki H, Ito T. Zonisamide. In: Pisani F, Perucca E, Avanzini G, Richens A, eds. *New Antiepileptic Drugs* (*Epilepsy Res*: suppl 3). New York: Elsevier, 1991; 169-74.
- Seino M, Naruto S, Tsugutaka I, Miyazaki H. Other anti-epileptic drug: zonisamide. In: Levy RH, Mattson RH, Meldrum BS eds. *Antiepileptic Drugs*, 4th ed. New York: Raven Press Ltd, 1995; 1011-23.
- Seino M, Okuma T, Miyasaka M, et al. Efficacy evaluation of AD-810(zonisamide); double-blind study comparing with carbamazepine. *J Clin Exp Med* 1988; 144: 275-91.
- Sivenius J, Kälviäinen R, Ylinen A, et al. Double-blind study of gabapentin in the treatment of partial seizures. *Epilepsia* 1991; 32: 539-42.
- Sivenius MRJ, Ylinen A, Murros K, et al. Double-blind dose reduction study of vigabatrin in complex partial epilepsy. *Epilepsia* 1987; 28: 688-92.
- Smith D, Barker G, Davies G, et al. Outcomes of add-on treatment with lamotrigine in partial epilepsy. *Epilepsia* 1993; 34: 312-22.
- Steiner TJ, Silviera C, Yuen AWC. Comparison of lamotrigine (Lamictal) and phenytoin monotherapy in newly diagnosed patients. *Epilepsia* 1994; 35(suppl 7): S61.
- Stolarek I, Blacklaw J, Thompson GG, Brodie MJ. Vigabatrin and lamotrigine in refractory epilepsy. *J Neurol Neurosurg Psychiatry* 1994; 57: 921-24.
- Tanganelli P, Regestra G. Vigabatrin versus carbamazepine in newly diagnosed epileptic patients. *Epilepsia* 1994; 35(suppl 7): S64.
- Tartara A, Manni R, Galimberti CA, et al. Six-year follow-up study on the efficacy and safety of vigabatrin in patients with epilepsy. *Acta Neurol Scand* 1992; 86: 247-51.
- Tartara A, Manni R, Galimberti CA, et al. Vigabatrin in the treatment of epilepsy: a double-blind placebo-controlled study. *Epilepsia* 1986; 27: 717-723.
- Tartara A, Manni R, Galimberti CA, et al. Vigabatrin in the treatment of epilepsy: a long-term follow-up study. *J Neurol Neurosurg Psychiatry* 1989; 52: 467-71.
- Tassinari CA, Michelucci R, Ambrosetto G, Salvi F. Double-blind study of vigabatrin in the treatment of drug resistant epilepsy. *Arch Neurol* 1987; 44: 907-10.
- Theodore WH, Jensen PK, Kwan RM. Felbamate: Clinical use. In: Levy RH, Mattson RH, Meldrum BS, eds. *Antiepileptic Drugs*, 4th ed. New York: Raven Press Ltd, 1995; 817-22.
- Theodore WH, Raubertas R, Porter RJ, et al. Felbamate: a clinical trial for complex partial seizures. *Epilepsia* 1991; 32: 392-7.
- Timmings PL, Richens A. Lamotrigine as add-on drug in the management of Lennox-Gastaut syndrome. *Eur Neurol* 1992; 32: 305-7.
- UK Gabapentin Study Group. Gabapentin in partial epilepsy. *Lancet* 1990; 335: 1114-7.
- US Gabapentin Study Group No.5. Gabapentin as add-on therapy in refractory partial epilepsy: a double-blind, placebo-controlled, parallel group study. *Neurology* 1994; 43: 2292-8.
- Warner T, Patsalos PN, Prevett M, et al. Lamotrigine-induced carbamazepine toxicity: an interaction with carbamazepine-10-11-epoxide. *Epilepsy Res* 1992; 11: 147-50.
- Wilder BJ, Ramsay RE, Guterman A, et al. A double-blind multicenter Placebo-controlled study of the efficacy and safety of zonisamide in the treatment of complex partial seizures in medically refractory patients. Internal report of Daiippon Pharmaceutical Co. 1986.
- Wilensky AJ, Temkin NR, Ojemann LM, et al. Gabapentin and carbamazepine as monotherapy and combined: A pilot study. *Epilepsia* 1992; 33(suppl 3): S77.
- Willmore LJ, Shu V, Wallin B, M88-194 Study Group. Efficacy and safety of add-on divalproex sodium in the treatment of complex partial seizure. *Neurology* 1996; 46: 49-53.
- Wolff DL, Kerrick JM, Rarick J, et al. Postmarket felbamate use in the refractory epilepsy population. *Epilepsia* 1994; 35(suppl 8): S32.
- Wolf SM, Shinnar S, Kang H, et al. Gabapentin toxicity in children manifesting as behavioral changes. *Epilepsia* 1996; 36: 1203-5.
- Yagi K, Seino M. Methodological requirements for clinical trials in refractory epilepsies-our experience with zonisamide. *Prog Neuropsychopharmacol Biol Psychiatry* 1992; 16: 79-85.
- Yagi K, Seino M, Watanabe Y, Kujo T. Change of urine volume and urinary components in patients with epilepsy before and after zonisamide treatment. Internal report of Daiippon Pharmaceutical Co, 1992.
- Yamatogi Y, Ohtahara S. Current topics of treatment. In: Ohtahara S, Roger J, eds. *Proceedings of the international symposium, new trends in pediatric epileptology*. Department of Child Neurology, Okayama Medical School, 1991; 136-48.
- Yuen AWC. Safety issues. In: Richens A, ed. *Clinical update on lamotrigine: a novel antiepileptic drug*. Tunbridge Wells: Wells Medical Ltd, 1992; 69-75.