Gamma Knife Radiosurgery for Malignant Tumors

C.-Jin Whang, M.D., F.A.C.S., Yang Kwon, M.D.

Department of Neurological Surgery, Asan Medical Center University of Ulsan, Seoul, Korea

Between May 1990, and June 1994, 79 patients with malignant tumors were treated radiosurgically using a Leksell gamma unit at Asan Medical Center. Of these patients, 57 were metastatic brain tumor, 12 were glioblastoma multiforme(GM), 4 were primitive neuroectodermal tumor, 3 were malignant germ cell tumor, 2 were recurrent lymphoma, and 1 was adenoid cystic carcinoma of the orbit. Among 57 patients with metastatic tumors, 28 patients harboring 60 tumors were followed clinically and radiographically. The median marginal dose for these tumors was 30 Gy and the median survival rate was 15 months. Twenty-one tumors disappeared and 32 tumors decreased in size during 2 to 6 months after radiosurgery on computed tomographic or magnetic resonance imaging scans. All 12 patients with GM were treated with conventional radiation(6,240~6,500 cGy) after surgical resection or biopsy prior to radiosurgery(13~15 Gy to margin). The results were varied. Radiosurgical treatment of two recurrent lymphomas and three recurrent mixed germ cell tumors after radiation and chemotherapy provided rapid clinical improvement with disappearance of the tumor. However, new lesions appeared in two lymphomas and one mixed germ cell tumor within 3 to 4 months. One patient with adenoid cystic carcinoma of the orbit, who was treated radiosurgically prior to resection, is alive without recurrence 31 months after the treatment. Gamma knife radiosurgery appears to be the best alternative method to surgical excision plus radiation therapy for single and multiple cerebral metastases. It also provides rapid palliation of symptoms due to recurrent malignant tumors. And it may have an adjuvant role in the treatment of some tumors delaying local recurrence, if given prior to resection. However, the preliminary results for the malignant gliomas were inconclusive.

Key Words: Gamma knife radiosurgery, Metastatic tumors, Recurrent malignant tumor

INTRODUCTION

Radiosurgery was first developed by Leksell to treat functional diseases, and recently it has been given

rapid acceptance along with improvement of neuroimaging technique. Most of the cases that are being treated at the majority of radiosurgery centers have been intracerebral arteriovenous malformations(AVMs) and benign tumors(Leksell, 1951; Steiner et al., 1992).

The efficacy of radiosurgery on AVMs and benign tumors is well established(Steiner et al., 1992). But others rather than these AVMs and benign tumors,

---

Address for correspondence: Choong-Jin Whang, M.D., Department of Neurological Surgery, Asan Medical Center University of Ulsan, 388-1, Poongnap-dong, Songpa-gu Seoul, 138-040, Korea. Tel: 82-2-224-3550.

metastatic tumors and primary malignant tumors are more common, as 50 % of all intracranial tumors are metastatic tumors (Young and Patchell, 1990; Kihlstrom et al., 1992; Cairncross and Posner, 1993). Metastatic brain tumors are a manifestation of advanced cancer and if not treated, the median survival will only be one month and even despite of getting optimal medical care, life expectancy won't be more than a few months (Leksell, 1951; Richards and McKissock, 1963; Deeley and Edwards, 1968; Order et al., 1968; Markesbery et al., 1978; Dosoretz et al., 1980; Galicich et al., 1980; Hendrickson et al., 1986; Mandell et al., 1986; Smalley et al., 1987; De Angelis et al., 1989; Coffey et al., 1992; Kihlstrom et al., 1992; Smalley et al., 1992; Steiner et al., 1992).

Today, surgical removal and radiation therapy offers the best results, as a standard treatment (Patchell et al., 1986; Patchell et al., 1990). But surgery has its own risks and the patient must be in good condition to have an open craniotomy under general anesthesia and it is limited to patients with single metastasis. Approximately fifty percent of patients have a single metastasis, and overall eighty-three percent of patients have 3 or less lesions (Delattre et al., 1988). The correlation between the number of metastasis and the systemic spread of malignant diseases are not clear (Delattre et al., 1988). So there should be other treatment method that are safer, simpler, easier, and have better or at least the same results. This would be beneficial to more patients.

Primary malignant tumors, as the prognoses are poor, even after a biopsy or open surgical removal and fractionated radiation therapy with other adjuvant therapy will still persist or will have a local recurrence in many patients. In these cases, there are reports on what role radiosurgery can play as one of the modalities in adjuvant treatment, but it doesn't seem so conclusive (Gutin and Wilson, 1990; Loeffler et al., 1990 a; Coffey et al., 1992; Dempsey et al., 1992; Loeffler et al., 1992 a,b; Steiner et al., 1992 a).

This study presents the cases who were treated with the Leksell gamma knife during the initial 4 years of its operation at our hospital in order to evaluate gamma knife radiosurgery for its role as an alternative to the primary treatment of metastatic tumors and its role as an adjuvant treatment modality to primary malignant tumors.

METHODS AND MATERIALS

Seventy-nine patients with primary and secondary malignant brain tumors were treated with a gamma knife at Asan Medical Center, from May 1990 through June 1994.

Metastatic tumors were the most common having 57 cases, followed by glioblastoma multiforme (GM) 12 cases, primitive neuroectodermal tumor (PNET) 4 cases, malignant germ cell tumor (GCT) 3 cases, recurrent lymphoma 2 cases and 1 case of adenoid cystic carcinoma involving the orbit (Table 1).

Of 57 patients with metastatic tumor, 28 were available for follow-up both clinically and radiologically. There were 18 males and 10 females. Their age ranged from 28 to 77 years with the mean age being 51.6 years old. These 28 patients had 60 metastatic lesions, 16 having single metastasis, 6 having 2 metastases, 3 having 3 metastases and 3 having over 4 metastases (Table 2). The primary tumor sites were the lung in 15 patients, gastrointestinal tract in 2, breast in 2, endometrium in 1, prostate in 1, kidney in 1, malignant melanoma in 1 and 4 unknown (Table 3). The volume of the tumor was 224 to 68,585 mm³ and

Table 1. Malignant tumors treated between May 1990 and June 1994.

	No. of patients
Metastatic tumors	57
Glioblastoma Multiforme	12
PNET*	4
Recurrent Lymphoma	2
Malignant Germ Cell Tumors	3
Adenoid Cystic Carcinoma, orbit	1
Total	79

*PNET : Primitive neuroectodermal tumor.

Table 2. Characteristics of 28 patients with metastatic tumors who were followed clinically and radiologically.

	Number
• Male : Female	18 : 10
• Age (years)	
Mean	51.6
Range	28~77
• Single Metastasis	16
Two Metastasis	6
Three Metastasis	3
Over 4 Metastasis	3
Total	60

the median value was 12,550 mm³. Marginal dose was 10 to 35 Gy with median dose being 30 Gy. The isodose line used was 50 % in most cases (Table 4).

Of 12 patients with GM, 3 were stereotactically biopsied, while 9 were subtotally resected (Table 5). All had radiation therapy before gamma knife. The treatment volume ranged from 3591 to 23,431 mm³ and marginal dose ranged from 13 to 20 Gy (Table 6).

Three of four patients with PNET went through radiosurgery, after subtotal resection with receiving radiation therapy to the residual tumor, and 15-25 Gy was given to the tumor margin.

We treated 2 patients with recurrent lymphoma. Both received radiation therapy and chemotherapy 3 to 4 years prior to recurrence. They were given 15 and 29 Gy respectively for palliation.

There were 3 patients with malignant GCTs (Table 7). One of them had chemotherapy after radiation therapy and gross total excision, followed by chemotherapy, however the tumor recurred in the brain stem, and progressive right hemiparesis and aphasia developed. As for palliative treatment, 15 Gy of marginal dose was given. The other had embryonal cell carcinoma in the pineal gland and was receiving radiation therapy up to 1,620 cGy when it had to be interrupted because of rapid deterioration of the patient's neurological status. So gamma knife radiosurgery was performed along with chemotherapy. A peripheral dose of 15 Gy was given.

The patient with the adenoid cystic carcinoma involving the orbit underwent gamma knife radiosurgery before surgical resection in order to prevent recurrence. As the size was larger, we used 33 % of the isodose curve, giving 33 Gy to the margin.

In all 79 patients, along with mild sedation and local anesthesia, Leksell stereotactic head frame (ELECTA Instruments) application and radiosurgical treatment was performed. We used high resolution CT scan or gadolinium enhanced MRI to define the tumor volume and to calculate target coordinates. Dose planning was done by Micro VAX computer (Digital Equipment Corp. Westminister, MA). Most of the patients were discharged the day after the treatment.

RESULTS

Twenty-eight patients had 60 metastatic tumors. Follow-up with CT or MRI to evaluate short term

Table 3. Primary site of metastatic tumors.

Primary Site	Number
Lung	15
G.I.*	2
Kidney	2
Breast	2
Melanoma	1
Endometrium	1
Prostate	1
Unknown	4
Total	28

*G.I.; Gastrointestinal Tract.

Table 4. Radiosurgical dosimetry for metastatic tumors.

• Volume (mm ³)	
Median	5750
Range	224-68585
• Marginal Dose (Gy)	
Median	30
Range	10-35
• Isodose at Margin (%)	
Median	50
Range	30-90

Table 5. Characteristics of 12 patients with glioblastoma multiforme (GM) treated with Gamma Knife.

Male : Female	7 : 5
Ages	43(4-58)
Pre G/K treatment (N=12)	
WBRT	5940-6500Gy
Stereotactic biopsy	3
Partial resection	9

Table 6. Radiosurgical dosimetry for glioblastoma multiforme (GM) (N=10).

Volume (mm ³)	
Median	13,083
Range	3,591-23,431
Marginal dose (Gy)	
Median	12.75
Range	10-25
Isodose at Margin (%)	
Median	50
Range	40-60

Table 7. Summary of patients with malignant germ cell tumors.

Sex/Age	Dx	Location	RT (cGy)	Marginal Dose(Gy)	Follow-up
M/19	Recurrent Mixed GCT	Brain Stem	5,300	15	disappearance of tumor in 3 weeks, but dissemination in 3 months
M/15	Embryonal cell ca.	Pineal	1,620	15	disappeared in 2 months, still alive(28 months)
M/19	Choriocarcinoma	Pineal		20	decreased in 3 months, disappeared in 12 months, still alive (17 months)

response to gamma knife radiosurgery on metastatic tumor was done after 2 to 6 months of treatment (Table 8). Of 60 tumors, 21(35 %) have completely subsided(Fig. 1), 32(53.3 %) have decreased in size, 1 showed no change and 6 had become a little larger. The ones that were not eradicated, but decreased or not changed in size, all showed some improvement with decrease of peritumoral edema. The ones whose size became larger showed an initial decrease in two month but increased thereafter.

During the follow-up period(median 12 months, ranging from 5 to 39 months), among the 60 tumors, 54(90 %) were controlled locally and of 28 patients, 11 developed recurrences from 5 to 17 months(Table 8). The median survival time by Kaplan-Meier method was 15 months(Fig. 2).

Table 8. Follow-up results of patients with metastatic tumors after gamma knife radiosurgery(n=60).

	Number	Percent
disappeared	21	35
decreased	32	53.3
no change	1	12
local recurrence	6	10
Total	60	100
Tumor control rate	54/60	90

Of 10 patients with GM, follow-up(median 10 months, ranging from 3 to 26 months) imaging studies showed 1 with central low density(Fig. 3), 4 with decreased tumor size, 2 with increased edema, 2 with increased tumor size, 1 with no change. The

Fig. 1. Left, Preoperative T1-weighted MRI with gadolinium enhancement showing a metastatic tumor from lung cancer. Right, CT scan with contrast enhancement 3 months after radiosurgery showing disappearance of the lesion.

Fig. 2. Kaplan-Meier analysis of survival data of 28 patients with metastatic brain tumor. Median survival, 15 months.

median survival time by Kaplan-Meier method was 12 months (Fig. 4).

One patient with PNET developed central low density (Fig. 5), but expired 17 months after treatment with progressive hydrocephalus (the parents declined to have any further treatment at this point). Three PNET patients had increased in size 4 to 7 months after radiosurgery.

In both cases of recurrent lymphoma, the tumor rapidly disappeared after gamma knife radiosurgery but recurred at other sites in 3 to 4 months.

One patient with mixed GCT which recurred in the brain stem, had no tumor appearance on CT scan

Fig. 3. Left, Stereotactic CT scan with contrast enhancement during the radiosurgical procedure of a 52 year old man with glioblastoma multiforme. Right, contrast-enhanced CT scan 3 months after Gamma Knife surgery showing central low density within the tumor.

Fig. 4. Kaplan-Meier analysis of survival data of 10 patients with glioblastoma multiforme (GM). Median survival 12 months.

and showed improvement in clinical symptoms 3 weeks after radiosurgery, but disseminated through CSF pathway in 3 months after treatment (Table 6, Fig. 6). In one patient with embryonal cell carcinoma in the pineal gland, the tumor resolved after 2 month of gamma knife treatment and the patient is doing well after 1 year of radiosurgery (Table 7). In one patient with choriocarcinoma in the pineal gland, the tumor decreased 3 months after radiosurgery and resolved in 12 months with chemotherapy.

The patient with adenoid cystic carcinoma of the orbit was alive without recurrence 31 months after gamma knife radiosurgery and surgical resection.

Fig. 5. Left, contrast-enhanced CT scan of 3 year old girl with PNET. Right, contrast-enhanced CT scan 5 months after radiosurgery showing loss of contrast-enhancement within the tumor.

Fig. 6. Left, T1-weighted MRI with gadolinium enhancement of a patient with recurrent mixed germ cell tumor. Right, contrast-enhanced CT scan 3 weeks after Gamma knife surgery showing disappearance of the tumor.

DISCUSSION

As almost all of the metastatic brain tumors are well defined radiographically and are relatively round, these are ideal lesions for microsurgical resection or radiosurgery (Kelley et al., 1988). And as most of the lesions are relatively small in size (<3.0 cm) at the time of presentation, and are minimally invasive, the entire extent of the tumor may well be included in the

radiosurgery field (Smalley et al., 1987; Loeffler et al., 1990 b; Loeffler et al., 1992 b).

Kihlstrom et al. (1992) treated 86 patients with metastatic tumor from 1975 through 1991, and among them, analyzed 33 patients who harbored 52 tumors. The minimal dose for the tumor was 13–50 Gy and the average was 29 Gy. Of 52 tumors, 50 showed a response, and among these, 23 were completely eradicated, 25 decreased in size, 2 showed no

change in size, and 1 increased in size with radiation necrosis within the tumor. Among the 23 completely eradicated tumors, 21 disappeared 3 months after gamma knife radiosurgery and 1 after 6 months of treatment. In most of the patients who had edema around the tumor at the time of presentation, the edema subsided with treatment. It could be said that the results are almost the same as those of our short term response. There are no differences in responses by histology of the tumor.

Coffey et al.(1992) did whole brain radiation with 30 Gy to metastatic tumors and to 2.5 cm size tumors and gave a boost radiosurgery with 16 Gy at the periphery of the tumor with a gamma knife. Among 31 patients treated, 23 patients were followed-up for a period of 1 to 24 months. Among these, 13 tumors disappeared or decreased in size, 7 showed no change, and 3 increased in size.

The local control rate of radiosurgery with gamma knife or linear accelerator to metastatic tumors varies from 81 % to 96 % according to reports from different centers(Adler et al., 1992 ; Coffey et al., 1992 ; Kihlstrom et al., 1992 ; Loeffler et al., 1992 b).

Sturm et al.(1987) treated 12 cases of deep seated solitary metastasis using a linear accelerator, and even though they were radioresistant, reported that in all 7 patients who had 20 to 30 Gy, local control was possible for more than 3 months of follow-up.

Loeffler et al.(1992 b) reported that they treated 111 patients who had 155 recurrent metastatic tumors with a linear accelerator for 5 years. The report showed 8 % of local recurrence(12/155), and 31 % of distant metastases(39/111).

Kihlstrom et al.(1992) treated 33 patients with metastases using a gamma knife. Among 25 patients who only had radiosurgery, 6 patients(24 %) developed distant metastases.

Patchell et al.(1990) conducted a randomized study recently on radiation therapy alone, and surgery with postoperative radiation therapy on single brain metastatic patients. Local recurrence occurred in 20 % in the group with surgery and postoperative radiation therapy, 52 % in the group with radiation therapy alone, and the median survival was 40 weeks and 15 weeks respectively.

Surgical resection followed by radiation therapy has some limitations : it has to be a single lesion and the patients should be able to undergo craniotomy under general anesthesia, and it has a high local recurrence rate. On the other hand, radiosurgery has no risk in

surgical complications such as hemorrhage, infection; wound problems, etc, and it can be used in multiple lesions and in patients in poor general condition. Radiosurgery offers a short hospitalization and a high local control rate.

Alexander et al.(1991) performed radiosurgery on 13 patients with GM and 4 patients with astrocytoma using a linear accelerator. They used radiosurgery as a primary therapy, which was followed by radiation therapy. The peripheral dose of radiosurgery was 10 to 20 Gy, and the average radiation therapy was 59.6 Gy. Among the 13 patients followed-up for 2 to 17 months, 10 patients survived up to 10 months after radiosurgery without clinical and radiographic progression.

Coffey et al.(1992) performed boost radiosurgery with a gamma knife after radiation therapy on 6 patients with GM, 4 patients with anaplastic astrocytoma, and 5 patients with ependymoma. For a period of 1 to 24 months of follow-up with CT or MRI among 15 patients, in 11(73 %) patients, tumors were under control.

Loeffler et al.(1992 a) treated 23 patients with GM and 14 patients with anaplastic astrocytoma. All of these patients went through external-beam radiotherapy of 5,490 cGy after surgical resection or biopsy and had radiosurgery 2 to 4 weeks later. The median tumor volume was 4.800 mm³ and the minimum dose was 1,000-2,000 cGy. They followed-up 37 patients for 7 to 41 months, and among these patients, the median survival was 26 months for GM patients.

At the Karolinska Institute, they used radiosurgery as an adjunctive treatment to operation or radiation therapy on 15 patients with anaplastic astrocytoma and GM. Even in high doses, not much efficacy could be seen and it was concluded that there should be more work done on this subject(Steiner et al., 1992).

The median survival of primary central nervous system lymphoma undergoing radiation therapy is not quite 1 year, and even if chemotherapy is added, the tumor will eventually recur, and the overall median survival time by Hochberg et al.(1988) was 13.5 months. In our study 2 patients with lymphoma, which recurred after radiation therapy and chemotherapy, had gamma knife radiosurgery. The tumors rapidly disappeared but recurred at other sites 3 to 4 months later.

The present standard treatment for pure germinoma is radiation therapy and in malignant nongerminomatous germ cell tumor, chemotherapy should be

added (Hoffman et al., 1991). Casentini et al. (1990) did radiosurgery with a dose of 10 to 20 Gy in 7 patients with germinoma and followed by standard external radiotherapy (24–36 Gy) to prevent tumor cell seeding. The study showed good results with small size tumors which regressed within days after radiosurgery and were completely eradicated within months. One of our patients who had recurrent mixed GCT had a recurrence after radiation therapy and surgical excision. Chemotherapy was added without any success to control the tumor. The tumor recurred in the brain stem, but disappeared in 3 weeks after gamma knife radiosurgery. After 3 months of radiosurgery, disseminated tumors were found. In one patient with embryonal cell carcinoma in the pineal gland, we had to terminate radiation therapy before completion because of a rapid deterioration of the patient's neurological condition. Instead, the patient had radiosurgery and continued chemotherapy, and has done very well without any recurrence for over two years of follow-up. In one patient with choriocarcinoma in the pineal gland, the tumor decreased in size 3 months after radiosurgery and resolved in 12 months with chemotherapy.

One patient with adenoid cystic carcinoma involving the orbit and temporal lobe had radiosurgery before surgical excision. Radiosurgery was done in order to prevent local recurrence after excision.

Gamma knife surgery is the best alternative method to surgical excision plus radiation therapy in treating metastatic brain tumor. In treatment of recurrent malignant tumors, gamma knife radiosurgery offered rapid symptomatic palliation in selected cases. In malignant nongerminomatous GCT, it has a role to play as an adjunctive treatment with chemotherapy.

This study concluded that there was a considerable decrease in local recurrence rate by performing radiosurgery before planned surgical excision in selected patients. However, the study was unable to draw any conclusions for malignant gliomas.

REFERENCES

- Alexander III E, Kooy HM, Nedzi L, Gelman RS, Loeffler JS. Linear accelerator radiosurgery for metastatic tumors. In : Alexander III E et al, eds. *Stereotactic and functional neurosurgery*, in press.
- Adler JR, Cox RS, Kaplan I, Martin DP. Stereotactic radiosurgical treatment of brain metastases. *J Neurosurg* 1992; 76: 444–9.
- Cairncross JG, Posner JB. The management of brain metastasis. In : Walker MD, ed. *Oncology of the nervous system*. Boston : Martinus Nijhoff, 1993; 341–77.
- Casentini L, Colombo F, Pozza F, Beneditti A. Combined radiosurgery and external radiotherapy of intracranial germinomas. *Surg Neurol* 1990; 34: 79–86.
- Coffey RJ, Lunsford LD, Flickinger JC. The role of radiosurgery in the management of malignant brain tumors. *Neurosurg Clin North Am* 1992; 3: 231–44.
- DeAngelis LM, Mandell LR, Thaler HT, Kimmel DW, Galicich JH, Fuks Z, Posner JB. The role of postoperative radiotherapy after resection of single brain metastases. *Neurosurgery* 1989; 24: 798–805.
- Delattre JY, Krol G, Thaler HT. Distribution of brain metastases. *Arch Neurol* 1988; 45: 741–4.
- Dempsey DK, Lunsford LD. Stereotactic radiosurgery for pineal region tumors. In : Lunsford LD, ed. *Stereotactic radiosurgery update*. Elsevier, 1992; 245–54.
- Dosoretz DE, Blitzer PH, Russell AH. Management of solitary metastasis to the brain: The role of elective brain irradiation following complete surgical resection. *Int J Radiat Oncol Biol Phys*. 1980; 6: 1727–30.
- Galicich JH, Sundaresan N, Arbit E, Passet S. Surgical treatment of single brain metastasis: Factors associated with survival. *Cancer* 1980; 45: 381–6.
- Gutin PH, Wilson DB. Radiosurgery for malignant brain tumors. *J Clin Oncol* 1990; 8: 571–3.
- Hendrickson FR, Lee MS, Larson M. The influence of surgery and radiation therapy on patients with brain metastasis. *Int J Rad Oncol Biol Phys* 1983; 9: 623–7.
- Hochberg FM, Miller DC. Primary central nervous system lymphoma. *J Neurosurg* 1988; 68: 835–53.
- Hoffman HJ, Otsubo H, Hendrick B, Humphreys RP, Drake JM, Becker LE, Greenberg M, Jenkin D. Intracranial germ-cell tumors in children. *J Neurosurg* 1991; 74: 545–51.
- Kelley PF, Kall BA, Goerss SJ. Results of computed tomography-based computer-assisted stereotactic resection of metastatic intracranial tumors. *Neurosurgery* 1988; 22: 7–17.
- Kihlstrom L, Karlsson B, Lindquist C. Gamma knife surgery in brain metastasis. In : Lunsford LD, ed. *Stereotactic Radiosurgery Update*, Elsevier, 1992; 429–34.
- Leksell L. The stereotaxic method and radiosurgery of brain. *Acta Chir Scand* 1951; 99: 316–9.
- Loeffler JS, Alexander III E, Hochberg FH. Clinical patterns of failure following stereotactic interstitial irradiation for malignant gliomas. *Int J Radiat Oncol Biol Phys* 1990; 19: 1455–62. a
- Loeffler JS, Alexander III E, Shea WM. Radiosurgery as part of the initial management of patients with malignant glioma. *J Clin Oncol* 1992; 9: 1379–85. a
- Loeffler JS, Alexander III E, Wen PY. Radiosurgery for brain metastasis five year experience at the Brigham and women's hospital. In : Lunsford LD, ed. *Stereotactic Radiosurgery Update*, Elsevier, 1992; 383–92. b.
- Loeffler JS, Kooy HM, Wen PY. The treatment of recurrent brain metastasis with stereotactic radiosurgery. *J Clin*

- Oncol* 1990; 8: 576-82 b.
- Mandell L, Hilaris B, Sullivan M. *The treatment of single brain metastasis from non-oat cell lung carcinoma. Cancer* 1986; 58: 641-9.
- Markesbery WR, Brooks WH, Gupta GD. *Treatment for patients with cerebral metastasis. Arch Neurol* 1978; 35: 754-56.
- Order SE, Hellman S, Essen CV. *Improvement in quality of survival following whole-brain irradiation for brain metastasis. Radiology* 1968; 91: 149-53.
- Patchell RA, Cirincione C, Thaler HT. *Single brain metastasis: Surgery plus radiation or radiation alone. Neurology* 1986; 36: 447-53.
- Patchell RA, Tibbs PA, Walsh JW. *A randomized trial of surgery in the treatment of single metastasis to the brain. N Engl J Med* 1990; 22: 494-500.
- Richards P, McKissock W. *Intracranial metastasis. Br Med J* 1963; 5: 15-8.
- Smalley SR, Laws ER, O'Fallon JR, Shaw EG, Schray MF. *Resection for solitary brain metastasis. J Neurosurg* 1992; 77: 531-40.
- Smalley SR, Schray MF, Laws ER, O'Fallon JR. *Adjuvant radiation therapy after surgical resection of solitary brain metastasis: Association with pattern of failure and survival. Int J Radiat Oncol Biol Phys* 1987; 13: 1611-6.
- Steiner L, Lindquist C, Stine M. *Radiosurgery. Adv Tech Stand in Neurosurg* 1992; 19: 19-95.
- Sturm V, Kober B, Hover KH, Schlegel W. *Stereotactic percutaneous single dose irradiation of brain metastasis with a linear accelerator. Int J Radiat Oncol Biol Phys* 1987; 13: 279-82.
- Young B, Patchell RA. *Surgery for a single brain metastasis. In: Wilkins RH, Rengachery SR, eds. Neurosurgery update I. New York: McGraw-Hill Inc, 1990; 473-6.*