Endoscopic Mucosal Resection

Chan-Sup Shim, M.D., Ph.D

*Department of Internal Medicine • Institute for Digestive Research
Soon Chun Hyang University College of Medicine, Seoul, Korea*

INTRODUCTION

With the progress in endoscopic equipments and techniques, endoscopic diagnosis and therapy of gastrointestinal diseases have advanced rapidly and dramatically over the past 15 years. Endoscopic mucosal resection (EMR), which is a recently introduced therapeutic method for resection of flat or depressed lesions of gastrointestinal tract, has come to play an increasingly important role in treatment of early cancer in gastrointestinal tract. EMR was developed for the purpose of obtaining a larger biopsy specimen, but is now used mainly radical endoscopic treatment of small, flat and depressed types of early cancer in gastrointestinal tracts. Full-thickness large specimen obtained by EMR provides the definite advantage of contributing to accurate histopathological analysis compared with other endoscopic treatments such as laser therapy and injection debulking therapy. In this review, the development of methods, complications, indications of endoscopic mucosal resection and the efficacy of endoscopic mucosal resection for curative treatment of early cancer in the gastrointestinal tracts and recent advances in the area are summarized.

DEVELOPMENT OF ENDOSCOPIC MUCOSAL RESECTION

Since the fiberscope with a working channel was devised by Machida, Japan, in 1964, the endoscopic forceps biopsy has been widely used to establish the diagnosis of gastrointestinal lesion. However, the specimens obtained by means of a forceps biopsy are too small to make an accurate diagnosis in some cases. For this reason, several methods to get larger specimens, for example, snare biopsy and jumbo biopsy etc., have been tried (Ottenjann et al., 1973; Tada et al., 1980). Ottenjann et al. (1973) devised a big particle biopsy method (snare biopsy) by cutting the elevated lesion by use of electrocoagulation. Tada et al. (1980)

developed a method of 'jumbo biopsy', which was a suction biopsy under direct endoscopic vision. But these techniques have some troublesome to be applied to the depressed lesions. To overcome this problem, Tada et al. (1984) devised a method, the so-called 'strip biopsy', which made it possible to obtain larger specimens (approximately 3 cm in diameter) even from flat or depressed lesions. And then they reported successful results of the strip biopsy for the treatment of early gastric cancer as well as for diagnosis of gastric borderline lesions. Thereafter, some authors have devised several methods, which have been based on the principles of the strip biopsy and made it possible to obtain larger specimens from the flat or depressed lesion without any serious complications (Hirao et al., 1988; Kawano et al., 1991; Makuuchi et al., 1991; Inoue et al., 1993; Lee et al., 1993; Fleischer et al., 1996; Lee et al., 1996). These all endoscopic techniques have been called as 'endoscopic mucosal resection', separately with endoscopic polypectomy, since they made it possible to resect the mucosa of flat or depressed lesions which cannot be resected by the conventional snare polypectomy technique.

METHODS OF ENDOSCOPIC MUCOSAL RESECTION

Although there are several different techniques for EMR, the principles of all techniques are based on the combination of endoscopic injection technique and snare polypectomy technique using the high frequency electric current.

1. Strip biopsy (Strip-off biopsy)

Strip biopsy, which had been devised by Tada et al. (1984) is a basic technique of endoscopic mucosal resection. For this procedure, the double-channel endoscope, the needle injection catheter, grasping forceps, snare catheter and electrosurgical unit should be prepared. The procedure are as follows: Under fiber-

endoscopic control, needle injection catheter is located near the target lesion through the endoscopic working channel. Two or four ml of physiological saline or saline-epinephrine solution are injected into the submucosa of the lesion through the needle injection catheter until the target lesion is elevated to form Yamada type II-III shape. And then this elevated target lesion is lifted by a grasping forceps inserted through the one working channel, and the snare inserted through the other working channel is tightened around the lesion. After confirming the snaring is correct position, this target lesion is removed by electrocoagulation. The resected tissue is retrieved by grasping forceps with endoscope. Fig. 1 and Fig. 2 are the schema and the endoscopic views representing the procedure.

In this procedure, the injection of physiological saline can provide some advantages with resect to following three points: 1) it makes easy the snare to entrap the flat or depressed lesion due to elevation of the target lesion, 2) it prevents the electrical damage to the deeper layer by maintaining the distance between the mucosal layer and the proper muscle layer, and 3) it prevents the bleeding due to the tamponade effect of saline or the vasoconstriction effect of the epinephrine contained in saline-epinephrine solution.

2. Endoscopic Resection with Local Injection of Hypertonic Saline-Epinephrine (ERHSE)

ERHSE, which has been devised by Hirao et al.

(1988) follows the similar principles as strip biopsy. But in ERHSE, after marking around the target lesion, 15 to 40 ml of a hypertonic saline-epinephrine solution (40 ml of 50% hypertonic glucose solution + 1 ml of 0.1% epinephrine + indigocarmine) are injected, instead of the saline solution used in strip biopsy. And then, the mucosa surrounding the target lesion is cut with a needle knife using electrical current along with marking and snared for removal of the lesion. Fig. 3 is the schema showing the procedures.

ERHSE permits the resection of larger lesion, but has higher risk of perforation by needle knife than in strip biopsy.

3. Endoscopic Mucosal Resection using Intraluminal Negative Pressure

In some techniques of EMR, intraluminal negative pressure induced by the endoscopic suction is used for lifting the target lesion, instead of grasping forceps used in strip biopsy. In these techniques, the double channel endoscope is not necessary to resect the lesion, in contrast to the strip biopsy and ERHSE techniques.

1) Endoscopic Mucosal Resection using EEMR Tube (EMRT)

EMRT, which has been developed by Makuuchi et al.(1991) is a technique that enables to resect the lesion by use of EEMR tube, which is a long transparent

Fig. 1. Schema of the procedures of strip biopsy

Fig. 2. Procedures of strip biopsy

A. Type IIa + IIc early gastric cancer is noted on lesser curvature site of antrum.

B. EUS shows ulcerative defect between the arrows with cancer invasion confined to mucosa.

C. After saline injection, the lesion is lifted with forceps and snaring with electrocautery loop wire.

D. Endoscopy shows post-EMR induced gastric ulcer.

E. Resected specimen shows well differentiated adenocarcinoma confined to mucosa.

Fig. 3. Schema of the procedures of ERHSE

- A. Identify the lesion, precisely.
- B. The marking around the lesion is made with electrocautery needle knife.
- C. Prior to resection, hypertonic saline with epinephrine is injected into the submucosa.
- D. Using the electrocautery knife, dissection of the lesion around the marks into lamina propria is performed.
- E. Retrieve the resected specimen with grasping forceps, or
- F-G. Resection of lesion with electrocautery and grasping forceps through 2 channel endoscope and retrieval of resected specimen are performed.

overtube designed to induce intraluminal negative pressure by endoscopic suction. The procedures are as follows: After injection of the saline-epinephrine solution around the lesion, the overtube is inserted near the lesion by the guidance of the endoscope. The snare is inserted through the side channel of the overtube and opened around the lesion. And then, the balloons located at the proximal end of the overtube are inflated in order to induce the intraluminally negative pressure by endoscopic suction. When the target lesion is aspirated into the overtube by negative pressure through the opened snare, the snare is closed, and the lesion entrapped by the snare is removed by electrocutting. Fig. 4 is a schema showing this procedure.

This technique has a advantage that enables to resect the larger lesion, but has a limitation that it cannot be used in the other site except the esophagus.

2) Endoscopic Mucosal Resection with a Cap Fitted Endoscope (EMRC)

EMRC had been devised by Inoue et al.(1993). In

EMRC, a small transparent cap fitted to the tip of endoscope is used. Fig. 5 is a schema representing the procedure. The tip of the endoscope is fitted with the specially designed transparent endoscope cap. The endoscope is inserted to the target lesion and more than 10 ml of saline-epinephrine solution is injected into the submucosal layer around the target lesion to elevate the lesion. After that, a small-sized snare is inserted through the working channel of endoscope and is opened inside the cap fitted at the tip of the endoscope. After the lesion is aspirated in the cap by full endoscopic suction, the lesion is snared and resected by electrocoagulation.

The advantages of EMRC are that this technique allows to resect the lesions on any part of the esophageal, gastric or colonic mucosa, and enables every part of the lesion to be observed easily regardless of the location and size of the lesion. But this technique has a disadvantage of the piecemeal resection being needed to deal with the lesion larger than 1.5 cm in diameter.

Fig. 4. Schema of the procedures of EEMR by EEMR tube
 A. Prior to resection, the saline mixed with epinephrine is injected into the submucosal layer.
 B-D. After the insertion of snare through the channel of EEMR tube, the target mucosa is snared and suctioned into the tube, and then strangulated by snare.
 E. Retrieval of resected specimen with retrieval device is performed.

Fig. 5. Schema of the procedures of EMRC
 A. The submucosal layer beneath the lesion is separated with an injection of saline.
 B-C. The target mucosa with a lesion is suctioned inside a cap and is strangulated by a snare.
 D. The strangulated mucosa is resected by electrocauterization.

3) Endoscopic Mucosal Resection using Band Ligator (EMRL)

In EMRL that has been devised by Lee et al.(1993), endoscopic variceal ligation kit (EVL kit) is used. The procedures are represented at Fig. 6. Under endoscopic guidance, saline or saline-epinephrine solution is injected into the submucosa of the target lesion until the lesion formed a polypoid bleb. The endoscope is then removed and fitted at the distal end with a friction adaptor preloaded with a rubber band. The endoscope

Fig. 6. Schema of the procedures of EMR with band ligation
 A. Prior to resection, saline is injected into the submucosal layer.
 B. The target mucosa with a lesion is suctioned inside a cap and ligated the mucosal lesion with band ligator.
 C-D. The ligated Yamada-III shaped lesion is resected by electrocauterization and retrieval of tissue is performed.

is reinserted to the target lesion and aspirate the lesion into the ligator. At that time, if the rubber band is deployed, the target lesion is ligated to create the polypoid lesion. And then the endoscope is withdrawn and the friction adaptor is removed. And the endoscope is reinserted, and the lesion formed a polyp is removed by snare polypectomy.

This technique is easier than other techniques, but has a limitation that it needs 3 times of insertion of the endoscope to resect the lesion larger than about 1.5 cm in diameter.

COMPLICATIONS OF ENDOSCOPIC MUCOSAL RESECTION

As in any other endoscopic procedures, EMR has a defined set of possible complications. The most significant complications of EMR are perforation and hemorrhage.

Most of the patient taking EMR do not complain the pain during resection. Since the pain during resection may indicate a transmural burn or perforation, the procedure should be immediately stopped when the patient complains the pain. The significant perforation has been very rarely reported(Tada et al.,1984 ; Hirao et al.,1988 ; Tada,1988 ; Takemoto et al., 1989 ; Kawano et al., 1991 ; Makuuchi et al., 1991 ; Inoue et al., 1993 ; Lee et al., 1993 ; Kaneko et al., 1995 ; Fleischer et al., 1996 ; Lee et al., 1996 ; Muto et al., 1996). Muto et al.(1996) reported that the incidence of perforation in 223 cases who had been performed EMR for gastric lesions was 0.45%. Takemoto et al.(1989) reported that the per-

foration did not occur in 445 lesions who had been performed EMR for gastric lesions. We had experienced the perforation in one(0.7%) of 149 cases(98 in stomach, 39 in colon, 12 in esophagus). However, perforation can occur if the deeper layers which are not sufficiently separated from the mucosa are entrapped in the snare, or the electrocoagulation current is used for a excessively great length of time. In order to prevent the perforation, the following three points should be carefully considered: 1) The sufficient amount of submucosal injection can prevent that electrocoagulation current is transmitted extensively to the deep layer. 2) The release-and-retightening method on snaring the lesion can allow that the deeper layers, which may be entrapped in the snare, are slipped-off. 3) Pure coagulation current on electrocoagulation increase the risk of perforation. Regarding the electrocoagulation current, the blend current is generally recommended. In our institute, pure cutting current or 4:1 blend (cutting : coagulation=4:1) current is used to resect the lesion in the esophagus or colon which has a thin wall. And 4:2 or 4:4 blend current is used to resect the lesion in stomach which has relatively thick wall or the lesion which is suspected to have high risk of hemorrhage on the procedure.

Although Muto et al.(1996) reported that the incidence of hemorrhage in 223 cases who has been performed EMR for gastric lesions was 14.3%, the incidence of procedure-induced significant hemorrhage has been generally reported 2 to 4%, and the incidence of major hemorrhage requiring transfusion has lower than 1% (Tada et al., 1984 ; Hirao et al., 1988 ; Tada, 1988 ; Takemoto et al., 1989 ; Kawano et al., 1991 ; Makuuchi et al., 1991 ; Inoue et al., 1993 ; Lee et al., 1993 ; Kaneko et al., 1995 ; Fleischer et al., 1996 ; Lee et al., 1996 ; Muto et al., 1996). Takemoto et al.(1989) reported that the incidence of hemorrhage during the procedure in 455 lesions performed endoscopic mucosal resection for the gastric lesions was 2.4%, and the incidence of the delayed hemorrhage after the procedure was 0.7%. In our experience, the incidence of hemorrhage during the procedure in 149 cases performed EMR for gastrointestinal tract was 1.3%, the incidence of the delayed hemorrhage was 2.0%. The reason, which the incidence of immediate hemorrhage in our data was lower than it in the reports of other authors, may be resulted from the fact that we usually used the epinephrine-saline solution for submucosal injection and performed endoscopic clipping for approximation of the large mucosal defect after endoscopic

mucosal resection. However, for immediate significant hemorrhage during the procedure, the endoscopic hemostatic techniques, such as hemoclippling, epinephrine injection, microwave hemostasis etc. should be performed (Choi et al., 1996).

The artificial ulcers resulting from endoscopic mucosal resection are moderate deep ulcer reaching to the submucosa, treated by the usual method of antiulcer drug therapy, and completely healed after 4-6 weeks.

INDICATIONS OF ENDOSCOPIC MUCOSAL RESECTION

The indications of EMR may be accomplished for diagnosis or therapeutic purposes. Table 1 is summarized of the indication of EMR.

In general, a biopsy carried out during endoscopy is best way to obtain accurate diagnostic information. Nevertheless, the correct histological diagnosis of some gastrointestinal lesion is often difficult, especially in excluding malignancy, since the conventional forceps biopsy specimens are too small to allow accurate diagnosis. On the contrary, the technique of EMR can potentially achieve the removal of the entire lesion, which is then available for thorough histological examination and allow accurate diagnosis. For example, the borderline adenoma, which are used the term as the adenoma in which conclusive diagnosis of malignancy cannot be made on the basis of histological examination of forceps biopsy specimens, can be correctly diagnosed by use of EMR. Fujiwara et al. (1996) reported that 28% of fifty cases, diagnosed as having borderline adenomas of the stomach on the basis of radiographic and endoscopic-biopsy findings, had been correct histological diagnosis as containing early gastric cancer after EMR.

Moreover, because EMR permits the resection of broad submucosal areas, the method makes it possible to histological diagnosis of the lesion, of which pathologic abnormalities are located in the deep mucosal

Table 1. Indications of endoscopic mucosal resection

1. Accurate diagnosis of the lesions in which the conventional forceps biopsy specimens are too small to allow diagnosis.
2. Histological diagnosis of the lesions which has pathologic abnormalities located in deep mucosa or submucosa (submucosal tumors, malignant lymphoma, benign lymphoid hyperplasia, Menetrier's disease, schirrhous type of GI cancer, etc.).
3. Diagnosis and therapy of gastrointestinal precancerous lesions (Flat adenoma, etc.).
4. Diagnosis and therapy of early gastrointestinal cancer.

or submucosal layers. Tada et al.(1988) reported that EMR had permitted the differential diagnosis between benign lymphoid hyperplasia and malignant lymphoma. We also had some experiences that the correct histological diagnosis had been possible in 44 cases with GI submucosal tumors, two cases with schirrhous type of gastric cancer, a case with melanoma, and a case with Menetrier's disease by use of EMR (Lee et al., 1991).

Since adenoma in gastrointestinal tract is a well-recognized precancerous lesion, the resection of entire adenoma is very important in the respect of preventing the cancer. Nevertheless, the flat or depressed adenomas can not be resected by conventional endoscopic polypectomy. On the contrary, since EMR makes it possible to resect the flat or depressed adenoma, the method may be very useful not only for a histological correct diagnosis but also for a curative treatment of flat or depressed lesions in gastrointestinal tract. In our unit, total 77 cases of flat or depressed adenomas of gastrointestinal wall were resected by EMR. 71(92.2%) of them were completely resected and finally diagnosed on the basis of histological examination of resected specimens. Moreover, EMR can be used for diagnosis and treatment of small early cancer without lymph node and distant metastasis. Regarding to this field, the follows are mentioned.

ENDOSCOPIC MUCOSAL RESECTION AS A CURATIVE THERAPY FOR EARLY CANCER

Progress in endoscopic technique and interest in early detection of gastrointestinal cancers has enabled early diagnosis of localized early cancers without any metastasis. Localized early cancers are likely to be cured by surgical resection, but there is the increasing evidence that endoscopic therapy may also be successful in cases with localized early cancer. Endoscopic therapy for early cancer is more advantageous than the conventional operative treatment with respect to relatively noninvasive, less expensive method. The techniques currently used for endoscopic treatment of early cancer in the gastrointestinal tract can be divided into two categories; tissue-destruction techniques(laser therapy, local injection therapy, etc) and tissue-resection techniques. Tissue-resection techniques have the advantages not only that the lesion can be examined histologically to reveal the extent and infiltration depth of the lesion, but also that the outcome of endoscopic therapy can be evaluated on the basis of

macroscopic and histologic findings. For these reasons, endoscopic resection technique has become the modality of the endoscopic therapy of early cancer in gastrointestinal tract.

Since endoscopic resection is completely ineffective in the lesion with lymph node or distant metastasis, a localized lesion without lymph node and distant metastasis is a prerequisite in order to cure early cancer by use of endoscopic therapy. Therefore, the indications of EMR for the purpose of curative treatment of early cancer should be considered carefully.

1. Indications of Endoscopic Mucosal as a Curative Treatment for Early Cancer

Curative treatment by EMR should only be performed for an early cancer without lymph node or distant metastasis. However, accurate detection of lymph node or distant metastasis before treatment may be difficult. Oguro(1991) reported that, on the basis of the review of histopathologic examination of 1430 surgical specimens the existence of lymph node metastasis in cases with gastric cancer was closely related to the depth of cancerous invasion, types and size. Thereafter, many similar reports(Watanabe,1988; Kato et al.,1990; Kudo, 1993; Endo et al.,1994) supported that the presence or absence of lymph node or distant metastasis was judged from the characteristics of the primary lesion and many investigator started to find the characteristic features of early cancer without lymph node or distant metastasis. In stomach, the characteristic features of early cancer without lymph node or distant metastasis reported by various investigators can be summarized at Table 2. On the basis of these data, many endoscopists have established the various criteria for application of EMR.

Generally, as the criteria for application of endos-

Table 2. The characteristic features of early gastric cancer without lymph node or distant metastasis

Well differentiated adenocarcinoma of type IIa not more than 2 cm in diameter
Well differentiated adenocarcinoma of type IIc, UL(-)*, not more than 1 cm in diameter
Mucosal cancer of type I
Focal cancer
Gastritis-like cancer lesion not more than 2 cm in diameter
Poorly differentiated cancer of type IIc, UL(-), not more than 5 mm in diameter, excluding lesions in the glands of the corpus ventriculi

* without disruption of the muscularis mucosa

copic mucosal resection in early gastric cancer, a well differentiated mucosal adenocarcinoma not more than 2 cm in size without ulceration is accepted.

In colon, the criteria for EMR are as follows; well differentiated adenocarcinoma of type Is, IIb, IIa, IIc, not more than 1 cm in size, M cancer (cancer confined to the mucosa) or SMla cancer (the cancer with minimally invasion to submucosa), without any invasion to lymphatic channels or vessels(Kudo, 1993).

In esophagus, the criteria for EMR are as follows; superficial esophageal cancer of type IIa, IIb, IIc, not more than 2 cm in size, intraepithelial cancer or mml cancer (cancer with minimally invasion to the muscularis mucosa) (Endo et al., 1994). Table 3 is summarized the criteria for curative treatment by EMR, used in our hospital. Endoscopic ultrasonography before EMR will help to define the depth of tumor invasion so that decision of endoscopic therapy should take into account the extent of the disease (Kim et al., 1991; Lee et al., 1995; Yasuda, 1996).

2. Estimation of the Outcome of Endoscopic Mucosal Resection for the Treatment of Early Cancer

Before EMR, the morphological characteristics of the lesion should be carefully evaluated by the endoscopy. If the lesion is compatible with the endoscopic criteria already mentioned, EMR could be performed. Computed tomography and endoscopic ultrasonography may be helpful to preoperatively evaluate the existence of the lymph node or distant metastasis.

After EMR, the histopathological evaluation of the resected specimen should be established in order to estimate the outcome of EMR as a curative treatment of early cancer. Regarding this point, histopathological evaluation of the resected specimen should be focused on whether the findings of the resected specimen are proper to the histopathological criteria of early cancer

without lymph node or distant metastasis, and whether the lesion is completely resected. Hamada et al.(1991) considered the lesion completely resected if the cancerous lesion was surrounded by normal mucosa not less than 2 mm in width. However, it is not practical to examine all serial slices from the specimen of 2 cm in size. Therefore Tada et al.(1993) used the following criteria to define the complete resection; no cancer tissue should be found in the first and last slices of the thin serial slices and, at the same time, normal mucosal layers including the muscularis mucosa should be observed on both sides of other slices. Recently, the criteria used by Tada et al.(1993) is widely accepted.

The complete resection rate of EMR in early cancer has been reported as 65-90% (Akamatsu et al., 1993; Endo, 1993; Igarashi et al., 1993; Takekoshi et al., 1994; Yokota et al., 1994). In our 45 cases with gastrointestinal early cancer (including cancers from adenomas) performed EMR, the complete resection rate was 86.7%.

If histological findings of the resected specimen are beyond the histological criteria, surgical operation should be necessary due to the possibility of lymph node or distant metastasis. And if the lesion is partially resected, further evaluation and treatment will be necessary. Fig. 7 shows a guideline for the further evaluation and treatment in the cases with incomplete resection (Takekoshi et al., 1994).

3. Follow-up Results after Endoscopic Mucosal Resection for the Treatment of Early Cancer

If the selection of cases for EMR is proper, the recurrence of cancer in lymph node or distant organ do not occur. Local recurrence of cancer after EMR are depended on whether the lesion was completely or incompletely resected on the basis of histological examination. Akamatsu et al.(1993) reported that the local recurrence was found in 8(27.6%) out of 29 early

Table 3. The criteria for curative treatment of early cancer by use of endoscopic resection

Site	Stomach	Colon	Esophagus
Endoscopic findings	Type I, any size IIa, IIb, < 2 cm IIc, < 1 cm	Type Is Ib < 1 cm IIa, IIa+IIc IIc, IIc+IIa	Type IIa IIb < 2 cm IIc
Histologic findings	Well defferentiated Mucosal cancer	Well defferentiated Mucosal cancer SMla cancer	ep cancer mm1 cancer

SMla cancer : the cancer which invades narrow and shallow into the submucosa

ep cancer : intraepithelial cancer

mm1 : cancer which invades shallow into the muscularis mucosa

Fig. 7. Follow-up scheme after endoscopic resection of early gastric carcinoma.

sm 1, 2 : degree of submucosal invasion

ly(0) : no lymph node metastases ; ly(1) : lymph node metastases

gastric cancer resected incompletely, in contrast to only one(3.8%) out of 26 resected completely.

Regarding the prognosis after EMR, the five-year survival rate after EMR in early gastric cancer is 88.1% not significantly different from 89.1% after surgical operation, according to the report by Tada et al.(1993). And the prognoses in the esophagus and colon after EMR are also similar with them after surgical operations (Endo, 1993 ; Igarashi et al., 1993 ; Yokota et al., 1994).

CONCLUSIONS

In conclusion, EMR is considered to be a useful procedure towards not only a correct diagnosis of the lesion that cannot be diagnosed by conventional forceps biopsy but a curative treatment of flat or depressed malignant mucosal lesions in gastrointestinal tract. Furthermore, since not only the frequency of early gastric cancer but cases suitable for EMR have been increasing steadily, EMR for treatment of early cancer seems to present a feasible alternative to surgery in selected cases. In practically, EMR for initial therapeutic method was practiced in 23.6% of cases with early gastric cancer from 1988 to 1990 in Japan (Nishi et al., 1995).

However, clearly careful follow-up at regular intervals after EMR is essential in order to establish whether EMR is a viable therapeutic option. And the further studies including large numbers of patients will be recommended in order to estimate the long-term results of EMR for curative treatment of early cancer.

REFERENCES

- Akamatsu T, Miyata K, Matsuzawa K, Fujimori Y, Hasebe O, Owa T, Suzuki A, Sakai H, Kiyosawa K, Furuta S, Katayama T. A long-term follow up study of cases undergoing endoscopic mucosal resection for early gastric cancer. *Gastroenterol Endosc* 1993 ; 5 : 97-104.
- Choi MG, Park SH, Bhang CS, Choi JY, Cho SH, Kim JK, Han JY, Chung KW, Sun HS. Endoscopic clip ligation on mucosal defect after endoscopic mucosal resection. *Kor J Gastrointest Endosc* 1996 ; 16 : 77-81.
- Endo M. Endoscopic resection as a local treatment of mucosal cancer of the esophagus. *Endoscopy* 1993 ; 25(suppl) : 672-4.
- Endo M, Takeshita K, Inoue H. Endoscopic mucosal resection of esophageal cancer. *Gan To Kagaku Ryoho* 1995 ; 22 : 192-5.
- Fleischer DE, Wang GQ, Dawsey S, Tio TL, Newsome J, Kidwell J, Pritti S. Tissue band ligation followed by snare resection (band and snare) : a new technique for tissue acquisition in the esophagus. *Gastrointest Endosc* 1996 ; 44 : 68-72.
- Fujiwara Y, Arakawa T, Fukuda T, Kimura S, Uchida T, Obata A, Higuchi K, Wakasa K, Sakura M, Kobayashi K. Diagnosis of borderline adenomas of stomach by endoscopic mucosal resection. *Endoscopy* 1996 ; 28 : 425-30.
- Hamada Y, Yoshimine H, Kubota H. Endoscopic resection for early gastric cancer measuring 1 cm or less in size. *Stomach Intest* 1991 ; 26 : 255-63.
- Hirao M, Masuda K, Asanuma T, Naka H, Noda K, Matsuura K, Yamaguchi O, Ueda N. Endoscopic resection of early gastric cancer and other tumors with local injection of hypertonic saline-epinephrine. *Gastrointest Endosc* 1988 ; 34 : 264-9.
- Igarashi M, Katsumata T, Naitho Y, Kobayashi K, Ohhara A, Saigenji K, Mitomi H, Atari E. Follow-up study of early colorectal cancer after endoscopic resection. *Dig Endosc* 1993 ; 5 : 359-65.
- Inoue H, Takeshita K, Hori H, Muraoka Y, Yoneshima H, Endo M. Endoscopic mucosal resection with a cap-fitted panendoscope for esophagus, stomach and colon mucosal lesions. *Gastrointest Endosc* 1993 ; 39 : 58-62.
- Kaneko E, Harada E, Kasugai T, Sakita T. The 2nd national survey of complications related to the gastrointestinal endoscopy from 1988 to 1992. *Gastroenterol Endosc* 1995 ; 37 : 642-52. (in Japanese)
- Kato H, Tachimori Y, Watanabe H, Yamaguchi H, Ishikawa T, Itabashi M. Superficial esophageal carcinoma : Surgical treatment and the results. *Cancer* 1990 ; 66 : 2319-23.
- Kawano T, Miyake S, Yasuno M, Takamatsu S, Katoh S, Nakamura H, Sugihara K, Hatano M, Yoshino K, Takeshita K, Inoue H, Yamagiwa A, Endo M. A new technique for endoscopic esophageal mucosectomy using a transparent overtube with intraluminal negative pressure(np-EEM). *Dig Endosc* 1991 ; 3 : 159-67.

- Kim JH, Park CW, Lee MS, Cho SW, Shim CS. Estimation for the depth of invasion in depressed types of early gastric cancer by endoscopic ultrasonography. *Korean J Gastrointest Endosc* 1991; 11: 261-71.
- Kudo S. Endoscopic mucosal resection of flat and depressed types of early colorectal cancer. *Endoscopy* 1993; 25: 455-61.
- Lee DK, Lee SW, Kwon SO, Jang WI, Shim YH, Cho MY. Endoscopic mucosectomy using an esophageal variceal ligation device for minute gastric cancer. *Endoscopy* 1996; 28: 386-9.
- Lee MS, Lee JS, Kim JS, Cho SW, Shim CS. The usefulness of suction ligation jumbo biopsy using an 'O'-band in esophago-gastric lesions. *Kor J Gastrointest Endosc* 1993; 13: 821.
- Lee MS, Park CW, Kim JH, Cho SW, Shim CS. Endoscopic enucleation of a gastric lipoma by strip biopsy. *Korean J Gastrointest Endosc* 1991; 11: 273-7.
- Lee MS, Rye KH, Lee JT, Kim JO, Bong HK, Cho YD, Kim YS, Lee JS, Hwang SG, Shim CS. The usefulness of endoscopic ultrasonography in endoscopic resection for gastric lesion. 10th International Symposium on Endoscopic Ultrasonography (Suppl), 1995.
- Makuuchi H, Machimura T, Mizutani K, Shimada H, Sugihara T, Tokuda Y, Sasaki T, Tajima T, Mitani T. Endoscopic mucosectomy for mucosal carcinomas in the esophagus. *J Gastroenterol Hepatol* 1991; 6(suppl 2): 5.
- Muto M, Saito Y, Koike T, Ikeya S, Sasaki T, Hoshino F. Complications of endoscopic polypectomy and endoscopic mucosal resection in stomach. *Gastroenterol Endosc* 1996; 38: 858-65. (in Japanese)
- Nishi M, Ishihara S, Nakajima T, Ohta K, Ohya S, Ohta H. Chronological change of characteristics of early gastric cancer and therapy; Experience in Cancer Institute Hospital of Tokyo 1950-1994. *J Cancer Res Clin Oncol* 1995; 121: 535-41.
- Oguro Y. Endoscopic treatment of early gastric cancer. *Dig Endosc* 1991; 3: 3-15.
- Ottenjann R, Lux G, Henke M, Strauch M. Big particle biopsy. *Endoscopy* 1973; 5: 139-43.
- Tada M. Strip biopsy. *Gastroenterol Endosc* 1988; 30: 2986-90. (in Japanese)
- Tada M, Iida Y, Sakaki N, Saito M, Oshita Y, Okazaki Y, Kawamura S, Takemoto T. A preliminary trial of jumbo biopsy using newly developed fiberscope. *Gastroenterol Endosc* 1980; 22: 450-7. (in Japanese)
- Tada M, Murakami A, Karita M, Yanai H, Okita K. Endoscopic resection of early gastric cancer. *Endoscopy* 1993; 25: 445-50.
- Tada M, Murata M, Murakami F. Development of the strip-off biopsy. *Gastroenterol Endosc* 1984; 26: 833-9. (in Japanese)
- Takekoshi Y, Baba Y, Ota H, Kato Y, Yanagisawa A, Takagi K, Noguchi Y. Endoscopic resection of early gastric carcinoma; Results of a retrospective analysis of 308 cases. *Endoscopy* 1994; 26: 352-8.
- Takemoto T, Tada M, Yanai H, Karita M, Okita K. Significance of strip biopsy, with particular reference to endoscopic mucosectomy. *Dig Endosc* 1989; 1: 4-9.
- Watanabe Y. Indication and limitation of endoscopic treatment for early gastric cancer. *Gastroenterol Endosc* 1988; 30: 1035-6. (in Japanese)
- Yasuda K. Endoscopic ultrasonic probes and mucosectomy for early gastric carcinoma. *Gastrointest Endosc* 1996; 43(suppl): 295.
- Yokota T, Sugihara K, Yoshida S. Endoscopic mucosal resection for colorectal neoplastic lesions. *Dis Colon Rectum* 1994; 37: 1108-11.