Published in final edited form as:

Nat Neurosci. 2007 December ; 10(12): 1578–1586. doi:10.1038/nn2005.

Specific requirement of NMDA receptors for long-term memory consolidation in *Drosophila* ellipsoid body

Chia-Lin Wu^{1,2,6}, Shouzhen Xia^{3,6}, Tsai-Feng Fu^{1,2,4}, Huaen Wang³, Ying-Hsiu Chen^{1,2}, Daniel Leong^{3,5}, Ann-Shyn Chiang^{1,2}, and Tim Tully³

¹Institute of Biotechnology and Department of Life Science, National Tsing Hua University, 101, Section 2, Kuang-Fu Road, Hsinchu 30013, Taiwan.

²Brain Research Center, National Tsing Hua University, University System of Taiwan, 101, Section 2, Kuang-Fu Road, Hsinchu 30013, Taiwan.

³Cold Spring Harbor Laboratory, 1 Bungtown Road, Cold Spring Harbor, New York 11724, USA.

⁴Department of Applied Chemistry, National Chi Nan University, University Road, Puli, Nantou 545, Taiwan.

Abstract

In humans and many other animals, memory consolidation occurs through multiple temporal phases and usually involves more than one neuroanatomical brain system. Genetic dissection of Pavlovian olfactory learning in *Drosophila melanogaster* has revealed multiple memory phases, but the predominant view holds that all memory phases occur in mushroom body neurons. Here, we demonstrate an acute requirement for NMDA receptors (NMDARs) outside of the mushroom body during long-term memory (LTM) consolidation. Targeted dsRNA-mediated silencing of *Nmdar1* and *Nmdar2* (also known as *dNR1* or *dNR2*, respectively) in cholinergic R4m-subtype large-field neurons of the ellipsoid body specifically disrupted LTM consolidation, but not retrieval. Similar silencing of functional NMDARs in the mushroom body disrupted an earlier memory phase, leaving LTM intact. Our results clearly establish an anatomical site outside of the mushroom body involved with LTM consolidation, thus revealing both a distributed brain system subserving olfactory memory formation and the existence of a system-level memory consolidation in *Drosophila*.

Pavlovian olfactory learning in *Drosophila* creates an elemental associative memory in a simple, accessible insect brain. This form of behavioral plasticity requires the normal function of NMDARs¹, as is the case in other invertebrate and vertebrate species^{2,3}. Memory formation in flies thereafter proceeds through several temporal phases and involves multiple biochemical cascades, which is again similar to findings from other animal models^{4–6}. In particular, spaced training produces stronger, longer-lasting memory than

© 2007 Nature Publishing Group

Correspondence should be addressed to A.-S.C. (aschiang@life.nthu.edu.tw) or T.T. (tully@cshl.edu).

⁵Present address: Department of Biomedical Engineering, City College of New York, 138th Street and Convent Avenue, New York, New York 10031, USA.

⁶These authors contributed equally to this work.

Note: Supplementary information is available on the Nature Neuroscience website.

AUTHOR CONTRIBUTIONS: C.-L.W. and S.X. conceived and designed the experiments. C.-L.W., S.X. and T.-F.F. carried out the experiments with technique support from H.W., D.L. and Y.-H.C. C.-L.W., S.X. and A.-S.C. analyzed the data. C.-L.W. and S.X. graphed the data. S.X. and T.T. wrote the paper.

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions>

massed training, which is a common property of memory formation^{7,8}. Contrary to the fact that memory storage involves multiple anatomical regions in other animal models⁹, olfactory memory has been proposed to be stored predominantly in the mushroom body neurons^{5,10}.

The mushroom body is a prominent neuropillar structure in the insect central brain. Intrinsic mushroom body cells send neurites ventrally into the calyx, a region of dendritic arborization that is innervated by efferents from several different regions, including projection neurons from the antennal lobes¹¹. Mushroom body axons project rostrally as a densely packed and stalk-like structure called the pedunculus to the anterior face of the brain, where they split and give rise to the dorsally projecting α and α' lobes and the medially projecting β , β' and γ lobes¹². Output neurons from the mushroom body project to many parts of the central brain. Another prominent neuropil is the central complex, which consists of four substructures, the ellipsoid body, the fan-shaped body, the nodulii and the protocerebral bridge. The central complex lies in the central brain between the pedunculi of the mushroom body and is bounded laterally by the two antennoglomerular tracts, dorsally by the pars intercerebralis, ventrally by the esophagus and the great commissure and frontally by the median bundle and the β -lobes of the mushroom bodies frontally¹³. The central complex forms intricate connections to a variety of brain centers, may mediate communication between the two hemispheres and is believed to be a control center for many different behavioral outputs^{13,14}.

The mushroom body does indeed have a central role in Pavlovian olfactory learning^{5,6,10}. Nevertheless, emerging evidence has hinted at the involvement of other extrinsic neurons and anatomical sites during olfactory memory consolidation. The *cer* gene, encoding a cathepsin inhibitor, most likely regulates LTM through its expression in glial cells surrounding the mushroom body¹⁵. In addition, several enhancer-trap transposon insertions with expression patterns exclusively outside of the mushroom body have been identified in a behavioral screen for olfactory memory mutants, although the functional relevance of these enhancer traps remains to be verified^{16,17}. Finally, an 'asymmetrical structure' in the central complex seems to correlate with the presence of LTM, but the functional relevance of this observation is also unknown¹⁸.

We recently have shown that NMDARs function in *Drosophila* during olfactory learning and thereafter during LTM consolidation¹. Here, using newly-constructed, dsRNA-mediated, UAS-driven transgenes to silence *dNR1* or *dNR2*, we have disrupted the normal function of NMDARs in the ellipsoid body or in the mushroom body, and have silenced synaptic transmission from both structures with UAS-driven dominant-negative *UAS-shits1* (ref. 19). We demonstrate dual dissociable roles for NMDARs in the ellipsoid body during LTM consolidation and in the mushroom body during the middle-term memory (MTM) phase of early memory processing. Our results suggest that LTM is stored in the ellipsoid body, outside of the mushroom body.

RESULTS

LTM is abolished by disrupting NMDARs in the ellipsoid body

Functional NMDARs in *Drosophila* consist of two subunits, dNR1 and dNR2, both of which are expressed widely in the adult brain, including in the mushroom body¹ and central complex (Supplementary Figs. 1 and 2 online). To address where functional NMDARs are required during olfactory memory formation, we used the binary GAL4/UAS gene-expression system²⁰ to target dsRNA-mediated knockdown of either the dNR1 or dNR2 subunits to various brain regions. For this purpose, we generated dsRNA-based transgenes (*UAS-dsNR1* and *UAS-dsNR2*) to silence gene expression in a sequence-specific manner²¹.

dsRNA triggers RNAi interference, an evolutionarily conserved process of sequence-specific post-transcriptional gene silencing^{22,23}. RNAi, however, can lead to the degradation of nontargeted mRNAs, commonly referred to as off-target effects. Using a recently established algorithm²⁴ and homology blast, we identified zero legitimate off-target sequences for *UAS-dsNR1* and only one (*CG17124*) for *UAS-dsNR2* (Supplementary Table 1 online). Therefore, dsRNA-mediated effects from both transgenes appeared to be highly specific for dNR1 and dNR2, and thus to functional NMDARs. Consistent with this expectation, the *UAS-dsNR1* transgene disrupted LTM in a quantitatively and qualitatively similar manner (Supplementary Fig. 3 online), as we previously have shown with a specific *dNR1* antisense message¹. We have not yet constructed a second dsRNA transgene for *dNR2*. dNR1 and dNR2 function together in *Drosophila*¹, however, so the silencing of *dNR2* constitutes a third disruption of NMDAR function. The fact that the LTM defect from our *UAS-dsNR2* transgene is similar to those from genetic disruptions of *dNR1* (see below) provides strong *in vivo* evidence against any different off-target effects from these transgenes. Taken together, the combined use of *UAS-dsNR1* and *UAS-dsNR2* provided us with a specific dsRNA-mediated silencing of functional NMDARs.

Western analysis revealed that the dNR2 protein was barely detectable in C155/+; *UAS-dsNR2*/+ flies (*elav/dsNR2*), whereas the heterozygous C155 (*elav*/+) and *UAS-dsNR2* (*dsNR2*/+) control flies produced a clear, single band that corresponded to the predicted sizes of the dNR2-2 and dNR2-3 proteins (Fig. 1a and Supplementary Fig. 2). Silencing of *dNR2* with *UAS-dsNR2* was further confirmed *in situ* with immunohistochemistry. Including our previously published antibody to *dNR2* (monoclonal α -5B10), we have developed four antibodies to *dNR2* that could be used for immunohistochemistry (see Supplementary Figs. 1 and 2 for details). All four antibodies labeled the ellipsoid body, suggesting that dNR2 was expressed preferentially in the ellipsoid body. Notably, two of these antibodies (α -820-2 and α -820-1) strongly labeled the ellipsoid body, suggesting an elevated expression of dNR2 in the ellipsoid body (Fig. 1b and Supplementary Fig. 2). This elevated expression was specifically diminished in *Feb170*/+; *UAS-dsNR2*/+ flies (Fig. 1c,d; see below for characterization of *Feb170*), whereas the expression of dNR2 in the protocerebral bridge (Fig. 1c,d) or in the optical lobes (Fig. 1e,f) was not affected. This reduction in dNR2 did not affect the gross morphology of the ellipsoid body, however, suggesting that a knockdown of dNR2 to this degree is not necessary for normal development of the ellipsoid body (Fig. 1g,h).

Western analysis also indicated that the dNR1 protein was reduced greatly in *UAS-dsNR1*/+; *hs-GAL4*/+ flies (*P26/dsNR1*) after heat shock, as compared with the heterozygous *hs-GAL4* (*P26*/+) or *UAS-dsNR1* (*dsNR1*/+) control flies (Supplementary Fig. 3). This confirmed our previous report that dNR1 could be acutely disrupted with heat shock using a heat shock-inducible GAL4 driver (*P26*)¹. Behavioral analyses suggested that LTM was abolished, but anesthesia-resistant memory (ARM) was normal in *UAS-dsNR1*/+; *hs-GAL4*/+ flies after heat shock-induced knockdown of *dNR1* (Supplementary Fig. 3), confirming our previous report using a different genetic method to knockdown dNR1.

The strong expression of dNR2 in ellipsoid body interested us, as another closely-related substructure of the central complex, the fan-shaped body, appears to subserve a short-term memory trace for visual learning²⁵ and may be involved with LTM formation after courtship conditioning²⁶. We identified several GAL4 drivers with preferential expression in these dNR2-positive ellipsoid body neurons (Fig. 2 and Supplementary Figs. 4 and 5 online). Three GAL4 drivers, *Feb170* (ref. 27), *Ruslan* (ref. 16) and *C42* (ref. 28), were chosen for further analyses, as their expression patterns overlapped only in these NR2-positive neurons of the ellipsoid body (Supplementary Fig. 5). Notably, all three GAL4 drivers showed no expression in the mushroom body (Supplementary Fig. 5). The ellipsoid body contains ten

types of small-field neurons that connect it to other central complex substructures (for example, the fan-shaped body)¹³. It also contains at least six types of large-field (R) neurons that arborize as a ring, mainly in the anterior part^{13,28}. Genetic mosaic flip-out analyses and immunolabeling indicated that the dNR2-positive neurons are the large-field, R4m subtype (H.-H. Lin, and A.-S.C., unpublished data). These dNR2-positive R4m neurons are likely cholinergic, as indicated by *Cha-GAL80* inhibition of their GAL4 expression (Supplementary Fig. 5).

Because we focused on genetic manipulations of *dNR1* in our previous study¹, we first disrupted the functional NMDARs in the ellipsoid body by knockdown of *dNR2*, and then confirmed our results with similar genetic manipulations of *dNR1* (which also is expressed in the ellipsoid body; ref. ¹ and Supplementary Fig. 2). Knockdown of *dNR2* using *Feb170-GAL4* to drive *UAS-dsNR2* specifically reduced the dNR2 protein expression in the ellipsoid body (Fig. 1b–f), without affecting the gross morphology of the ellipsoid body (Fig. 1g,h). The same knockdown also produced a 50% reduction in 1-d memory after spaced training (Fig. 2d). Similarly, disruption of NMDARs by Feb170-driven transgenic expression of *UAS-dsNR1* specifically disrupted 1-d memory after spaced training (Fig. 2f). We further confirmed this disruptive effect on 1-d memory after spaced training with Ruslan and C42 to drive *UAS-dsNR2* in the ellipsoid body (Supplementary Fig. 4). In contrast, the same genetic manipulations had no effects on 1-d memory after massed training (right panels, Fig. 2d,f), Feb170-driven expression of *UAS-dsNR2* (Fig. 2e) or *UAS-dsNR1* (Fig. 2g) had no effect on 3-h memory after one session of training and Feb170-driven expression of *UAS-dsNR2* (Supplementary Fig. 6 online) or *UAS-dsNR1* (Supplementary Fig. 6) had no effect on memory immediately after one session of training. Together with other behavior-genetic dissections dividing olfactory memory formation into four functionally distinct memory phases—short-term memory (STM), MTM, ARM and LTM^{8,17,29}—these observations suggest that genetic disruption of NMDAR function in the ellipsoid body specifically abolishes LTM, without affecting other earlier memory phases.

Protein synthesis–dependent LTM is specifically abolished

To rule out the possibility that a developmental defect underlay this LTM defect, we used the *tubulin* (*tub*)-*GAL80^{ts}* transgene with the binary GAL4/UAS system³⁰. Groups of flies, raised at 18 °C, were kept for 3 d at either 18 °C or 29 °C before spaced training. We determined that incubation for 3 d at 29 °C was sufficient for optimal disinhibition of a Feb170-driven *UAS-mCD8::GFP* reporter by *tub-GAL80^{ts}* (Supplementary Fig. 7 online). We observed that 1-d memory after spaced training was reduced significantly in transgenic *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies when kept at 29 °C for 3 d, whereas such memory was normal in the same flies when kept at 18 °C ($P < 0.05$; Fig. 3a), suggesting an acute physiological role for functional NMDARs in the ellipsoid body during LTM processing.

LTM uniquely depends on protein synthesis. Similar to previous studies^{8,29}, 1-d memory was reduced by 50% in normal flies kept at 29 °C for 3 d and fed cycloheximide (CXM) overnight before spaced training (left, Fig. 3b). However, in transgenic *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies kept at 29 °C for 3 d, CXM feeding did not further disrupt 1-d memory after spaced training (right, Fig. 3b). This result confirms that disruption of NMDARs specifically impairs protein synthesis–dependent LTM in the ellipsoid body.

Consolidation of LTM requires functional NMDARs

When kept at 29 °C (restrictive) for 4 d, shifted to 18 °C (permissive for inhibition of GAL4 by *GAL80*) for 3 d, subjected to spaced training and tested for 1-d memory at 18 °C, *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies showed normal 1-d memory (Fig. 3c). This result suggests that adult-specific disruption of dNR2 during the 4-d incubation at 29 °C was

restored to a normal functional level when the flies were shifted to 18 °C for 3 d. If flies were subjected to spaced training on day 3 of a 4-d incubation at 29 °C, however, 4-d memory (again at 18 °C for 3 d) was severely disrupted in *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies (Fig. 3d). The dNR2 expression was restored to a normal functional level at the time of testing (Fig. 3c), thereby revealing a disruption of consolidation and storage, rather than of retrieval, of LTM by knockdown of dNR2 in *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies.

A ‘reversal’ training protocol was designed to further distinguish a consolidation and storage defect from a retrieval failure. We subjected flies to a second reversal spaced training session 4 d after the first spaced training, where the original CS– became the CS+ (nonshocked odor and shocked odor, respectively), and vice versa. Thereafter, 1-d memory was quantified (performance indexes were calculated relative to the second CS+). In the absence of memory formation after the first spaced training session, reversal memory would be expected to be near a performance index of 35. If, on the other hand, memory formation after the first spaced training was normal, then it would counteract memory formation after reversal training, thereby producing a lower performance index that would be near zero. When *UAS-dsNR2/tub-GAL80^{ts}* and *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies were subjected first to spaced training on day 3 of a 4-d incubation at 29 °C and then to reversal spaced training after 3 d at 18 °C, 1-d reversal memory was significantly higher in *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies than in *UAS-dsNR2/tub-GAL80^{ts}* controls (Fig. 3e). This result demonstrates normal memory retrieval and implies weaker-than-normal memory storage from the first spaced training in *Feb170/+; UAS-dsNR2/tub-GAL80^{ts}* flies. Therefore, functional NMDARs contribute specifically to the consolidation and storage, but not to the retrieval of a protein synthesis-dependent LTM in the ellipsoid body.

Overexpression of dNR2 in the ellipsoid body enhances LTM

We observed that 1-d memory after one or two sessions of training was significantly enhanced when dNR2 was overexpressed in *Feb170*-targeted brain regions (Fig. 4a,b). This outcome is fully complementary to our dsRNA-based experiments: gene disruptions produce loss-of-function effects, whereas overexpression of the normal gene produces gain-of-function effects. Thus, NMDA receptors in *Drosophila* function in a manner that is quite analogous to a previous study in mice³¹.

LTM retrieval requires neural activity from the ellipsoid body

A physiological requirement for NMDARs in the ellipsoid body during LTM consolidation argues that a newly acquired olfactory experience is initially processed in or upstream of the mushroom body^{5,6,10}, but is then transferred from the mushroom body to the ellipsoid body for LTM storage. Such a hypothesis predicts that blocking synaptic output from the mushroom body, but not from the ellipsoid body, during memory consolidation (that is, during training and in a time window after training) would abolish LTM, and that blocking synaptic output from the ellipsoid body after memory consolidation is complete would disrupt LTM retrieval. To test these two predictions, we combined the *UAS-shi^{ts1}* transgene with *Feb170* or *OK107* GAL4 drivers. *OK107* is a GAL4 driver targeting transgenic expression in all lobes of the mushroom body (Supplementary Fig. 5). The *UAS-shi^{ts1}* transgene has been shown to block neuronal transmission in a temperature-dependent, dominant-negative fashion¹⁹.

To test the first prediction, we subjected *OK107/+; UAS-shi^{ts1}/+* flies, *Feb170/+; UAS-shi^{ts1}/+* flies or *UAS-shi^{ts1}/+* controls to spaced training and then maintained them for the first 6 h thereafter at a restrictive temperature (29 °C). Compared with the results from control experiments carried out at 18 °C (Fig. 5a) or with control flies, 1-d memory after

spaced training was abolished in *OK107/+; UAS-shi^{ts1}/+* flies, but was normal in *Feb170/+; UAS-shi^{ts1}/+* flies (Fig. 5b). To rule out the possibility that the 9-h blocking may cause irreversible disruption of synaptic transmission from the mushroom body, *OK107/+; UAS-shi^{ts1}/+* and *UAS-shi^{ts1}/+* flies were kept at 29 °C for 9 h and then shifted back to 18 °C for 18 h before being tested for initial learning after one session of training. Compared to *UAS-shi^{ts1}/+* controls, learning was normal in *OK107/+; UAS-shi^{ts1}/+* flies (68 ± 3.3 versus 61 ± 2.3 ; $P = 0.1$). This observation suggests that the abolition of 1-d memory in *OK107/+; UAS-shi^{ts1}/+* flies is specific to memory consolidation. These results demonstrate that neural activity from the mushroom body, but not from the ellipsoid body, is required during memory consolidation.

To test the second prediction, we subjected the same three genotypes to spaced training at 18 °C, maintained them at 18 °C for 23 h after training and then shifted them to 29 °C for 1 h before testing them at 29 °C. Under these conditions, 1-d memory after spaced training was blocked in both *OK107/+; UAS-shi^{ts1}/+* and *Feb170/+; UAS-shi^{ts1}/+* flies (Fig. 5c). Under the same conditions, however, sensorimotor responses to the odors were normal in both *OK107/+; UAS-shi^{ts1}/+* and *Feb170/+; UAS-shi^{ts1}/+* flies, as compared to *UAS-shi^{ts1}/+* controls (48 ± 4.1 or 40 ± 5.3 versus 49 ± 5.1 , $P = 0.34$ for 3-octanol; 48 ± 3.3 or 39 ± 4.9 versus 39 ± 5.1 , $P = 0.28$ for 4-methyl-cyclohexanol). This observation suggests that memory retrieval is specifically disrupted in both *OK107/+; UAS-shi^{ts1}/+* and *Feb170/+; UAS-shi^{ts1}/+* flies. Therefore, after LTM is consolidated, neural activity from either the mushroom body or ellipsoid body is required for retrieval. Together, these observations argue that consolidation and storage of olfactory LTM occurs upstream of synaptic transmission from the R4m neurons of the ellipsoid body. Considering that memory may be transferred outside of the mushroom body (during and within the first 6 h after training) for LTM consolidation (Fig. 5b) and that NMDARs are specifically required in the ellipsoid body for such a process (Figs. 2–4), our results suggest that early memory resides in (or upstream of) the mushroom body and then is transferred to the ellipsoid body during the consolidation of LTM.

NMDARs in the mushroom body are not required for LTM

The specific requirement of functional NMDARs in the ellipsoid body for LTM consolidation and storage prompted us to explore the role of NMDARs in the mushroom body (Fig. 6), the neuroanatomical structure known to mediate olfactory associations, STM and probably LTM^{5,10}. dNR2 or dNR1 in the mushroom body were specifically diminished when *UAS-dsNR2* (Fig. 6a–d) or *UAS-dsNR1* (Fig. 6g–j) were targeted to the mushroom body with OK107, whereas their expression in the optical lobes were not affected. We observed that 1-d memory after spaced or massed training was normal when dNR2 (Fig. 6e) or dNR1 (Fig. 6k) was knocked down, suggesting that functional NMDARs in the mushroom body are not required for the formation of LTM or ARM.

The latter conclusion was further confirmed by assessing 3-h memory after one session of training, which can be broken down into an anesthesia-sensitive memory (ASM) and ARM⁸. Knockdown of dNR2 (Fig. 6f) or dNR1 (Fig. 6l) via the OK107 mushroom body GAL4 driver similarly disrupted 3-h memory after one session of training. In contrast, 3-h memory was normal in these same transgenic flies when subjected to a 2-min cold shock 2 h after training, which provides a direct measure of ARM⁸. Similar effects were also observed when *UAS-dsNR1* was targeted to the mushroom body with two additional mushroom body GAL4 drivers, C739 and 201Y (Supplementary Figs. 5 and 8 online). Together, these data indicate that functional NMDARs in the mushroom body contribute to ASM. Memory retention immediately after one session of training was normal in transgenic flies expressing *UAS-dsNR2* or *UAS-dsNR1* via the OK107 mushroom body GAL4 driver (Supplementary Fig. 6), suggesting that STM is also not affected by knockdown of NMDARs in the

mushroom body. Therefore, MTM, but not STM, is the component memory phase of ASM that was specifically disrupted.

OK107 is a GAL4 driver labeling almost all of the mushroom body neurons³². Some of these neurons were probably cholinergic, including the γ lobe neurons that are primarily targeted by 201Y and a subset of the α/β lobe neurons that are primarily labeled by C739 (Supplementary Fig. 5). The rest, a subset of the α/β lobe neurons, were likely noncholinergic (Supplementary Fig. 5). Therefore, the cholinergic γ lobe neurons are targeted by both OK107 and 201Y, whereas the cholinergic α/β lobe neurons are targeted by both OK107 and C739. We observed that 3-h memory without cold shock in OK107/*dsNR2* flies did not differ from that with cold shock in control flies (Fig. 6f), suggesting that MTM was completely abolished. In contrast, MTM was only partially disrupted in 201Y/*dsNR1* or in C739/*dsNR1* flies (Supplementary Fig. 8). These results argue that MTM depends on cholinergic transmission and is completely abolished when NMDARs are silenced in all of these cholinergic α/β and γ cells.

DISCUSSION

The predominant view of olfactory memory is that it is processed in the mushroom body^{5,10}. Our data demonstrate a role for R4m neurons in the ellipsoid body during LTM consolidation, which now suggests a much broader and more complex neuronal circuitry sub-serving olfactory memory consolidation in *Drosophila*. Because genetic modulations of NMDARs in the ellipsoid body produce effects that are specific to LTM, components of this complex neural circuitry appear to subserve at least one specific temporal phase (LTM) of memory processing. Consistent and complementary to this notion, consolidation of LTM remains normal when NMDAR function is disrupted in the mushroom body (Table 1). Thus, the transference of memory from one anatomical location to another as consolidation progresses to LTM appears to occur in *Drosophila*, as in various other species^{6,9,33}.

NMDARs in the ellipsoid body and LTM consolidation

Specific disruption of functional NMDARs by dsRNA-mediated knockdown of either *dNR2* or *dNR1* in the R4m neurons of the ellipsoid body specifically abolishes protein synthesis-dependent LTM (Figs. 2 and 3 and Table 1), suggesting that the ellipsoid body is important during memory consolidation. NMDAR function is physiological rather than developmental, as induction of dsRNA transgenes in adults was sufficient to abolish LTM (Fig. 3). This role for NMDARs is also specific for memory phase (LTM) and brain region (ellipsoid body). Initial learning and early memories were not affected when NMDARs were knocked down in the ellipsoid body, and LTM was not affected when NMDARs were knocked down in the mushroom body (Table 1). These results, together with recent observations that LTM formation may require neuronal activity from the vertical lobes of the mushroom body and correlates with the appearance of a Fas II-immunoreactive asymmetrical body near the central complex^{18,34,35}, support a broader neuroanatomical circuitry involving both the mushroom body and ellipsoid body that subserves olfactory memory consolidation.

We also postulate that NMDARs in the ellipsoid body are involved with LTM consolidation and storage, but not retrieval. Disruption of NMDAR function during training, but not during testing, severely diminished LTM, excluding the possibility that memory retrieval was affected (Fig. 3c,d). Reversal memory was stronger when NMDARs were disrupted during an initial spaced training session compared with disruption during reversal training, directly demonstrating these flies' ability to retrieve memory (Fig. 3e). That retrieval is normal and that consolidation and storage have failed is consistent with our initial study¹ and with various studies of mammals³⁶.

Blocking synaptic output from the mushroom body, but not from the ellipsoid body, during training and in the first 6 h after training abolished consolidation of LTM (Fig. 5b), suggesting that consolidation of LTM occurs downstream of the mushroom body (or possibly in the mushroom body if the efferents from the ellipsoid body were to project back to the mushroom body). In contrast, blocking synaptic output from the ellipsoid body or mushroom body disrupted retrieval, but not acquisition and consolidation (Fig. 5c). This result is functionally analogous to STM, which is formed in or upstream of the mushroom body, probably without the involvement of extrinsic neurons or structures^{30,32,37}, and synaptic output from the mushroom body is required specifically for its retrieval, but not its acquisition^{38,39}. Taken together, our observations support a model where memory is first acquired in the mushroom body and is then transferred to the ellipsoid body for storage during memory consolidation, in agreement with various observations from other species where memory transfer from one brain region to another may occur^{6,33,36}. Combined with other recent studies, we would propose (i) that acquisition involves an initial association of an odor and shock in and/or upstream of the mushroom body^{5,10}, (ii) that STM resides in the mushroom body^{30,32,37}, (iii) that MTM is acquired in mushroom body α/β neurons and persists there via recurrent activity involving the mushroom body α'/β' DPM (dorsal-paired medial) neurons and the mushroom body α/β neurons themselves⁴⁰, (iv) that LTM consolidation involves the transference of memory to the ellipsoid body and (v) that LTM retrieval requires neural activity output from the ellipsoid body and from mushroom body neurons³⁴. Therefore, our results support a broader neuroanatomical circuitry involving both the mushroom body and ellipsoid body that subserves memory processing and retrieval.

Consolidation is the progressive stabilization of memory from a short, labile form to a long-lasting, stable form. Memory consolidation commonly refers to two types of processes, early and late consolidation. Early (or cellular or local) consolidation is accomplished within the first few minutes to hours after learning and occurs in all of the species studied to date⁴¹. This relatively fast type of consolidation takes place in local nodes in the neuronal circuit(s) and depends on cross talk among synapses, somata and nuclei^{41,42}. Late consolidation, which so far has only been demonstrated in humans and mammals, takes much longer and involves multiple brain systems^{36,41–43}. Late consolidation is initiated in parallel to, or as a consequence of, early consolidation, and is characterized by much slower temporal kinetics. Recent molecular and cellular studies in rodents have shown that a memory initially depends on the hippocampus, but eventually becomes independent of hippocampal function and may be consolidated into neocortical circuits^{36,41–43}. To our knowledge, the existence of multiple memory systems has not clearly been demonstrated in any invertebrate model system. Our demonstration that NMDARs are specifically required in the ellipsoid body rather than in the mushroom body during LTM consolidation shows that memory consolidation is a systems-level phenomenon in *Drosophila*.

It has recently been shown that LTM requires branch-specific neural activity and CREB in α/β mushroom body neurons³⁵. Notably, the only GAL4 driver used in that study, C739, is not specific to the mushroom body; it also labels some ellipsoid body neurons that appear to be different from the NMDAR-positive R4m neurons (Supplementary Figs. 5 and 9 online). Consistent with this observation, C739-targeted knockdown of functional NMDARs left LTM intact (Supplementary Fig. 8). When neural activity was silenced in *UAS-shi^{ts1}/+*; C739/+ flies, however, memory retrieval was disrupted³⁴. Considering that the cyclic AMP-dependent neuronal activity in the mushroom body is required for retrieval of LTM (S.X., H.W. and T.T., unpublished data), these results raise the interesting possibility that neural activity in the mushroom body α lobes may be correlated with memory retrieval rather than memory consolidation.

An intriguing difference exists in systems-level memory processing between *Drosophila* and rodents. Though LTM may eventually recruit the ellipsoid body, the mushroom body appears to be crucial for both memory consolidation and retrieval in *Drosophila*. In contrast, the hippocampus is required for consolidation, but not for retrieval, of long-lasting memories transferred to cortical systems^{36,41–43}. This difference may highlight an unusual aspect of olfactory memory in *Drosophila*. We recently have found that the encoding of odor identity and odor intensity is experience-dependent and mushroom body-dependent⁴⁴. Thus, the mushroom body appears to be central to olfactory discrimination as a perceptual task, providing a possible explanation as to why the mushroom body has to be involved with retrieval of LTM.

NMDARs in the mushroom body and MTM formation

The mushroom body is clearly involved in olfactory learning, memory consolidation and retrieval^{5,6,10}. Nevertheless, consolidation of LTM occurred normally even when the function of NMDARs in the mushroom body was disrupted (Table 1). MTM depends on NMDAR function in the mushroom body (Fig. 6) and on *amnesiac* (*amn*)-encoded neuropeptides⁴⁵ expressed in DPM neurons¹⁷. DPM neurons are extrinsic to the mushroom body, but nevertheless send extensive arborizations to all the mushroom body lobes¹⁷. Synaptic output from DPM neurons is required for the persistence of MTM, but does not appear to be involved in its acquisition and retrieval⁴⁶. Consistently, MTM is proposed to be formed in the mushroom body α/β neurons, and blocking mushroom body output diminishes the retrieval of MTM^{34,40}. Taken together, these observations suggest that the involvement of the mushroom body in LTM consolidation may be through an NMDAR-independent pathway, and that the mushroom body and ellipsoid body may participate independently in distinct temporal stages of memory consolidation, a neurobehavioral phenomena that has been reported in other species^{9,47}.

Understanding how the NMDAR-containing protein complexes are independently involved, at the cellular level, with different memory phases in distinct brain regions, and how these distinct anatomical sites communicate with each other to yield adaptive behavior from prior experience will be of particular importance in the future.

METHODS

Fly stocks

Flies used were wild-type Canton-S w¹¹¹⁸ (iso1CJ), the ‘Cantonized’ *UAS-dsNRI*, *UAS-dsNR2*, *UAS-shi^{ts1}*, *tub-GAL80^{ts}*, Feb170, Ruslan, C42, OK107, 201Y, C739, C155 (*elav-GAL4*), *hs-GAL4* (*P26*) and *UAS-mCD8::GFP*. Feb170 and C42 were from G. Korge (Freie Universitaet Berlin), *hs-GAL4* from Y. Zhong (Cold Spring Harbor Laboratory), *UAS-mCD8::GFP* from L. Luo (Stanford University) and *tub-GAL80^{ts}* from Bloomington Fly Center. The genetic backgrounds for all stocks were Cantonized by outcrossing their heterozygous virgins with w¹¹¹⁸ (iso1CJ) males for six generations.

Antibody production and western blotting

Rabbit antibodies to *dNRI* have been described previously¹ or are described in Supplementary Figs. 1 and 2. Rabbit polyclonal antibodies to *dNR2* were raised against various peptides corresponding to different regions of *dNR2* (Supplementary Figs. 1 and 2).

Western analysis was carried out as previously described¹. In brief, adult heads were homogenized with lysis buffer, centrifuged at 14,000 rpm at 4 °C for 50 min, and the supernatant was saved. Lysate proteins were electrophoresed on a 6% SDS-PAGE gel and then electroblotted onto PVDF membranes. Immobilized proteins were probed with a rabbit

polyclonal antibody to *dNR1* (α -85S at 1:5,000 dilution) polyclonal antibodies to *dNR2* (α -820-2 at 1:5,000 dilution, α -820-1 at 1:5,000 dilution or α -84S at 1:5,000 dilution) or a mouse monoclonal antibody to actin (1:2,500 dilution, Developmental Studies Hybridoma Bank) as a loading controls and the membrane was incubated with HRP-conjugated goat antibody to rabbit (1:10,000 dilution) or mouse (1:5,000 dilution) IgG. The positive signal was visualized with Qentix Western signal enhancer and SS West Pico Substrate detection (Pierce).

Behavior

Olfactory associative learning was measured by training 2–3-day-old adult flies in a T-maze with a Pavlovian conditioning procedure⁴⁸. Groups of about 100 flies received one or ten sessions of massed or spaced training during which they were exposed sequentially to one odor (conditioned stimulus, CS+; 3-octanol or 4-methyl-cyclohexanol) paired with electric shock and then to a second odor (CS–; 4-methyl-cyclohexanol or 3-octanol) without electric shock. Learning was measured immediately after one training session¹. We tested 3-h memory after one training session⁸. We evaluated 1-d or 4-d memory after spaced or massed training, which induces strong, long-lasting memory for conditioned avoidance⁸. Spaced training consisted of ten cycles of one session of training with a 15-min rest interval between each cycle. Massed training consisted of ten cycles of one session of training with no rest interval between each. Afterwards, flies were tested for memory retention of conditioned avoidance at the choice point of the T-maze after 1 or 4 d. A performance index was calculated as the number of flies avoiding the CS+ minus that avoiding the CS–, divided by the total number of flies and finally multiplied by 100.

Heat-shock regimen

Heat shock induction was carried out according to an established protocol with the same hs-GAL4 driver (P26)¹.

Drug feeding

CXM feeding was carried out according to an established protocol⁸.

Statistics

As a result of the nature of their mathematical derivation, performance indexes are distributed normally. Hence, the data were evaluated via one- or two-way ANOVAs. Subsequent pair-wise planned comparisons were adjusted for experiment-wise error (α'), keeping the overall $\alpha = 0.05$. All data were presented as mean $\pm$ s.e.m.

Methods used for the generation of flies carrying *UAS-dsNR1* and *UAS-dsNR2* transgenes, antibody production, immunohistochemistry and imaging whole-mount mCD8::GFP expression are described in the Supplementary Methods online.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank G. Korge, Y. Zhong, L. Luo and Bloomington Fly Center for fly stocks. We also thank M. Heisenberg, H. Cline and J. Dubnau for comments and discussion. This work was supported by grants to T.T. from the US National Institutes of Health and Dart Neurosciences, LLC, and to A.-S.C. from the National Science Council, the Brain Research Center of the University System of Taiwan and the Technology Development Program of Ministry of Economy.

References

1. Xia S, et al. NMDA receptors mediate olfactory learning and memory in *Drosophila*. *Curr. Biol* 2005;15:603–615. [PubMed: 15823532]
2. Roberts AC, Glanzman DL. Learning in *Aplysia*: looking at synaptic plasticity from both sides. *Trends Neurosci* 2003;26:662–670. [PubMed: 14624850]
3. Tsien JZ. Linking Hebb's coincidence-detection to memory formation. *Curr. Opin. Neurobiol* 2000;10:266–273. [PubMed: 10753792]
4. Allweis, C. The congruity of rat and chick multiphase memory-consolidation models. In: Andrew, RJ., editor. *Neural and Behavioral Plasticity*. New York: Oxford University Press; 1991. p. 370–393.
5. Davis RL. Olfactory memory formation in **Drosophila**: from molecular to systems neuroscience. *Annu. Rev. Neurosci* 2005;28:275–302. [PubMed: 16022597]
6. Margulies C, Tully T, Dubnau J. Deconstructing memory in *Drosophila*. *Curr. Biol* 2005;15:R700–R713. [PubMed: 16139203]
7. Kogan JH, et al. Spaced training induces normal long-term memory in CREB mutant mice. *Curr. Biol* 1997;7:1–11. [PubMed: 8999994]
8. Tully T, Preat T, Boynton SC, Del Vecchio M. Genetic dissection of consolidated memory in *Drosophila*. *Cell* 1994;79:35–47. [PubMed: 7923375]
9. Eichenbaum, H.; Cohen, NJ. *From Conditioning to Conscious Recollection: Memory Systems of the Brain* 583. New York: Oxford University Press; 2001.
10. Gerber B, Tanimoto H, Heisenberg M. An engram found? Evaluating the evidence from fruit flies. *Curr. Opin. Neurobiol* 2004;14:737–744. [PubMed: 15582377]
11. Stocker RF. The organization of the chemosensory system in *Drosophila melanogaster*: a review. *Cell Tissue Res* 1994;275:3–26. [PubMed: 8118845]
12. Jefferis GS, Marin EC, Watts RJ, Luo L. Development of neuronal connectivity in *Drosophila* antennal lobes and mushroom bodies. *Curr. Opin. Neurobiol* 2002;12:80–86. [PubMed: 11861168]
13. Hanesch U, Fischback K-F, Heisenberg M. Neuronal architecture of the central complex in *Drosophila melanogaster*. *Cell Tissue Res* 1989;257:343–366.
14. Strauss R, Heisenberg M. A higher control center of locomotor behavior in the **Drosophila** brain. *J. Neurosci* 1993;13:1852–1861. [PubMed: 8478679]
15. Comas D, Petit F, Preat T. **Drosophila** long-term memory formation involves regulation of cathepsin activity. *Nature* 2004;430:460–463. [PubMed: 15269770]
16. Dubnau J, et al. The *staufer/pumilio* pathway is involved in *Drosophila* long-term memory. *Curr. Biol* 2003;13:286–296. [PubMed: 12593794]
17. Waddell S, Armstrong JD, Kitamoto T, Kaiser K, Quinn WG. The *amnesiac* gene product is expressed in two neurons in the *Drosophila* brain that are critical for memory. *Cell* 2000;103:805–813. [PubMed: 11114336]
18. Pascual A, Huang KL, Neveu J, Preat T. Neuroanatomy: brain asymmetry and long-term memory. *Nature* 2004;427:605–606. [PubMed: 14961111]
19. Kitamoto T. Conditional modification of behavior in *Drosophila* by targeted expression of a temperature-sensitive shibire allele in defined neurons. *J. Neurobiol* 2001;47:81–92. [PubMed: 11291099]
20. Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* 1993;118:401–415. [PubMed: 8223268]
21. Carthew RW. Gene silencing by double-stranded RNA. *Curr. Opin. Cell Biol* 2001;13:244–248. [PubMed: 11248560]
22. Fire A, et al. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 1998;391:806–811. [PubMed: 9486653]
23. Hannon GJ. RNA interference. *Nature* 2002;418:244–251. [PubMed: 12110901]
24. Flockhart I, et al. Fly RNAi: the *Drosophila* RNAi screening center database. *Nucleic Acids Res* 2006;34:D489–D494. [PubMed: 16381918]
25. Liu G, et al. Distinct memory traces for two visual features in the *Drosophila* brain. *Nature* 2006;439:551–556. [PubMed: 16452971]

26. Sakai T, Tamura T, Kitamoto T, Kidokoro Y. A clock gene, period, plays a key role in long-term memory formation in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 2004;101:16058–16063. [PubMed: 15522971]
27. Siegmund T, Korge G. Innervation of the ring gland of *Drosophila melanogaster*. *J. Comp. Neurol* 2001;431:481–491. [PubMed: 11223816]
28. Renn SC, et al. Genetic analysis of the *Drosophila* ellipsoid body neuropil: organization and development of the central complex. *J. Neurobiol* 1999;41:189–207. [PubMed: 10512977]
29. Yin JC, et al. Induction of a dominant-negative CREB transgene specifically blocks long-term memory in *Drosophila*. *Cell* 1994;79:49–58. [PubMed: 7923376]
30. McGuire SE, Le PT, Osborn AJ, Matsumoto K, Davis RL. Spatiotemporal rescue of memory dysfunction in *Drosophila*. *Science* 2003;302:1765–1768. [PubMed: 14657498]
31. Tang YP, et al. Genetic enhancement of learning and memory in mice. *Nature* 1999;401:63–69. [PubMed: 10485705]
32. Lee T, Lee A, Luo L. Development of the *Drosophila* mushroom bodies: sequential generation of three distinct types of neurons from a neuroblast. *Development* 1999;126:4065–4076. [PubMed: 10457015]
33. Dash PK, Hebert AE, Runyan JD. A unified theory for systems and cellular memory consolidation. *Brain Res. Brain Res. Rev* 2004;45:30–37. [PubMed: 15063098]
34. Isabel G, Pascual A, Preat T. Exclusive consolidated memory phases in *Drosophila*. *Science* 2004;304:1024–1027. [PubMed: 15143285]
35. Yu D, Keene AC, Srivatsan A, Waddell S, Davis RL. *Drosophila* DPM neurons form a delayed and branch-specific memory trace after olfactory classical conditioning. *Cell* 2005;123:945–957. [PubMed: 16325586]
36. Nakazawa K, McHugh TJ, Wilson MA, Tonegawa S. NMDA receptors, place cells and hippocampal spatial memory. *Nat. Rev. Neurosci* 2004;5:361–372. [PubMed: 15100719]
37. Akalal DB, et al. Roles for *Drosophila* mushroom body neurons in olfactory learning and memory. *Learn. Mem* 2006;13:659–668. [PubMed: 16980542]
38. Dubnau J, Grady L, Kitamoto T, Tully T. Disruption of neurotransmission in *Drosophila* mushroom body blocks retrieval, but not acquisition, of memory. *Nature* 2001;411:476–480. [PubMed: 11373680]
39. McGuire SE, Le PT, Davis RL. The role of *Drosophila* mushroom body signaling in olfactory memory. *Science* 2001;293:1330–1333. [PubMed: 11397912]
40. Krashes MJ, Keene AC, Leung B, Armstrong JD, Waddell S. Sequential use of mushroom body neuron subsets during *Drosophila* odor memory processing. *Neuron* 2007;53:103–115. [PubMed: 17196534]
41. Dudai Y. The neurobiology of consolidations, or, how stable is the engram? *Annu. Rev. Psychol* 2004;55:51–86. [PubMed: 14744210]
42. Frankland PW, Bontempi B. The organization of recent and remote memories. *Nat. Rev. Neurosci* 2005;6:119–130. [PubMed: 15685217]
43. Wiltgen BJ, Brown RA, Talton LE, Silva AJ. New circuits for old memories: the role of the neocortex in consolidation. *Neuron* 2004;44:101–108. [PubMed: 15450163]
44. Xia S, Tully T. Segregation of odor identity and intensity during odor discrimination in *Drosophila* mushroom body. *PLoS Biol* 2007;5:e264. [PubMed: 17914903]
45. Feany MB, Quinn WG. A neuropeptide gene defined by the *Drosophila* memory mutant amnesiac. *Science* 1995;268:869–873. [PubMed: 7754370]
46. Keene AC, et al. Diverse odor-conditioned memories require uniquely timed dorsal paired medial neuron output. *Neuron* 2004;44:521–533. [PubMed: 15504331]
47. Emptage NJ, Carew TJ. Long-term synaptic facilitation in the absence of short-term facilitation in *Aplysia* neurons. *Science* 1993;262:253–256. [PubMed: 8211146]
48. Tully T, Quinn WG. Classical conditioning and retention in normal and mutant *Drosophila melanogaster*. *J. Comp. Physiol. [A]* 1985;157:263–277.

Figure 1.

dsRNA-mediated knockdown of *dNR2* in adult brain. (a) A schematic of the *UAS-dsNR2* transgenic construct is shown (left panel; see Methods for details). Western blot analysis revealed an appreciable knockdown of the dNR2 protein in adult heads of *C155/+; UAS-dsNR2/+* flies (*elav/dsNR2*) compared with *C155* (*elav/+*) and *UAS-dsNR2* (*dsNR2/+*) control genotypes (right panel). (b,c) Immunostaining for dNR2 was reduced in R4m neurons of the ellipsoid body in *Feb170/+; UAS-dsNR2/+* flies (*Feb170; dsNR2*, c) compared with *Feb170/+* control flies (*Feb170*, b). (d) Immunostaining of dNR2 was quantified in R4m neurons of the ellipsoid body and in the nearby protocerebral bridge for *Feb170/+* (black bars) and *Feb170/+; UAS-dsNR2/+* (white bars) flies. (e,f) Immunostaining of dNR2 was similar in optical lobes of *Feb170/+* (*Feb170*; e) and *Feb170/+; UAS-dsNR2/+* (*Feb170; dsNR2*; f) flies. (g,h) In spite of the knockdown of *dNR2* in *Feb170/+; UAS-dsNR2/+* flies (see above), the gross morphology of the ellipsoid body was normal in *Feb170/+; UAS-dsNR2, UAS-mCD8::GFP/+* flies (*Feb170; GFP, dsNR2*, g), as compared to *Feb170/+; UAS-mCD8::GFP/+* flies (*Feb170; GFP*, h). Scale bar, 50 μ m.

Figure 2.

Functional NMDARs in the ellipsoid body are required for LTM. (a–c) Preferential expression of dNR2 and *Feb170 GAL4* in R4m neurons of the ellipsoid body. Confocal imaging of dNR2 immunostaining (with α -820-2) in whole-mount adult brain revealed preferential expression (red) in R4m neurons of the ellipsoid body (posterior view, a). The *Feb170 GAL4*-driven expression of a *UAS-mCD8::GFP* reporter (green) in the ellipsoid body is shown (b). Merged images indicated overlap (yellow) of dNR2 staining (red) and GFP reporter (green) in R4m neurons of the ellipsoid body (EB, c). CB, cell body; LTR, lateral triangle. Scale bar, 50 μ m. (d) dsRNA-mediated knockdown of *dNR2* via *Feb170 GAL4*-driven expression of a *UAS-dsNR2* transgene specifically abolished LTM. We observed that 1-d memory after spaced training was significantly disrupted (* $P < 0.05$), whereas 1-d memory after massed training was normal in *Feb170/+*; *UAS-dsNR2/+*

(*Feb170/dsNR2*) flies. Heterozygous *Feb170* (*Feb170/+*) and *UAS-dsNR2* (*dsNR2/+*) control flies produced similar scores to wild-type flies (+/+) after either spaced or massed training ($n = 8$ performance indexes per group). (e) dsRNA-mediated knockdown of *dNR2* via *Feb170* *GAL4*-driven expression of the same *UAS-dsNR2* transgene had no effect on 3-h memory after one session of training in *Feb170/+*; *UAS-dsNR2/+* (*Feb170/dsNR2*) flies, as compared to control flies (+/+, *Feb170/+* or *dsNR2/+*) ($n = 8$ performance indexes per group). (f) dsRNA-mediated knockdown of *dNR1* via *Feb170* *GAL4*-driven expression of a *UAS-dsNR1* transgene specifically abolished LTM, similar to knockdown of *dNR2* ($n = 8$ performance indexes per group). (g) dsRNA-mediated knockdown of *dNR1* via *Feb170* *GAL4*-driven expression of a *UAS-dsNR1* transgene had no effect on 3-h memory after one session of training, similar again to knockdown of *dNR2* ($n = 8$ performance indexes per group).

Figure 3.

Inducible knockdown of dNR2 specifically blocks consolidation and storage, but not retrieval, of protein synthesis-dependent LTM. **(a)** Adult-specific knockdown of *dNR2* using the *GAL80^{ts}* repressor of *GAL4*-mediated *UAS-dsNR2* expression specifically abolished LTM at the restrictive temperature. After spaced training, 1-d memory was significantly disrupted in *Feb170/+; UAS-dsNR2/+; tub-GAL80^{ts}/+* (*Feb170/dsNR2; GAL80^{ts}*) flies when they were shifted from 18 °C to 29 °C for 3 d (right panel; *n* = 16 performance indexes per group) before training (SP, red arrow) and tested (black arrow) at 29 °C, but was normal when they were kept at 18 °C throughout the experiment (left panel; *n* = 8 performance indexes per group). **(b)** Adult-specific knockdown of *dNR2* abolished protein synthesis-dependent LTM. After spaced training, 1-d memory was reduced by 50% in CXM-fed wild-type flies (+/+ +CXM; left panel), but was not disrupted further in CXM-fed *Feb170/+; UAS-dsNR2/+; tub-GAL80^{ts}/+* (*Feb170/dsNR2; GAL80^{ts}* +CXM) flies, kept for 3 d at 29 °C before training and testing (*n* = 8 performance indexes per group). **(c)** After spaced training, 1-d memory was normal in *Feb170/+; UAS-dsNR2/+; tub-GAL80^{ts}/+* (*Feb170/dsNR2; GAL80^{ts}*) flies after 4 d at 29 °C, shifted to 18 °C for 3 d and followed by training and testing at 18 °C, indicating a recovery of NMDAR function at the permissive temperature. **(d)** Four-day memory after spaced training was significantly impaired in the same flies (*Feb170/dsNR2; GAL80^{ts}*) when incubated at 29 °C for 3 d, subjected to spaced training, maintained at 29 °C for 1 d and then shifted back to 18 °C for 3 d before testing, indicating that knockdown of NMDARs for 1 additional day after training is sufficient to block LTM consolidation and storage. **(e)** After reversal spaced training, 1-d memory (RS, blue arrow) was higher in the same flies (*Feb170/dsNR2; GAL80^{ts}*) than in controls when incubated at 29 °C for 3 d, subjected to spaced training, maintained at 29 °C for 1 d, shifted to 18 °C for 3 d, subjected to reversal training and tested at 18 °C 1 d later, indicating normal retrieval for the former (*n* = 8 performance indexes per group). * *P* < 0.05.

Figure 4.

Overexpression of dNR2 in the ellipsoid body enhances 1-d memory. Males homozygous for *UAS-dNR2-2* (a transgene expressing one of three dNR2 isoforms) were crossed with *Feb170* or wild-type (+/+) virgins, and wild-type males were also crossed with *Feb170* females. **(a)** We observed significantly higher 1-d memory after one session of training in *Feb170/+; UAS-dNR2-2/+* flies (*Feb170/dNR2*) than in wild-type (+/+) flies or in *Feb170/+* (*Feb170/+*) or *UAS-dNR2-2/+* (*dNR2/+*) control flies ($n = 18$ performance indexes per group). **(b)** Similarly, 1-d memory after two sessions of training was also higher in *Feb170/+; UAS-dNR2-2/+* flies (*Feb170/dNR2*) than in wild-type (+/+) controls ($n = 12$ performance indexes per group).

Figure 5.

Transference of memory from the mushroom body to the ellipsoid body during LTM consolidation. Wild-type (+/+), *Feb170* or *OK107* males were crossed to *UAS-shi^{ts1}* females, and all of the progeny were raised at 18 °C to minimize any potential leaky effect of *UAS-shi^{ts1}* on development. (a) Flies were kept at a permissive temperature (18 °C) throughout the entire experiment. After spaced training (SP), 1-d memory did not differ among *UAS-shi^{ts1}/+* (+/*shi^{ts1}*), *UAS-shi^{ts1}/+*; *OK107/+* (*OK107/shi^{ts1}*) or *Feb170/+*; *UAS-shi^{ts1}/+* (*Feb170/shi^{ts1}*) flies. (b) Flies were shifted to a restrictive temperature (29 °C), subjected to spaced training, maintained at 29 °C for another 6 h, shifted back to 18 °C for 18 h and tested at 18 °C. We observed that 1-d memory after spaced training was abolished in *UAS-shi^{ts1}/+*; *OK107/+* flies, but not in *Feb170/+*; *UAS-shi^{ts1}/+* flies, compared with *UAS-shi^{ts1}/+* controls. (c) Flies were subjected to spaced training at 18 °C, kept at 18 °C for 23 more h, shifted to 29 °C and then tested 1 h later. *UAS-shi^{ts1}/+*; *OK107/+* and *Feb170/+*; *UAS-shi^{ts1}/+* flies had impaired 1-d memory after spaced training, compared with *UAS-shi^{ts1}/+* controls, indicating that the output of neural activity from both the mushroom body and ellipsoid body is required during retrieval of LTM. In each of the above experiments, wild-type flies were also tested under the same conditions and showed consistent, higher scores (performance index = 48 ± 3 , 40 ± 2 , and 48 ± 5 for experiments a–c) than *UAS-shi^{ts1}/+* flies, as reported by others³⁹. $n = 8$ performance indexes per group.

Figure 6. NMDAR function in the mushroom body is required for MTM but not for LTM. (a,b) Immunostaining of dNR2 was specifically reduced in the mushroom body calyx (dendrites) in *OK107/+; UAS-dsNR2/+* flies (*dsNR2; OK107*, b), as compared with *OK107/+* controls (*OK107*, a). (c,d) dNR2 expression in optical lobes was similar between these flies. Scale bar, 50 μ m. (e) dsRNA-mediated knockdown of *dNR2* in the mushroom body did not affect 1-d memory after spaced or massed training in *OK107/dsNR2* flies, compared to controls (*+/+*, *OK107/+* or *dsNR2/+*), indicating that NMDAR function in the mushroom body is not required for LTM. (f) dsRNA-mediated knockdown of *dNR2* in the mushroom body disrupted 3-h memory after one session of training (without cold shock) in *OK107/dsNR2* flies, compared with control (*+/+*, *OK107/+* or *dsNR2/+*) flies. In contrast, 3-h memory after one session of training did not differ among these genotypes if the flies were subjected to a 2-min cold shock 2 h after training. Together, these data indicate that knockdown of *dNR2* in the mushroom body impairs MTM, but not ARM⁸ or LTM. (g–j) Similarly, immunostaining of dNR1 was specifically reduced in the mushroom body calyx in *UAS-dsNR1/+; OK107/+* flies (*dsNR1; OK107*, h), as compared with *OK107/+* controls (g), but was not affected in optical lobes in the same flies (i,j). (k) dsRNA-mediated knockdown of *dNR1* via *OK107 GAL4*–driven expression of the same *UAS-dsNR1* transgene did not affect 1-d memory after spaced or massed training, similar to knockdown of *dNR2*. (l) dsRNA-mediated knockdown of *dNR1* via *OK107 GAL4*–driven expression of the *UAS-dsNR1* transgene impaired MTM, but not ARM, again as with knockdown of *dNR2*. *n* = 8 performance indexes per group for 1-day memory and *n* = 6 performance indexes per group for 3-h memory.

Table 1

Effects of specific disruption of functional NMDARs in ellipsoid body or mushroom body on four different memory phases after olfactory learning

	Disruption of NMDARs in ellipsoid body (with <i>Feb170</i> as the GAL4 driver)		Disruption of NMDARs in mushroom body (with <i>OK107</i> as the GAL4 driver)	
	<i>UAS-dsNR2</i>	<i>UAS-dsNR1</i>	<i>UAS-dsNR2</i>	<i>UAS-dsNR1</i>
LTM	++	++	–	–
MTM	–	–	++	+
ARM	–	–	–	–
STM	–	–	–	–

++ indicates complete abolition; +, severe disruption; –, no effect at all.