CORRESPONDENCE

Lewy Body and Parkinsonian Dementia: Common, but Often Misdiagnosed Conditions

by PD Dr. med. Brit Mollenhauer, Prof. Dr. med. Hans Förstl, Prof. Dr. med. Günther Deuschl, Prof. Dr. med. Alexander Storch, Prof. Dr. med. Wolfgang Oertel, Prof. Dr. med. Claudia Trenkwalder in volume 39/2010

Hydrocephalus Should not Be Forgotten

Normal pressure hydrocephalus should be added to this successful overview of important differential diagnoses from the range of dementias. The condition affects at least 60 000 adult patients in Germany alone, and the estimated number of undetected cases is huge. According to current estimates, one in every 10 patients with dementia also has normal pressure hydrocephalus. The disease is often misinterpreted as part of the ageing process, since two-thirds of those affected are older than 60, with men affected twice as often as women. In contrast to the observational diagnosis in children with hydrocephalus, adults do not display any external signs or characteristics.

The lead symptoms of normal pressure hydrocephalus correspond to those of Alzheimer's disease and Parkinson's syndrome, with deterioration in performance, personality changes, memory loss, headaches, and unclear quasi-epileptic seizures. Without treatment the patients will develop, in addition to dementia, gait disturbances including a total inability to walk. The development of urinary incontinence (occasionally fecal incontinence) is typical. If a hydrocephalus persists for years and no treatment is given, a reduction in mental capacity to less than 20% has been documented, without the reflex status showing gross neurological deficits.

A timely diagnosis of the disorder is essential for the patient especially because effective and confirmed surgical therapeutic options are available these days—for example, regular lumbar punctures or neuro

surgical shunt therapy—which may not result in cure but which delay the inevitable disease course dramatically. The correct diagnosis is also of particular interest for the surgeon and anesthetist in the context of surgical interventions in these patients in order to ensure that deteriorations of the mental status of such patients are not hastily interpreted as the result of the anesthesia or surgery

DOI: 10.3238/arztebl.2011.0131a

REFERENCES

1. Mollenhauer B, Förstl H, Deuschl G, Storch A, Oertel W, Trenkwalder C: Lewy body and parkinsonian dementia: common, but often misdiagnosed conditions. *Dtsch Arztebl Int* 2010; 107(39): 684–91.

Prof. Dr. med. Christian-Friedrich Vahl

Universitätsmedizin Mainz
Klinik und Poliklinik für Herz-,
Thorax- und Gefäßchirurgie
Langenbeckstr. 1
55131 Mainz, Germany
cvahl@gmx.net

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

Evidence Is not Satisfactory

We learned from this article that only one study (*N Engl J Med* 2004) exists of ChE1 treatment in parkinsonian dementia. The ChE1 drug rivastigmine had the following effects: “[...]slight improvements in cognitive functioning, by 2.1 points in a scale of 0–70 of the ADAS-cog.”

Two of 70 points is such a small amount as to be hardly worth mentioning since such small improvements may easily be due to a measuring error or a statistical inaccuracy.

Further, the article mentions “notable clinical improvements” without explaining in more detail what this means.

It is interesting that the so called clinically notable improvements were observed in only 20% of patients and such “clinically notable improvements” were also observed in 15% of patients who were included in the study but received placebo.

This means that the drug had a positive effect that exceeded the placebo effect in only 5% of all patients. Does this justify administration of an expensive and barely tested medication to everybody, although the probability of its effectiveness is only 1:20?

DOI: 10.3238/arztebl.2011.0131b

REFERENCES

1. Mollenhauer B, Förstl H, Deuschl G, Storch A, Oertel W, Trenkwalder C: Lewy body and parkinsonian dementia: common, but often misdiagnosed conditions. *Dtsch Arztebl Int* 2010; 107(39): 684–91.

Dr. phil. Roman Landau

Maria-Louisenstr. 63
22301 Hamburg, Germany
info@winfried-vollmer.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.

In Reply:

Professor Vahl reminds us of the importance of normal pressure hydrocephalus as a differential diagnosis for dementia with motor symptoms. This disorder is indeed common and is often not recognized, although it is very well treatable by means of repeated lumbar puncture or surgical shunt implantation (1). This also underlines the importance of cerebral imaging (magnetic resonance imaging [MRI] is the technique of choice) in dementia and/or gait disturbances, which is often not performed. Without MRI the diagnosis of normal pressure hydrocephalus will be missed.

Dr Landau questions the use of cholinesterase inhibitors in Parkinson's disease with dementia. We mentioned in our article the controlled study using rivastigmine (2). A clinically notable improvement was seen in 19.8% of patients in the group receiving the drug but also in 14.5% in patients receiving placebo. Deterioration was noted in 13% in the "verum" group and 23.1% in the placebo group. These effects are low with respect to the total study group, but currently no clinically relevant alternative treatments are available. The study reported by Emre and colleagues is the only larger one with rivastigmine. A second study of a similar order of magnitude would be desirable. No controlled studies are currently available for comparable substances.

Personally in our clinical practice we often see clinical stabilization of patients with Lewy body dementia

after starting treatment with rivastigmine, in the form of cognitive improvement and reduced hallucinations

DOI: 10.3238/arztebl.2011.0132

REFERENCES

1. Relkin N, Marmarou A, Klinge P, Bergsneider M, Black PM: Diagnosing idiopathic normal-pressure hydrocephalus. *Neurosurgery* 2005;57(3 Suppl): 4–16; discussion ii–v.
2. Emre M, Aarsland D, Albanese A, et al.: Rivastigmine for Dementia associated with Parkinson's Disease. *N Engl J Med* 2004; 351(24): 2509–18.
3. Mollenhauer B, Förstl H, Deuschl G, Storch A, Oertel W, Trenkwalder C: Lewy body and parkinsonian dementia: common, but often misdiagnosed conditions. *Dtsch Arztebl Int* 2010; 107(39): 684–91.

Prof. Dr. med. Claudia Trenkwalder

PD Dr. med. Brit Mollenhauer

Paracelsus Elena-Klinik
Klinikstr. 16
34128 Kassel, Germany
brit.mollenhauer@pk-mx.de

Conflict of interest statement

Professor Trenkwalder has received honoraria for presentations or advisory board meetings from Cephalon, Boehringer Ingelheim, GlaxoSmithKline, Novartis, Solvay, TEVA, and Schwarz/UCB Pharma. Dr Mollenhauer receives financial funding from the Michael J Fox Foundation for Parkinson's Research, the American Parkinson Disease Association, and through a Dr Werner Jackstädt scholarship (Stifterverband für die Deutsche Wissenschaft—a German industry initiative promoting science and learning) and is in receipt of research funds from TEVA. She has received travel expenses from Boehringer- Ingelheim and Novartis and honoraria for speaking from GlaxoSmithKline and Bayer Schering Pharma AG.