Published in final edited form as:

EuroIntervention. 2011 March ; 6(8): 936–941. doi:10.4244/EIJV6I8A163.

Risk-Benefit Trade-offs in Revascularization Choices

Jerome J. Federspiel, BA¹, Sally C. Stearns, PhD², Ron T. van Domburg, PhD³, Brett C. Sheridan, MD⁴, Jennifer L. Lund, BA⁵, and Patrick W. Serruys, MD PhD³

¹School of Medicine, University of North Carolina, Chapel Hill, NC, USA ²Department of Health Policy and Management, University of North Carolina School of Public Health, Chapel Hill, NC, USA ³Thoraxcenter, Erasmus Medical Center Rotterdam, the Netherlands ⁴Department of Surgery, Division of Cardiothoracic Surgery, University of North Carolina Hospitals, Chapel Hill, NC, USA ⁵Department of Epidemiology, University of North Carolina School of Public Health, Chapel Hill, NC, USA

Abstract

Aims—When patients choose percutaneous coronary intervention (PCI) over coronary artery bypass grafting (CABG), they accept an increased long-term risk of repeat revascularization in exchange for short term morbidity benefits. This paper quantifies the risk-benefit trade-off faced by patients with multiple vessel coronary artery disease.

Methods and Results—Data from the Arterial Revascularization Therapies Study are used to generate risk-benefit acceptability curves for PCI versus CABG. Risks are measured by the long-term likelihood of repeat revascularization while benefits are measured by short term reductions in pain or improvements in health-related quality of life (HRQL). PCI patients faced a risk of 0.81 additional revascularization events over three years in exchange for being pain-free at one month. A patient would need to be willing to tolerate a risk of 1.06 additional revascularization events at three years, in exchange for being pain free at one month to be 95% confident that choosing PCI over CABG is risk-effective for him/her.

Conclusions—The risk-benefit framework outlined in this study provides information to enable physicians to help their patients weigh directly each procedure's risks and benefits. While trade-offs are typically measured in quality-adjusted life years, using pain reduction to reflect benefits may provide a more tangible framework for patients.

Keywords

Percutaneous stents; coronary artery bypass graft

Introduction

The optimal mechanical revascularization technique for multivessel coronary disease (MVD) remains contentious, with advocates supporting both percutaneous coronary intervention with insertion of stents (PCI) and coronary artery bypass grafting (CABG). Patients choose a procedure after consultation with physicians and may weigh a number of considerations in making their decision. Trends in the United States and elsewhere show that the volume of percutaneous procedures has been increasing while CABG rates decline.[1,2]

Corresponding Author: Sally C Stearns, PhD, CB 7411, 1104C McGavran-Greenberg Hall, Chapel Hill 27599-7411, (919) 843-2590 (Voice), (919) 966-6961 (Fax), sstearns@email.unc.edu.

This analysis was conducted at The University of North Carolina at Chapel Hill

These data indicate a growing preference for PCI over CABG for many patients and their physicians, especially for persons age 65 and older.[3]

Traditional cost-effectiveness analysis is useful for determining the societal benefit of these procedures, and this avenue of research has been incorporated in a number of comparisons of PCI and CABG.[4] [5] However, for patients with low cost sharing requirements for both procedures, the choice may be based on the valuation of benefits and risks rather than financial consideration. Clinical trial evidence indicates that both CABG and PCI increase health-related quality of life (HRQL) for patients with MVD. In the long-run, however, CABG shows substantially lower rates of revascularization when compared with bare metal stenting (PCI).[6] Receiving these deferred benefits of reduced rates of repeat revascularization requires accepting higher morbidity such as delayed relief from pain or improvement in HRQL in the time period immediately following the procedure.

Economists have become interested in using probabilistic methods to portray the trade-off between risks and benefits as well as the implicit valuation that patients make when one technique is chosen over a competitor. For example, probabilistic simulation modeling has been used to estimate the joint density of therapeutic risks and benefits from two pharmacological prophylactic treatments for deep-vein thrombosis.[7] In this study, we use clinical data from the Arterial Revascularization Therapy Study (ARTS) to estimate the trade-off between the benefit of quick reductions in post-procedure pain and HRQL when choosing PCI versus the increased risk of repeat revascularization. Specifically, this analysis uses a risk-benefit acceptability curve (RBAC), which is analogous to the cost-effectiveness acceptability curves used to portray uncertainty in the economic evaluation of health care technologies.[8] The RBAC provides estimates of the probability that a particular approach such as PCI is viewed by the consumer as being risk-effective (i.e., having greater value of benefits than cost of risks, analogous to cost-effectiveness) over different preference thresholds for the risk-benefit trade-offs identified. The estimates provided may be useful to physicians in explaining trade-offs between risks and benefits of alternative treatments to their patients, and the methodology can be applied to the current generation of PCI versus CABG studies as more complete data become available.

Methods

Between April 1997 and June 1998, ARTS randomized 1205 patients with MVD to receive PCI with bare metal stent implantation (n=600) or CABG (n=605).[5] Key details of the original study relevant for this analysis are the inclusion/exclusion criteria and the data collection strategy. The indications for revascularization for enrollment in ARTS included silent ischemia, stable or unstable angina pectoris, and the presence of at least 2 *de novo* lesions located in different major epicardial coronary arteries. Exclusion criteria included left ventricular ejection fraction <30%, left main stenosis, history of cerebrovascular accident (CVA), transmural myocardial infarction (MI) within the preceding week, severe hepatic or renal disease, and need for concomitant major surgery.[9] All patients gave written, informed consent. Follow-up information was obtained for patients at 1, 6, 12, and 36 months post-randomization, and the EuroQol EQ-5D instrument was administered to patients at each time point. More PCI patients completed the EQ-5D at 1 month than did CABG patients (93.5% versus 88.8%, $p = 0.005$); by 6 months, the difference was not significant and remained so for the rest of the follow-up period (90.2% vs 87.8%, $p=0.22$). All other data completion rates were comparable between treatment groups.

A binary variable indicating the presence or absence of clinical events was used to represent the risk in the decision to undergo PCI. Risk variables were calculated for three outcomes at 12 and 36 months post-randomization: (1) additional mechanical revascularization after the

randomized therapy; (2) additional PCI; and (3) additional CABG. Separate analysis of death, CVA and MI are not provided because their incidence did not differ in the ARTS trial, though disagreement remains about whether these outcomes differ between the two procedures. We used three measures to quantify the quality of life benefits of PCI. The pain/discomfort question on the EQ-5D was recoded to a binary variable indicating absence or presence of pain at the time of interview. For a more comprehensive view, a preference-weighted composite health-related quality of life measure (HRQL) was calculated from the EQ-5D survey responses.[10] Patients who did not complete the EQ-5D because they died prior to its administration were assumed to have a HRQL of zero for all subsequent time points. The HRQL measure was also used to estimate quality-adjusted life years (QALYs) post-intervention; since HRQL was only measured at four points post-randomization, we approximated QALYs using a linear area under the curve approximation (except for the first month, where the patient's HRQL at 1 month post randomization was used to calculate the entire first month of QALY since PCI offers such immediate pain relief relative to CABG).

Incremental risk was calculated as the difference in the proportions of patients experiencing additional revascularization for patients assigned to PCI versus CABG. Similarly, incremental benefit was calculated as the difference in proportions of patients reporting being free from pain and the difference in mean HRQL scores. Incremental risk-benefit ratios (IRBR) for various follow-up points in time were calculated by dividing incremental risk by incremental benefit:

$$IRBR = \frac{Risk_{PCI} - Risk_{CABG}}{Benefit_{PCI} - Benefit_{CABG}} \quad (1)$$

Bootstrapping with 5,000 replicates was conducted to form risk-benefit acceptability curves (RBAC) for each combination of selected risk time points (12 and 36 months) and one month benefit.[8] A longer time frame (up to 3 years) was chosen for risks while a shorter time frame was chosen for benefits because differences in benefits (pain or HRQL) were concentrated in the first few months after initial revascularization (see Figures 1A and 1B). All inference statistics were conducted at a two-tailed alpha level of 0.05, and results were analyzed on an intention to treat basis. Data analysis was conducted in SAS version 9.1.3 (SAS Institute, Cary, NC) and Stata version 10.1 (Statacorp, College Station, TX).

Rather than attempt to explicitly discount pain or clinical events benefits, we compared the long-term risks of PCI with its short-term benefits. As discussed later, discounting would increase the relative value of PCI because of greater short term pain reduction benefits in contrast to the benefits from CABG that take longer to accrue.

Results

As previously reported, rates of revascularization were higher among patients randomized to PCI than CABG, with the incremental risk for PCI versus CABG increasing from 17.2% to 20.2% when moving from 1 to 3 years post-randomization (Table 1).[6] [11] At three years post randomization, of the 160 PCI patients who had experienced additional revascularization 15 (2.5% of trial arm) had experienced both an additional PCI and an additional CABG while of the 39 CABG patients with repeat revascularizations only 2 (0.3%) had experienced both procedures after their initial treatment.

PCI patients were more likely to be free from pain and had higher HRQL scores shortly after randomization, though the effects were no longer statistically significant by six months after intervention. Table 1 shows that the incremental benefit of PCI decreases (from 0.25 to 0.04

on the freedom from pain measure and from 6.3 to -0.9 on the HRQL measure) when moving from the 1 month to 6 month time point, as many CABG patients recovered from the immediate post-procedure pain and HRQL impairment that occurs during the first few months following CABG. Figures 1A and 1B illustrate this trend graphically and show that composite HRQL from CABG reaches that of PCI slightly earlier than does freedom from pain, with both measures statistically equal from both procedures by 12 months post randomization. When QALYs are calculated using the HRQL scores, the difference in QALYs is minimal and not statistically significant after six months. Since repeat revascularizations accrue over time but significant differences in benefits occur only at one month after initial revascularization, we focus our comparisons on the trade-off of long-term risk for short-term benefit.

Together, the increased risk of revascularization from PCI and benefit trends for both procedures lead to increasing risk-benefit ratios as time from initial treatment randomization increases (Table 2). Bootstrapped estimates of the incremental risks and benefits and the distribution of the incremental risk/benefit ratio illustrate the trade-off more completely. All of the sampled incremental risk and benefit estimates were positive, meaning there was always a tradeoff between PCI and CABG and the IRBRs were always positive. As shown in the last column of Table 2 (bottom half), our data indicate that a patient or his/her physician would need to be willing to tolerate a risk of 1.06 additional revascularizations, including risk of 0.76 additional repeat PCI events and 0.44 additional CABG events, at 3 years in exchange for being pain free at 1 month in order to be 95% confident that choosing PCI over CABG is risk-effective for him/her. (By risk-effective, we mean that the risk is worth the benefit.) Similarly, for each 1 percentage point improvement in HRQL at 1 month post randomization, a patient would need to be willing to tolerate 4.8 additional revascularizations to be 95% confident that they are making a risk-effective choice in choosing PCI.

Figure 2 demonstrates the information in Table 2 visually through the formation of the risk-benefit acceptability curves (RBACs) from the distributions of incremental risk-benefit ratios. The RBAC illustrates the probability that PCI is risk-effective given a particular risk-benefit ratio threshold. [12] Each RBAC in Figure 2 represents the cumulative density of a single ratio of risk and benefit variables (the proportion of sampled ratios less than or equal to the specified ratio). Consequently, movements along the curve from the left to the right represent a willingness to accept a higher level of risk of revascularization from PCI in return for the immediate relief (i.e., reduction in pain or improvement in HRQL) from PCI relative to CABG. Discounting of benefits and risks (not done for simplicity of presentation) would shift the curves more to the left since the pain reduction from CABG is delayed relative to PCI.

We noted earlier that more PCI patients completed the EQ-5D at 1 month than did CABG patients. The lower response rates for the EQ-5D at 1 month post randomization among CABG patients likely leads to an underestimation of the pain reduction benefits of PCI, as patients with high pain at one month were probably less likely to complete surveys. To assess this possible bias, we recalculated the risk benefit ratios under the assumption that the difference in response rate between CABG and PCI patients at 1 month after randomization (4.7 percent) was comprised entirely of patients in pain (the least favorable case for CABG). Our point estimates of the risk benefit ratios would in this case overestimate the ratio by only 7.4%, indicating that any bias from differential response rates for CABG and PCI patients at 1 month is not substantial.

Discussion

This report uses risk-benefit ratio estimation and the corresponding risk-benefit acceptability curves to model the tradeoff between PCI and CABG. We model the choice of PCI over CABG as that of accepting greater long-term risk of additional revascularization in exchange for reduced post-procedure morbidity, specifically pain and composite HRQL. Rather than discount risks or benefits, we explicitly compared short term benefits and long term risks by comparing benefits at 1 and 6 months with risks at 12 and 36 months. In order for patients or their physicians to be 95 percent confident that their decision is risk-effective, a patient choosing PCI over CABG would need to be willing to accept the risk of 1.06 additional revascularizations over the next 3 years per patient benefiting by being free from pain at 1 month post-randomization. Analogously; a patient would need to accept 4.8 additional procedures per percentage point increase in HRQL.

Although the trial data are one decade old, the objective of this analysis is to develop a methodology that can enable more informed decision making on the part of patients and their physicians in choosing between PCI and CABG when neither procedure has a compelling indication or contraindication. Individual patients will value the trade-off differently; for some, exchanging the increased risk of repeat PCI or CABG to obtain short-term pain relief and HRQL increases will be acceptable, while others may prefer to endure short-term pain to obtain a higher probability of avoiding a subsequent revascularization. Additionally, patients may prefer to risk undergoing multiple PCI procedures rather than a single CABG, or they may prefer to avoid the risk of requiring CABG subsequent to PCI and instead have CABG initially.

The interpretation of RBACs has generally followed Bayesian guidelines. This approach has advantages from a policymaker's perspective, because it allows for the determination of whether an intervention is risk effective given a specified tolerance for risk. This delineation has clear value in exercises such as determining how much risk of severe adverse events would have to be tolerated to receive benefits. Composite HRQL measures generated from the EQ-5D provide a more comprehensive view of a patient's status than outcomes based strictly on freedom from pain. Yet the interpretation of HRQL is less intuitive, especially for patients. We suggest that the freedom of pain measure, while a simplification of HRQL, provides a useful way for physicians to work with patients to compare these two procedures, as presence and absence of pain is intuitive for most patients and providers. Distributions of risk-benefit ratios can provide patients and physicians with a sense of the trade-off between risks and benefits in direct comparison, not simply the separate magnitudes of risks and benefits. Patients would be best informed by being given both the risk-benefit ratio and the size of the absolute risks and benefits. The changes at the margin are likely most relevant for policy makers, and the fact that the statistically significant differences in outcome occur for just a short period after initial revascularization mean that the absolute increase in QALYs is very small.

Our study is not without limitations, especially in terms of applying the results in current clinical practice. Conceptually, our model depends on the assumption that the primary risk a patient experiences when choosing PCI over CABG is that of repeat revascularization, and the primary benefit is decreased pain. While most clinicians accept those risks and benefits as correct, especially with the use of bare metal stents, assertions about differences in rates of death and more severe complications are more contentious.[13–16] Furthermore, drug eluting stents have reduced risk of restenosis compared with that of bare metal stents, although controversy remains about the magnitude of this effect and potential increased risk of thrombotic events,[17–19] though longer-term follow-up data are not yet available. The results of the SYNTAX and FREEDOM trials, among others, will explore in detail the

different risks and benefits of PCI with drug eluting stent insertion when compared with CABG in patients with MVD. One year data from SYNTAX indicate the incremental risk of any repeat revascularization was reduced to 7.6%, compared with 17.2% in ARTS, although differences in inclusion criteria preclude direct comparison. [20]

An additional important consideration is the fact that the heterogeneity of treatment effects within populations is always of concern when applying trial results to clinical practice. [21] [22] For patients with less extensive disease or who have a shorter life expectancy, PCI may be more appealing, with the opposite being true for patients in better general health with more extensive cardiac disease. As described above in the study methods, ARTS only included patients with double and triple vessel disease (predominantly double) and excluded patients with a number of medical complications. The risk-benefit acceptability curves presented in this analysis are consequently most applicable to relatively healthy patients with MVD without several other medical complications, a relatively small portion of the universe of CAD patients. Yet the potential value of the approach for improving information available for decision-making by physicians and patients remains. Ultimately, as more comprehensive data are available, risk-benefit determination can become increasingly personalized.

Acknowledgments

This study was supported in part by NIGMS (5T32-GM008719) and NIA (1R01-AG025801). Suzanne West provided early inspiration for the analysis.

References

1. Ulrich MR, Brock DM, Ziskind AA. Analysis of trends in coronary artery bypass grafting and percutaneous coronary intervention rates in Washington state from 1987 to 2001. *The American Journal of Cardiology*. 2003; 92(7):836–839. [PubMed: 14516888]
2. Dawkins KD, Gershlick T, de Belder M, Chauhan A, Venn G, Schofield P, Smith D, Watkins J, Gray HH. S. Joint Working Group on Percutaneous Coronary Intervention of the British Cardiovascular Intervention, and S. the British Cardiac. Percutaneous coronary intervention: recommendations for good practice and training. *Heart*. 2005; 91 suppl_6:vi1–vi27. [PubMed: 16365340]
3. Gerber Y, Rihal CS, Sundt Iii TM, Killian JM, Weston SA, Therneau TM, Roger VL. Coronary Revascularization in the Community: A Population-Based Study, 1990 to 2004. *Journal of the American College of Cardiology*. 2007; 50(13):1223–1229. [PubMed: 17888838]
4. Stroupe KT, Morrison DA, Hlatky MA, Barnett PG, Cao L, Lyttle C, Hynes DM, Henderson WG. P. for the Investigators of Veterans Affairs Cooperative Studies. Cost-Effectiveness of Coronary Artery Bypass Grafts Versus Percutaneous Coronary Intervention for Revascularization of High-Risk Patients. *Circulation*. 2006; 114(12):1251–1257. [PubMed: 16966588]
5. Serruys PW, Unger F, Sousa JE, Jatene A, Bonnier HJ, Schonberger JP, Buller N, Bonser R, van den Brand MJ, van Herwerden LA, Morel MA, van Hout BA. Comparison of coronary-artery bypass surgery and stenting for the treatment of multivessel disease. *N Engl J Med*. 2001; 344(15): 1117–1124. [PubMed: 11297702]
6. Legrand VMG, Serruys PW, Unger F, van Hout BA, Vrolix MCM, Fransen GMP, Nielsen TT, Paulsen PK, Gomes RS, de Queiroz e Melo JMG, Marques dos Santos Neves JP, Lindeboom W, Backx B. I. on behalf of the Arterial Revascularization Therapy Study. Three-Year Outcome After Coronary Stenting Versus Bypass Surgery for the Treatment of Multivessel Disease. *Circulation*. 2004; 109(9):1114–1120. [PubMed: 14993134]
7. Lynd LD, O'Brien B J. Advances in risk-benefit evaluation using probabilistic simulation methods: an application to the prophylaxis of deep vein thrombosis. *J Clin Epidemiol*. 2004; 57(8):795–803. [PubMed: 15485731]
8. Fenwick E, O'Brien BJ, Briggs A. Cost-effectiveness acceptability curves - facts, fallacies and frequently asked questions. *Health Economics*. 2004; 13(5):405–415. [PubMed: 15127421]

9. Serruys PW, Unger F, Van Hout BA. The ARTS (Arterial Revascularization Therapies Study): Background, goals and methods. *Int J Cardiovasc Intervent.* 1999; 2(1):41–50. [PubMed: 12623386]
10. Greiner W, Weijnen T, Nieuwenhuizen M, Oppe S, Badia X, Busschbach J, Buxton M, Dolan P, Kind P, Krabbe P, Ohinmaa A, Parkin D, Roset M, Sintonen H, Tsuchiya A, Charro F. A single European currency for EQ-5D health states. *The European Journal of Health Economics.* 2003; 4(3):222–231. [PubMed: 15609189]
11. van Domburg R, Daemen J, Pedersen S, Bressers M, van Herwerden L, Firth B, Unger F, Serruys PW. Short- and Long- term health related quality-of-life and anginal status after randomisation to coronary stenting versus bypass surgery for the treatment of multivessel disease: results from the Arterial Revascularization Therapies Study (ARTS). *EuroIntervention.* 2007; (3):506–511. [PubMed: 19755292]
12. Briggs AH. A Bayesian approach to stochastic cost-effectiveness analysis. *Health Economics.* 1999; 8(3):257–261. [PubMed: 10348420]
13. Serruys PW, Ong ATL, van Herwerden LA, Sousa JE, Jatene A, Bonnier JJRM, Schonberger JPMA, Buller N, Bonser R, Disco C, Backx B, Hugenholtz PG, Firth BG, Unger F. Five-Year Outcomes After Coronary Stenting Versus Bypass Surgery for the Treatment of Multivessel Disease: The Final Analysis of the Arterial Revascularization Therapies Study (ARTS) Randomized Trial. *J Am Coll Cardiol.* 2005; 46(4):575–581. [PubMed: 16098418]
14. Hannan EL, Wu C, Walford G, Culliford AT, Gold JP, Smith CR, Higgins RSD, Carlson RE, Jones RH. Drug-Eluting Stents vs. Coronary-Artery Bypass Grafting in Multivessel Coronary Disease. *N Engl J Med.* 2008; 358(4):331–341. [PubMed: 18216353]
15. SoS Investigators. Coronary artery bypass surgery versus percutaneous coronary intervention with stent implantation in patients with multivessel coronary artery disease (the Stent or Surgery trial): a randomised controlled trial. *Lancet.* 2002; 360(9338):965–970. [PubMed: 12383664]
16. Daemen J, Boersma E, Flather M, Booth J, Stables R, Rodriguez A, Rodriguez-Granillo G, Hueb WA, Lemos PA, Serruys PW. Long-Term Safety and Efficacy of Percutaneous Coronary Intervention With Stenting and Coronary Artery Bypass Surgery for Multivessel Coronary Artery Disease: A Meta-Analysis With 5-Year Patient-Level Data From the ARTS, ERACI-II, MASS-II, and SoS Trials. *Circulation.* 2008; 118(11):1146–1154. [PubMed: 18725490]
17. Pfisterer M, Brunner-La Rocca HP, Buser PT, Rickenbacher P, Hunziker P, Mueller C, Jeger R, Bader F, Osswald S, Kaiser C. Late Clinical Events After Clopidogrel Discontinuation May Limit the Benefit of Drug-Eluting Stents: An Observational Study of Drug-Eluting Versus Bare-Metal Stents. *Journal of the American College of Cardiology.* 2006; 48(12):2584–2591. [PubMed: 17174201]
18. Malenka DJ, Kaplan AV, Lucas FL, Sharp SM, Skinner JS. Outcomes following coronary stenting in the era of bare-metal vs the era of drug-eluting stents. *JAMA.* 2008; 299(24):2868–2876. [PubMed: 18577731]
19. Abbott JD, Voss MR, Nakamura M, Cohen HA, Selzer F, Kip KE, Vlachos HA, Wilensky RL, Williams DO. Unrestricted Use of Drug-Eluting Stents Compared With Bare-Metal Stents in Routine Clinical Practice: Findings From the National Heart, Lung, and Blood Institute Dynamic Registry. *Journal of the American College of Cardiology.* 2007; 50(21):2029–2036. [PubMed: 18021868]
20. Serruys PW, Morice M-C, Kappetein AP, Colombo A, Holmes DR, Mack MJ, Stahle E, Feldman TE, van den Brand M, Bass EJ, Van Dyck N, Leadley K, Dawkins KD, Mohr FW, S.I. the. Percutaneous Coronary Intervention versus Coronary-Artery Bypass Grafting for Severe Coronary Artery Disease. *N Engl J Med.* 2009; 360(10):961–972. [PubMed: 19228612]
21. Kraemer HC, Frank E, Kupfer DJ. Moderators of Treatment Outcomes: Clinical, Research, and Policy Importance. *JAMA.* 2006; 296(10):1286–1289. [PubMed: 16968853]
22. Horwitz RI, Singer BH, Makuch RW, Viscoli CM. Can treatment that is helpful on average be harmful to some patients? A study of the conflicting information needs of clinical inquiry and drug regulation. *Journal of Clinical Epidemiology.* 1996; 49(4):395–400. [PubMed: 8621989]

Figure 1.

Figure 1A: Proportion of patients free from pain after randomization.

Figure 1B: Mean HRQL scores obtained by value weighting EQ-5D results.

Figure 2.

Figure 2A: Risk benefit acceptability curve for incremental risk of additional revascularization within 3 years versus freedom from pain at 1 month.

Figure 2B: Risk benefit acceptability curve for incremental risk of additional revascularization within 3 years of randomization versus HRQL benefits at 1 month.

Table 1

Risk and Benefit Variables Used in Analysis

	PCI	CABG	Incremental Difference ^a	95% CI for increment
Risks				
Additional Revascularization (PCI or CABG) ^b	N (%)		(%)	
12 mo	126 (21.0)	23 (3.8)	17.2	* (13.6, 20.8)
36 mo	160 (26.7)	39 (6.4)	20.2	* (16.2, 24.3)
Additional PCI				
12 mo	94 (15.7)	20 (3.3)	12.4	* (9.1, 15.5)
36 mo	120 (20.0)	36 (6.0)	14.0	* (10.3, 17.8)
Additional CABG				
12 mo	40 (6.7)	4 (0.7)	6.0	* (3.9, 8.1)
36 mo	55 (9.2)	7 (1.2)	8.0	* (5.5, 10.5)
Benefits				
% of Patients Free from Pain				
Baseline	151 (25.8)	150 (25.7)	0.1	(-4.9, 5.1)
1 Month	359 (64.0)	210 (39.1)	24.9	* (19.2, 30.6)
6 Months	348 (64.3)	322 (60.6)	3.7	(-2.1, 9.5)
12 Months	363 (67.5)	362 (67.2)	0.3	(-5.3, 5.9)
36 Months	323 (62.5)	321 (62.0)	0.5	(-5.4, 6.4)
HRQL at Time mean, full health = 100				
Baseline	63.9	63.5	0.4	(-2.2, 3.0)
1 Month	80.3	74.0	6.3	* (3.8, 8.9)
6 Months	81.2	82.1	-0.9	(-3.5, 1.7)
12 Months	81.7	83.0	-1.3	(-4.1, 1.4)
36 Months	79.3	79.1	0.2	(-2.8, 3.3)
QALYs since Randomization mean in years				
1 Month ^c	0.067	0.062	0.005	* (0.003, 0.007)
6 Months	0.400	0.391	0.013	* (0.002, 0.025)
12 Months	0.816	0.816	0.000	(-0.022, 0.022)

	PCI	CABG	Incremental Difference ^a	95% CI for increment
36 Months	2.450	2.495	-0.044	(-0.115, 0.027)

^a Differ slightly due to rounding

^b The sum of PCI and CABG exceeds the total revascularization estimate because some patients have both procedures after initial treatment.

^c Applies the QOL at 1 month for the entire first month. All other calculations done by averaging beginning and end QOLs

* p < 0.05

Table 2

Estimates of Incremental Risk / Benefit Ratios for PCI and CABG

	Sample Estimate	95% Probability Risk Effective
<i>One Year Post-Randomization Clinical Event Risk</i>		
Incremental risk of repeat revascularization at 12 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.66	0.91
Inc. benefit in HRQL, 1 mo	2.7	4.1
Inc. risk of repeat PCI at 12 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.48	0.66
Inc. benefit in HRQL, 1 mo	2.0	3.0
Inc. risk of repeat CABG at 12 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.23	0.34
Inc. benefit in HRQL, 1 mo	0.95	1.5
<i>Three Year Post-Randomization Clinical Event Risk</i>		
Incremental risk of repeat revascularization at 36 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.81	1.06
Inc. benefit in HRQL, 1 mo	3.2	4.8
Inc. risk of repeat PCI at 36 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.56	0.76
Inc. benefit in HRQL, 1 mo	2.3	3.4
Inc. risk of repeat CABG at 36 mo vs		
Inc. benefit in pain free proportion, 1 mo	0.31	0.44
Inc. benefit in HRQL, 1 mo	1.3	2.0