Unexpected outcome (positive or negative) including adverse drug reactions

Small intestinal intussusceptions due to the placement of a percutaneous endoscopic jejunostomy tube

Takayuki Satoh,¹ Kazue Sawada,¹ Miyuki Satoh,¹ Kikuchi Yohko,¹ Masataka Yamada,¹ Masaaki Zaitso,¹ Tadahiro Osada,¹ Reiji Sawaya,¹ Toshie Nata,² Nobuhiro Ueno,² Kentaro Moriichi,² Katsuya Ikuta,² Yusuke Mizukami,² Jiro Watari,³ Mikihiro Fujiya,² Yutaka Kohgo²

¹Shibetsu City Hospital, Shibetsu, Japan;

²Department of Medicine, Division of Gastroenterology and Hematology/Oncology, Asahikawa Medical University, Asahikawa, Japan;

³Hyogo College of Medicine, Nishinomiya, Japan

Correspondence to Mikihiro Fujiya, fjym@asahikawa-med.ac.jp

Summary

Percutaneous endoscopic jejunostomy (PEJ) has been developed and is considered to be a better method than percutaneous endoscopic gastrostomy for preventing the occurrence of aspiration pneumonia. However, the incidence of other complications associated with this procedure is less clear. We herein report a rare case with a small intestinal intussusception due to a PEJ placement. In this case, a radiologic examination with gastrografen was useful to detect the typical findings of a small intestinal intussusception, a beak-like filling defect, and identify the location of the lesion. An endoscopic examination that was carefully performed with a thin scope was effective to observe the ischaemic change of the small intestine and immediately determine the indication for surgical treatment. This case highlights the necessity to carefully manage patients with a PEJ placement, considering the risk of small intestinal intussusceptions when the patient complains of symptoms that are suspicious for an intestinal obstruction.

BACKGROUND

Artificial enteral nutrition has arisen as a major concern in clinical situations and home care because of the increase in both the elderly population and the patients with head and neck and esophageal disorders.¹ Percutaneous endoscopic gastrostomy (PEG) has become a popular procedure for establishing enteral feeding, although the procedure has possible complications including hiatus hernia, reflux esophagitis, and aspiration pneumonia.² Percutaneous endoscopic jejunostomy (PEJ) has subsequently been developed and is considered to be a better method than PEG for preventing aspiration pneumonia because the PEJ tube is placed at a more distal area through the stomach. While PEJ contributes to a decreased risk of refractory pneumonia due to the aspiration, the incidence of other complications in this procedure is less clear.^{3 4} The present case report describes an unusual case of small intestinal intussusception caused by a PEJ tube placement.

CASE PRESENTATION

An 84-year-old male with a 20-year history of chronic obstructive pulmonary disease visited our hospital due to dyspnea and dysphagia in December 2009. Because of the complaints of severe dysphagia and malnutrition, a PEJ was performed for enteral feeding. Sixty-five days after the insertion of PEJ, the patient complained of nausea, vomiting, and abdominal distention. A radiological examination with the injection of 50% gastrografen from the PEJ tube detected beak-like filling defect (figure 1). Ultrasonography (figure 2) revealed a target sign in the

corresponding site of the small intestine. Taken together, the observations led to a diagnosis of a small intestinal intussusception caused by the PEJ tube placement. To determine if surgery was needed to cure the small intestinal intussusceptions, an endoscopic examination using a thin scope (XP260N, Olympus Medical Systems, Tokyo, Japan) was carefully performed through the fistula beside the PEJ tube after obtaining a written informed consent. The disappearance of the vascular networks and mucosa discolored as dark red or grey were observed (figure 3), thus suggesting that ischaemic changes had occurred in this area of the small intestine. Surgery was immediately performed. The intraoperative findings showed that the dark red jejunum was dilated and invaginated (figure 4). A 2 m section of the jejunum was removed and a histological examination showed severe ischaemic change in the small intestine.

OUTCOME AND FOLLOW-UP

The patient died due to a cardiac failure 1 month after the surgical operation.

DISCUSSION

This report presents a rare case of small intestinal intussusception due to a PEJ tube placement, which required a jejunectomy to treat the severe ischaemic change in the small intestine. While the common complications of PEJ include pain, bleeding and cutaneous ulceration at the insertion site, peristomal leak and infection, tube extrusion, and aspiration pneumonia,²⁻⁴ only one case with the small intestinal intussusceptions and

Figure 1 Results of the radiologic examination. Radiologic examination with 50% of gastrografin revealed a beak-like filling defect in the proximal jejunum, thus suggesting a small intestinal intussusception due to the placement of the PEJ tube.

incomplete intestinal obstruction caused by a placement of PEJ tube has been described to date.⁵ This previously reported case was cured with conservative treatment because a severe obstruction or ischaemic change in the small intestine was not evident. The present case provides an alert that small intestinal intussusceptions due to PEJ may have the potential to cause a severe ischaemic change of the small intestine. Patients with a PEJ placement have to be carefully managed when they complain of symptoms that are suspicious for an intestinal obstruction.

In the current case, radiological examination with the injection of 50% gastrografin from the PEJ tube was useful to detect the typical finding of the intussusceptions, beak-like filling defect, and to identify the lesion's location, and subsequent endoscopic examination with a thin scope was effective to diagnose the ischaemic change of the small intestine and determine the indication of surgical operation. Consequently, a jejunectomy was immediately

performed and the intussusceptions and ischaemic change of the small intestine were relieved. It should be important to immediately grasp the condition of the small intestine by radiologic and endoscopic examinations when a patient complains of abdominal symptoms such as distention, nausea, and vomiting.

In summary, a rare case with a small intestinal intussusception due to the placement of a PEJ tube is reported. A radiologic examination with gastrografin was useful to detect the typical finding of the small intestinal intussusception, beak-like filling defect, and identify the location of the lesion. Endoscopic examination that was carefully performed with a thin scope was effective to observe the ischaemic change of the small intestine and immediately determine the indication for surgical treatment. Therefore, it is important to carefully manage patients with a PEJ placement under considering a risk of the small intestinal intussusceptions when the patient complains of symptoms that are suspicious for an intestinal obstruction.

Figure 2 Results of the ultrasonographic image. Typical target sign suggestive of small intestinal intussusceptions was apparent in the middle area of the abdomen.

Figure 3 Endoscopic image of the intussusception site. An endoscopic examination, which was carefully performed through the fistula beside the PEJ tube, detected the disappearance of the vascular networks and mucosa that was discolored dark red or grey. These observations indicated severe ischaemic change in the small intestine.

Figure 4 Intraoperative findings of the small intestine. An intestinal intussusception due to the PEJ tube and a 2 m portion of the dark red jejunum was observed, suggesting intestinal damages due to severe ischaemic change throughout almost the entire jejunum.

Learning points

- ▶ PEJ may cause small intestinal intussusceptions.
- ▶ It is important to carefully manage patients with a PEJ placement when the patient complains of symptoms that are suspicious for an intestinal obstruction.
- ▶ A radiologic examination was useful to detect the typical finding of the small intestinal intussusception and identify the location of the lesion.
- ▶ Endoscopic examination that was carefully performed with a thin scope was effective to observe the ischaemic change of the small intestine and immediately determine the indication for surgical treatment.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Gauderer MW**, Ponsky JL, Izant RJJr. Gastrostomy without laparotomy: a percutaneous endoscopic technique. *J Pediatr Surg* 1980;**15**:872–5.
2. **Nishiwaki S**, Araki H, Goto N, *et al*. Clinical analysis of gastroesophageal reflux after PEG. *Gastrointest Endosc* 2006;**64**:890–6.
3. **Kadakia SC**, Sullivan HO, Starnes E. Percutaneous endoscopic gastrostomy or jejunostomy and the incidence of aspiration in 79 patients. *Am J Surg* 1992;**164**:114–18.
4. **Shetzline MA**, Suhocki PV, Workman MJ. Direct percutaneous endoscopic jejunostomy with small bowel enteroscopy and fluoroscopy. *Gastrointest Endosc* 2001;**53**:633–8.
5. **Ishii M**, Teramoto S, Yakabe M, *et al*. Small intestinal intussusceptions caused by percutaneous endoscopic jejunostomy tube placement. *J Am Geriatr Soc* 2007;**55**:2093–4.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Satoh T, Sawada K, Satoh M, Yohko K, Yamada M, Zaitzu M, Osada T, Sawaya R, Nata T, Ueno N, Moriichi K, Ikuta K, Mizukami Y, Watari J, Fujiya M, Kohgo Y. Small intestinal intussusceptions due to the placement of a percutaneous endoscopic jejunostomy tube. *BMJ Case Reports* 2011;10.1136/bcr.07.2010.3169, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow