Rare disease

Diagnosis of longitudinally extensive transverse myelitis

Helen Nightingale, Jonathan Witherick, Alastair Wilkins

University of Bristol, Bristol, UK

Correspondence to Alastair Wilkins, alastair.wilkins@bristol.ac.uk**Summary**

Longitudinally extensive transverse myelitis (LETM) is a neurological condition characterised by a contiguous inflammatory lesion of the spinal cord. LETM is often associated with the autoimmune central nervous system disease neuromyelitis optica (NMO) and rarely with multiple sclerosis. The discovery of the NMO-IgG antibody, provides a useful serological marker of LETM associated with the NMO disease spectrum (LETM and/or optic neuritis). Here, the authors report two cases of LETM, which differ in disease severity and NMO-IgG antibody serological status.

BACKGROUND

The clinical course of longitudinally extensive transverse myelitis (LETM) is characterised by single or multiple attacks of paraparesis or tetraparesis, sensory deficits and bowel/bladder disturbances,¹ and in severe cases can lead to respiratory failure. LETM is defined by a spinal MRI indicating a lesion extending over three or more vertebral segments.² The early distinction between possible LETM aetiologies is crucial in providing accurate prognosis and guiding therapeutic strategy. The diagnosis of LETM as part of the neuromyelitis optica (NMO) disease spectrum, is suggested by NMO-IgG antibody seropositivity.² LETM patients with NMO-IgG seropositivity are considered at increased risk of future relapses and development of the whole NMO disease spectrum.³ Thus, early serological testing for NMO-IgG is important.

CASE PRESENTATION**Case 1**

A 78-year-old Caucasian man was admitted with a 1-week history of progressive bilateral leg weakness that had left him bed-bound. He also complained of sensory changes in his feet and difficulty with micturition. There were no reported visual symptoms.

Examination of his cranial nerves and upper limbs was unremarkable. Lower limb examination revealed bilateral weakness (medical research council (MRC) grade 1 on the right and 2–3 on the left) with areflexia and flexor plantar responses. A sensory level was noted at T4 but distal proprioception remained intact. He had reduced anal tone and a urinary catheter was required on admission to hospital.

Case 2

A 31-year-old woman presented with progressive bilateral leg weakness and numbness over a 5-day period. She also complained of a 'band like' sensation around her trunk. Her vision, bowel and bladder were normal. Five years previously she had an episode of left leg numbness followed by progressive weakness over a 5-week period resulting in difficulty walking. A spinal MRI at the time showed an area of high intensity extending from C5/C6 to T3/T4 and visual evoked potentials were normal. Following a course

of intravenous steroids, power and sensation in her leg improved, and repeat spinal MRI had shown reduced cord swelling. She had no symptoms in the interim and no previous neurological problems.

On examination, cranial nerves and upper limbs were normal. She had bilateral weakness of hip flexion, right knee extension and ankle dorsiflexion (MRC grade 4). Reflexes were brisk throughout the lower limbs with flexor plantar responses. She had reduced sensation to light touch and vibration in her left lower limb and a sensory level at T10. Proprioception in the left foot was reduced up to the ankle. She remained ambulant.

INVESTIGATIONS**Case 1**

MRI of his spine revealed an area of high signal intensity extending from C5/6 to T10 on T2 sequences (figure 1). No lesions were shown on brain MRI. Standard blood markers for vasculitic conditions, vitamin B12 and syphilis serology tests were negative. He was seropositive for NMO-IgG antibodies. Cerebrospinal fluid (CSF) was acellular with a raised protein level of 1.03 g/l and immunoglobulin bands which matched with serum.

Case 2

An area of high intensity extending over 4½ segments was found on MRI spine (figure 2), with a normal MRI brain. CSF analysis was normal with no oligoclonal bands present. Standard blood markers for vasculitic conditions, vitamin B12 and syphilis serology tests were negative. Visual evoked potentials were normal. NMO-IgG antibodies were negative.

DIFFERENTIAL DIAGNOSIS

- ▶ NMO
- ▶ Infectious myelitis
- ▶ Myelitis in association with systemic autoimmunity, for example, systemic lupus erythematosus
- ▶ Paraneoplastic myelitis
- ▶ MS

Figure 1 T2-weighted MRI of the spine of case 1. A continuous demyelinating lesion can be seen extending from C5/6 to T10.

Figure 2 T2-weighted MRI of the spine of case 2. A continuous demyelinating lesion can be seen extending from C5/C6 to T3/T4.

TREATMENT

Case 1

He received several courses of intravenous methylprednisolone and plasma exchange, causing some improvement in sensation (returned to T10 level) but no improvement in motor function in his legs.

Case 2

Following a 3-day course of intravenous methylprednisolone she improved quickly and returned to normal in 3 months.

DISCUSSION

LETM is an inflammatory condition of the spinal cord causing variable levels of morbidity and mortality. Previous case reports of LETM have focused upon the early distinction of LETM from MS to enable prompt and appropriate therapy, here we additionally emphasise the clinical importance of NMO-IgG serological status.

Knowledge of LETM aetiology is helpful to initiate appropriate therapy and predict prognosis. A recent study of the NMO disease spectrum described a 77% prevalence of LETM among 13 patients,⁴ whereas another study reported diffuse spinal cord lesions in only 13% of 104 conventional MS patients.⁵ Features distinguishing NMO disease spectrum from MS associated LETM include, spinal cord lesions involving both grey and white matter, which span three or more vertebral segments and are prone to cavitation; brain lesions which are linear rather than oval without a Dawson finger configuration; the presence of serum NMO-IgG and absence of CSF oligoclonal bands.⁶ Other rare causes of LETM include infectious myelitis (including viral, fungal, parasitic and bacterial); other autoimmune conditions including neurological lupus, neurosarcoidosis, Behcet's disease, antiphospholipid syndrome and Sjögren's syndrome; paraneoplastic conditions; and vascular abnormalities, such as arteriovenous malformations.⁶ Careful history taking, clinical examination, blood serology, CSF analysis and diagnostic imaging are helpful in differentiating between these numerous causes.

The differentiation between MS and NMO disease spectrum associated LETM is important, as NMO has a worse prognosis and earlier and more frequent relapses.⁷ Furthermore, immuno-modulatory therapy is preferential for MS, whereas immuno-suppressive therapy is suggested for the NMO disease spectrum;⁷ intravenous corticosteroids are recommended for acute presentation and plasmapheresis is suggested if disease is refractory to steroids. A diagnosis of LETM as part of the NMO disease spectrum is based on MRI spine indicating a lesion extending over three or more spinal cord segments, MRI brain showing no features of MS and NMO-IgG seropositivity.⁸

The NMO-IgG antibody binds to the aquaporin-4 water channel protein, which resides on astrocyte foot processes.⁹

Learning points

- ▶ LETM is a neurological condition characterised by a contiguous inflammatory lesion of the spinal cord.
- ▶ LETM is often associated with the autoimmune central nervous system disease NMO and rarely associated with MS.
- ▶ Morbidity and mortality levels associated with LETM are variable.
- ▶ Early serological testing for NMO-IgG antibodies in suspected LETM is important for prompt diagnosis, indicating prognosis and establishing appropriate therapy to reduce the risk of future relapses or disease progression.

The pathogenesis of NMO is suggested to involve binding of autoantibodies to aquaporin-4 leading to blood brain barrier hyperpermeability, perivascular inflammation, astrocyte injury and inflammation culminating in spinal cord and/or optic nerve demyelination and cavitation.

A previous report has suggested NMO-IgG antibodies are present in approximately 38% of LETM patients following an initial attack, and is associated with >60% risk of relapse within 1 year and development of optic neuritis.³ Thus all suspected cases of LETM should undergo NMO-IgG serological testing, as NMO relapses are associated with increased disability and morbidity.¹⁰

In conclusion, we have highlighted two cases of LETM with different disease severity and NMO-IgG status and described the importance of the NMO-IgG antibody as diagnostic and prognostic biomarker for LETM.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Pearce JM.** Neuromyelitis optica. *Spinal Cord* 2005;**43**:631–4.
2. **Wingerchuk DM, Lennon VA, Lucchinetti CF, et al.** The spectrum of neuromyelitis optica. *Lancet Neurol* 2007;**6**:805–15.
3. **Weinshenker BG, Wingerchuk DM, Vukusic S, et al.** Neuromyelitis optica IgG predicts relapse after longitudinally extensive transverse myelitis. *Ann Neurol* 2006;**59**:566–9.
4. **Bizzoco E, Lolli F, Repice AM, et al.** Prevalence of neuromyelitis optica spectrum disorder and phenotype distribution. *J Neurol* 2009;**256**:1891–8.
5. **Bot JC, Barkhof F, Polman CH, et al.** Spinal cord abnormalities in recently diagnosed MS patients: added value of spinal MRI examination. *Neurology* 2004;**62**:226–33.
6. **Nandhagopal R, Al-Asmi A, Gujjar AR.** Neuromyelitis optica: an overview. *Postgrad Med J* 2010;**86**:153–9.
7. **Pandit L.** Transverse myelitis spectrum disorders. *Neural India* 2009;**57**:126–33.
8. **Wingerchuk DM, Lennon VA, Pittock SJ, et al.** Revised diagnostic criteria for neuromyelitis optica. *Neurology* 2006;**66**:1485–9.
9. **Saikali P, Cayrol R, Vincent T.** Anti-aquaporin-4 auto-antibodies orchestrate the pathogenesis in neuromyelitis optica. *Autoimmun Rev* 2009;**9**:132–5.
10. **Argyriou AA, Makris N.** Neuromyelitis optica: a distinct demyelinating disease of the central nervous system. *Acta Neurol Scand* 2008;**118**:209–17.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Nightingale H, Witherick J, Wilkins A. Diagnosis of longitudinally extensive transverse myelitis.
BMJ Case Reports 2011;10.1136/bcr.10.2010.3444, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow