Unusual presentation of more common disease/injury

An unusual case of pneumomastia: mammographic appearances

David Henry Green

Wirral Breast Screening Unit, Wirral Teaching Hospital, Bebington, UK

Correspondence to Dr. David Henry Green, davidhenrygreen@hotmail.com**Summary**

Gas in the breast parenchyma (pneumomastia) is a common occurrence after breast interventional procedures but is a rare phenomenon otherwise and not previously recorded mammographically. We present a case of pneumomastia presenting after laparoscopic abdominal surgery. The juxtaposition of laparoscopic surgery with a mammogram appointment is likely to be a rare event; however, recognition of this entity will avoid potential diagnostic pitfalls and allow the radiologist to alert the examining breast surgeon to the possible presence of subcutaneous crepitation.

BACKGROUND

Gas in the breast parenchyma (pneumomastia) is a common occurrence after breast interventional procedures but is a rare phenomenon otherwise and not previously recorded mammographically.

CASE PRESENTATION

A 66-year-old woman attended the breast clinic with symptoms of a lump in the lower inner quadrant of the left breast. Bilateral mammography was performed and revealed an unusual pattern of parenchymal gas shadows in the left breast (figure 1). Clinical examination was normal but on direct questioning the patient indicated that she had undergone laparoscopic pelvic surgery 7 days earlier for right oophorectomy and adhesion lysis. A diagnosis of pneumomastia secondary to subcutaneous emphysema was made.

DIFFERENTIAL DIAGNOSIS

The differential diagnosis of pneumomastia is:

- i Interventional breast procedure
- ii Penetrating
- iii Subcutaneous emphysema
- iv Anaerobic infection

DISCUSSION

Pneumomastia seen on a mammogram can result from diagnostic breast procedures or pathology. In the past, CO₂ gas has been used as a contrast agent by injecting it into the fascial spaces of the breast; the resulting pneumomammogram was said to improve the diagnostic value of the examination.¹ Pneumocystography may still occasionally be useful as an adjunct to ultrasound in the further evaluation of non-simple breast cysts,² while iodinated contrast ductography has largely supplanted pneumoductography in the investigation of nipple discharge. Needle biopsy commonly results in the introduction of air

into the breast tissue. In all of the above cases, the resulting gas shadows will localise to an anatomical structure. Anaerobic infection of the breast, while a rare cause of pneumomastia, is unlikely to cause diagnostic confusion since the clinical presentation is different and mammography is contraindicated.³

If an interventional breast procedure has been excluded, pneumomastia will be the result of subcutaneous emphysema tracking from a more distant source. It is recognised that subcutaneous emphysema is a rare complication of laparoscopic surgery⁴ as a result of the large volumes of high pressure CO₂ used to induce pneumoperitoneum. In this particular case, the presence of anterior abdominal wall adhesions between omentum and peritoneum had necessitated a Palmer's point entry (ie, a laparoscopic porthole 3 cm below the left costal margin in the midclavicular line). This allows the surgeon safe access to the pelvis.⁵ It seems reasonable to assume that the proximity of the porthole to the left breast explains why, in this case, air tracked to the left breast and not the right. The juxtaposition of laparoscopic surgery with a mammogram appointment is likely to be a rare event; however, recognition of this entity will avoid potential diagnostic pitfalls and allow the radiologist to alert the examining breast surgeon to the possible presence of subcutaneous crepitation.

Learning points

- Pneumomastia is common after percutaneous interventional breast procedures but rare otherwise.
- Pneumomastia can occur after laparoscopic surgery.
- Pneumomastia produces characteristic changes on mammography.

Competing interests None.

Patient consent Obtained.

Figure 1 (A) Mediolateral oblique, (B) craniocaudal and (C) magnified coned craniocaudal mammograms of the left breast showing an incidental finding of parenchymal gas (outlined by arrows).

REFERENCES

1. **Rozin DL.** [Pneumomammography in the diagnosis of breast diseases]. *Vopr Onkol* 1975;**21**:6–14.
2. **Fajardo LL,** Jackson VP, Hunter TB. Interventional procedures in diseases of the breast: needle biopsy, pneumocystography, and galactography. *AJR Am J Roentgenol* 1992;**158**:1231–8.
3. **Delotte J,** Karimjee BS, Cua E, *et al.* Gas gangrene of the breast: management of a potential life-threatening infection. *Arch Gynecol Obstet* 2009;**279**:79–81.
4. **Badawy SZA,** Dalpe C, Romano D. Pneumomediastinum and subcutaneous emphysema during carbon dioxide laser laparoscopy for tuboplasty. *J Gynecol Surg* 2002;**18**:159–163.
5. **Aust TR,** Kayani SI, Rowlands DJ. Direct optical entry through Palmer's point: a new technique for those at risk of entry-related trauma at laparoscopy. *Gynecol Surg* 2009;**7**:315–17.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Green DH. An unusual case of pneumomastia: mammographic appearances. *BMJ Case Reports* 2011;10.1136/bcr.03.2010.2852, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow