Published in final edited form as:

Org Lett. 2011 April 1; 13(7): 1742–1745. doi:10.1021/ol200256a.

Mechanistic Investigation of the Enantioselective Intramolecular Stetter Reaction: Proton Transfer is the First Irreversible Step

Jennifer L. Moore[§], Anthony P. Silvestri, Javier Read de Alaniz[‡], Daniel A. DiRocco, and Tomislav Rovis^{*}

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523

Abstract

A study on the mechanism of the asymmetric intramolecular Stetter reaction is reported. This investigation includes the determination of the rate law, kinetic isotope effects and competition experiments. The reaction was found to be first order in aldehyde and azolium catalyst or free carbene. A primary kinetic isotope effect was found for the proton of the aldehyde. Taken together with a series of competition experiments, these results suggest that proton transfer from the tetrahedral intermediate formed upon nucleophilic attack of the carbene onto the aldehyde is the first irreversible step.

The seminal example of the reversal of functional group polarity, the benzoin reaction, dates to 1832, when Wöhler and Liebig reported that cyanide catalyzes the formation of benzoin from two equivalents of benzaldehyde (eq. 1).^{1,2} In 1943, Ukai *et al* showed that thiazolium salts catalyze the homodimerization of aldehydes in the presence of base.³ A related *Umpolung*⁴ transformation is the Stetter reaction, the conjugate addition of the aldehyde into a Michael acceptor.⁵ Utilizing cyanide or thiazolylidene carbenes as catalysts, Stetter demonstrated that a variety of aromatic and aliphatic aldehydes are competent nucleophilic coupling partners with a wide range of α , β -unsaturated ketones, esters, and nitriles.⁶ The ability to bring two different electrophilic partners together and form a new carbon—carbon bond enhances the potential utility of this transformation.

Our group has developed chiral triazolynylidene carbenes and precursors, **1-3**, for a variety of carbene mediated transformations.⁷ We have shown that the carbenes derived from **1** and **2** are capable of inducing the cyclization of aromatic and aliphatic aldehydes to α , β -unsaturated esters, ketones, thioesters, amides, aldehydes, and nitriles.⁸ More recently, we

rovis@lamar.colostate.edu.

[§]Current address: Natural Science Division, Pepperdine University, Malibu, CA.

[‡]Current address: Department of Chemistry and Biochemistry, University of California, Santa Barbara, CA.

Supporting Information Available: Experimental procedures, kinetic and competition experiment data are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

have extended this reaction to preliminary reports of asymmetric intermolecular reactions.^{9,10} Nevertheless, limitations remain.

Mechanistic insight into organocatalytic reactions is important for the development of general transformations.¹¹ To the best of our knowledge a detailed study probing the mechanism of the Stetter reaction has not been reported. In the absence of such a study the working model of the Stetter reaction is based on the Breslow mechanism for the thiamin-catalyzed benzoin reaction.^{12,13} The mechanism is closely related to Lapworth's mechanism for cyanide anion catalyzed benzoin reaction.¹⁴ As with the cyanide catalyzed benzoin reaction, the thiazolinyldene catalyzed reaction is reversible.¹⁵

The proposed catalytic cycle is as follows: the carbene **I** (Scheme 1), formed *in situ* by base deprotonation of the corresponding azolium salt, adds to the aldehyde to form **II**. A proton transfer event generates acyl anion equivalent **III**, termed the nucleophilic alkene or Breslow intermediate. Subsequent addition into the Michael acceptor forms a new carbon—carbon bond to generate **IV**. A second proton transfer event then provides **V**. Finally, collapse of this tetrahedral intermediate **V** to form Stetter product is accompanied by liberation of the active catalyst. As we strive to understand differences in catalysts and continue to work toward the development of the enantioselective intermolecular reactions, we believe that the results from a detailed mechanistic study may provide insight toward the rational attainment of these goals. Herein we report a series of mechanistic experiments that shed light on the nuances that govern reactivity in the intramolecular Stetter reaction.

(1)

As salicylaldehyde derived aldehyde **4** is used as a benchmark to measure the efficiency and selectivity of newly developed catalysts for the Stetter reaction, it was chosen as the substrate for this study, eq 1. Under standard reaction conditions, aldehyde **4** is subjected to 20 mol % **2a** and 20 mol % KHMDS in toluene (0.025 M) at 0 °C; the observed rate of the reaction is $2.65 \times 10^{-3} \text{M}^{-1} \text{s}^{-1}$. Gas chromatography was utilized for the analysis of cyclized product **5** by using 4,4'-di-*tert*-butyl biphenyl (DBB) as an internal standard (t_R 4.8 min) and monitoring the disappearance of aldehyde **4** (t_R 2.0 min) and concurrent appearance of keto-ester **5** (t_R 2.9 min). Standard kinetic analysis using the conversion of aldehyde **4** to keto-ester **5** exhibits a first order dependence as a function of aldehyde concentration versus time over four half lives.¹⁶

The catalyst dependence was found to be first order, determined by varying the concentration of catalyst from 0.0025M to 0.0100M. These experiments establish a second order rate law (eq 1).¹⁷

The ²H kinetic isotope effect (KIE) study was conducted under standard reaction conditions from triplicate runs with **4** and its deuterated isotopologue (ArCDO), and the k_H/k_D was found to be 2.62 (Scheme 2). These experiments suggest that proton transfer is turnover limiting.

In order to shed further light on the mechanism, we conducted a series of competition experiments. We reasoned that the sterics and electronics of the Michael acceptor should have a profound role on the reaction if the first irreversible step is the second proton transfer event (**IV** to **V**) but a negligible role if it is the first (**II** to **III**, Scheme 1). Equimolar amounts of aldehydes **4** and **6** were subjected to the reaction in the presence of an internal standard (DBB) and the reaction monitored by GC. We found that the two substrates are consumed at nearly the same rate ($k_{\text{rel}} = 1.25$, k_4/k_6 , eq 2). This result suggests that the initial proton transfer is the first irreversible step.

(2)

We further conducted a series of competition experiments to investigate the role of aldehyde electronics on the reaction. A more electron-deficient aldehyde is consumed ~10.1 times faster than the parent substrate **4**, Scheme 3a. Conversely, the more electron-rich substrate **10** proved more sluggish than **4** providing a k_4/k_{10} of ~7.7.

The above observations are all consistent with the initial proton transfer (**II** to **III**, Scheme 1) being the first irreversible step.¹⁸ The more electron-deficient substrate **8** bears a methine of increased acidity relative to the parent substrate. Similarly, substrate **10** bearing the electron-releasing *para*-methoxy group thus has a less acidic methine in its corresponding intermediate **II**, leading to a slower reaction.

What remains is to determine the mechanism of the proton transfer. A direct 1,2-proton shift is symmetry forbidden and unlikely to occur.¹⁹ In a DFT study of a model Stetter reaction mechanism, Yates has suggested that the proton transfer event must proceed intermolecularly, most likely via a second zwitterionic intermediate (similar to **II**, Scheme 1).²⁰ However, if this were the mechanism for the proton transfer in our reaction, kinetics should be second order in catalyst and substrate. We have further found that the reaction involving the free carbene **3a** is similar in all respects to the **2a**/KHMDS system suggesting HMDS and KBF₄ play no role in the reaction.²¹

The most likely scenario that remains is that there is an agent of proton transfer within the molecule, either on the catalyst or the substrate. We envisioned two scenarios, one involving the ethereal oxygen linking the aldehyde and Michael acceptor and the second involving the aryl ring on the azolium precatalyst (Scheme 4). As a test of the latter hypothesis, we note that subjection of deuterioaldehyde **4-D** to N-phenyl triazolium precatalyst under typical reaction conditions results in no observed deuteration in the catalyst architecture. A kinetic isotope effect should result in some deuteration of the ortho position on the catalyst aryl ring (compare **XIII** in Scheme 4, H vs D). The potential role of substrate in accelerating the proton transfer event was also evaluated. One can imagine that the proximity of the ether linker leads to deprotonation of the methine, and generation of the derived oxonium (**X** in Scheme 4). This intermediate is presumably very short lived, undergoing abstraction by the alkoxide generating **XI**, analogous to the nucleophilic alkene **III** in Scheme 1. As a test of

this hypothesis, we conducted a competition experiment between ether-linked aldehyde **4** and substrate **12** bearing a methylene linker. Experiments shown in Scheme 3 suggest that the more electrophilic substrate **12** (bearing the more acidic methine – contrast **4** and **10**, Scheme 3b) should be consumed faster if the oxygen linker has no role in the mechanism. We found that the oxygen-linked substrate **4** is consumed in high preference over the methylene-linked substrate **12** implicating the oxygen in the proton transfer event.²²

In summary, we have provided convincing evidence that proton transfer is the first irreversible step in the triazolinyldine carbene catalyzed asymmetric intramolecular Stetter reaction. This finding will have an impact on the design of future catalysts for the Stetter reaction. Furthermore, given the importance of generating the enamine or nucleophilic alkene in other NHC-catalyzed reactions of aldehydes, this mechanistic nuance should also have a bearing on those areas.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Science Foundation and NIGMS (GM72586) for support, and Roche and Amgen for unrestricted support. J. R. and J. L. M. thank the National Institutes of Health (Ruth L. Kirschstein Minority Pre-doctoral fellowship). J. R. also thanks Colorado PEAKS AGEF (graduate fellowship). We thank Profs. Jeffrey B. Johnson (Hope College) and Daniel A. Singleton (Texas A&M) for many helpful discussions.

References

1. Woehler F, Liebig J. *Ann Pharm.* 1903; 3:249.
2. For reviews, see: (a) Moore JL, Rovis T. *Top Curr Chem.* 2009; 290:77. (b) Read de Alaniz J, Rovis T. *Synlett.* 2009; 8:1189. [PubMed: 20585467] (c) Enders D, Niemeier O, Henseler A. *Chem Rev.* 2007; 107:5606. [PubMed: 17956132]
3. Ukai T, Tanaka R, Dokawa T. *J Pharm Soc Jpn.* 1943; 63:296.
4. Seebach D. *Angew Chem.* 1979; 91:259.
5. Stetter H. *Angew Chem, Int Ed Engl.* 1976; 15:639.
6. Stetter H, Kuhlmann H. *Org React.* 1991; 40:407.
7. (a) Read de Alaniz J, Rovis T. *Synlett.* 2009; 1189. [PubMed: 20585467] (b) Moore JL, Rovis T. *Top Curr Chem.* 2009; 290:77.
8. (a) Kerr MS, Read de Alaniz J, Rovis T. *J Am Chem Soc.* 2002; 124:10298. [PubMed: 12197730] (b) Kerr MS, Rovis T. *J Am Chem Soc.* 2004; 126:8876. [PubMed: 15264801] (c) Read de Alaniz J, Rovis T. *J Am Chem Soc.* 2005; 127:6284. [PubMed: 15853335] (d) Liu Q, Rovis T. *J Am Chem Soc.* 2006; 128:2552. [PubMed: 16492036] (e) Read de Alaniz J, Kerr MS, Moore JL, Rovis T. *J Org Chem.* 2008; 73:2033. [PubMed: 18302407]
9. (a) Liu Q, Perreault S, Rovis T. *J Am Chem Soc.* 2008; 130:14066. [PubMed: 18834123] (b) Liu Q, Rovis T. *Org Lett.* 2009; 11:2856. [PubMed: 19507841] (c) DiRocco DA, Dalton DM, Oberg KM, Rovis T. *J Am Chem Soc.* 2009; 131:10872. [PubMed: 19722669]
10. Contributions from others: (a) Enders D, Han J, Henseler A. *Chem Commun.* 2008:3989. (b) Enders D, Han J. *Synthesis.* 2008:3864. (c) Jousseume T, Wurz NE, Glorius F. *Angew Chem Int Ed.* 2011; 50:1410. For an enzyme-catalyzed asymmetric Stetter, see: (d) Dresen C, Richter M, Pohl M, Lüdeke S, Müller M. *Angew Chem Int Ed.* 2010; 49:6600.
11. (a) Kunz RK, MacMillan DWC. *J Am Chem Soc.* 2005; 127:3240. [PubMed: 15755116] (b) Franzén J, Marigo M, Fielenback D, Wabnitz TC, Kjærsgaard A, Jørgensen KA. *J Am Chem Soc.* 2005; 127:18296. [PubMed: 16366584] (c) Halland N, Lie MA, Kjærsgaard A, Marigo M, Schjøtt B, Jørgensen KA. *Chem Eur J.* 2005; 11:7083. (d) Grossman RB, Comesse S, Rasne RM, Hattori K, DeLong MN. *J Org Chem.* 2003; 68:871. [PubMed: 12558409] (e) Lemay M, Ogilvie WW. *Org Lett.* 2005; 7:4141. [PubMed: 16146372] (f) Armstrong A. *Angew Chem Int Ed.* 2004; 43:1460.

- (g) Bulman Page PC, Barros D, Buckley BR, Marples BA. *Tetrahedron: Asymmetry*. 2005; 16:3488.
12. (a) Breslow RJ. *J Am Chem Soc*. 1958; 80:3719. (b) Knight RL, Leeper FJ. *Tetrahedron Lett*. 1997; 38:3611. (c) Knight RL, Leeper FJ. *J Chem Soc, Perkin Trans*. 1998; 1:1891. (d) Dudding T, Houk KN. *Proc Natl Acad Sci USA*. 2004; 101:5770. [PubMed: 15079058] (e) Sheehan JC, Hunneman DH. *J Am Chem Soc*. 1966; 88:3666. (f) White MJ, Leeper FJ. *J Org Chem*. 2001; 66:5124. [PubMed: 11463265] (g) Mahathananchai J, Zheng P, Bode JW. *Angew Chem Int Ed*. 2011; 50:1673.
13. Lopez-Calahorra and co-workers proposed an alternative mechanism involving thiazolium dimer **VI** with the reaction occurring through nucleophilic attack of this species on benzaldehyde to yield **VII**, followed by formation of intermediate type **VIII**, lacking stabilization of the carbanion. See, for example: Lopez-Calahorra F, Rubires R. *Tetrahedron*. 1995; 51:9713. For a counterpoint, see: Breslow R, Schmuck C. *Tetrahedron Lett*. 1996; 37:8241.

14. Lapworth A. *J Chem Soc*. 1903; 83:995.
15. Buck JS, Ide WS. *J Am Chem Soc*. 1931; 53:2350.
16. See SI for the details of the kinetic experiments and the corresponding graphs.
17. The same kinetic analysis was conducted on catalyst **3a** showing little difference in the corresponding catalytic cycle.
18. This is consistent with Berkessel's observation that the catalyst resting state in the triazolyldene-catalyzed benzoin in THF is a dioxolane resulting from the tetrahedral alkoxy intermediate analogous to **II** adding another molecule of aldehyde in a reversible process. Thus, if proton transfer to generate **III** is slow, the alkoxide would prefer adding to an aldehyde to avoid the charge on oxygen in the absence of solvation. See: Berkessel A, Elfert S, Etzenbach-Effers K, Teles JH. *Angew Chem Int Ed*. 2010; 49:7120.
19. 1,2-proton shift is a symmetry forbidden transformation (Kemp DS. *J Org Chem*. 1971; 36:202. and references therein) but has been calculated to have a ~29 kcal/mol barrier (thiazolyldine) and ~51 kcal/mol barrier (cyanide) in the formation of the acyl anion equivalent from formaldehyde; see: Goldfuss B, Schumacher M. *J Mol Model*. 2006; 12:591. [PubMed: 16258739]
20. Hawkes KJ, Yates BF. *Eur J Org Chem*. 2008:5563.
21. The protocol for generating the free carbene **3a** involves removal of volatiles including HMDS. Added HMDS did not appreciably affect the rate of the reaction. Separation of the insoluble KBF₄ shows no appreciable differences in reactivity. Furthermore, in situ ¹⁹F NMR of the reaction medium shows no fluorine signals suggesting KBF₄ is completely insoluble under these conditions.
22. That substrate **12** still undergoes reaction means that there must be another mechanism available for proton transfer, presumably either via Hypothesis II above or involving another molecule of catalyst or intermediate, as argued by Yates (see ref. 20).

Figure 1.
Catalysts used in the Stetter reaction

Scheme 1.
Proposed General Mechanism for the Stetter Reaction

Scheme 2.
Kinetic Isotope Effect Studies

Scheme 3.
Aldehyde Competition Experiments

• Hypothesis 1 - substrate shuttles proton:

• Hypothesis 2 - catalyst shuttles proton:

Scheme 4.
Probing the Mechanism of the 1,2-Proton Shift

Scheme 5.
Competition Experiment