Published in final edited form as:

J Am Diet Assoc. 2011 April ; 111(4): 532–541. doi:10.1016/j.jada.2011.01.017.

Nutrient Intake and Anemia Risk in the WHI Observational Study

Cynthia A. Thomson, PhD, RD,

Associate Professor Nutritional Sciences, University of Arizona, Shantz Bldg -1177 E 4th Street, Tucson, AZ 85724, (520) 626-1565, FAX: (520) 621-9446

Jeffrey Stanaway, MPH,

University of Washington, Department of Epidemiology, 1959 NE Pacific Street, Health Sciences Building F-262, Box 357236, Seattle, WA 98195-7236, (206) 919-9001

Marian L. Neuhouser, PhD, RD,

Affiliate Associate Professor, Epidemiology, Fred Hutchinson Cancer Research Center, 1100 Fairview Avenue, North, MP-702, PO Box 19024, Seattle, WA 98109, (206) 667-4797, FAX: (206) 667-7850

Linda G. Snetselaar, RD, PhD, LD,

Interim DEO of Community and Behavioral Health, Professor Department of Epidemiology, College of Public Health, University of Iowa, E225-A General Hospital, Iowa City, IA 52242, (319) 384-5011, FAX: (319) 384-5385

Marcia L. Stefanick, PhD,

Professor of Medicine, Stanford Prevention Research Center, Medical School Office Building, 251 Campus Drive Room X308, Stanford, CA 94305, (650) 725-5041, FAX: (650) 725-6247

Leslie Arendell, MS, and

Mel and Enid Zuckerman College of Public Health, University of Arizona, PO Box 245203, Tucson, AZ 85724, 520-626-9621, FAX: (520) 626-2767

Zhao Chen, PhD, MPH

Professor, Division Director, Epidemiology/Biostats, Mel and Enid Zuckerman College of Public Health, University of Arizona, PO Box 245203, Tucson, AZ 85724, (520) 626-9011, FAX: (520) 626-2767

Cynthia A. Thomson: cthomson@u.arizona.edu; Jeffrey Stanaway: Stanaway@uw.edu; Marian L. Neuhouser: mneuhaus@fhcrc.org; Linda G. Snetselaar: linda-snetselaar@uiowa.edu; Marcia L. Stefanick: Stefanick@stanford.edu; Leslie Arendell: lesliea@email.arizona.edu; Zhao Chen: zchen@email.arizona.edu

Abstract

Background—Nutritional anemia among post-menopausal women is preventable; recent data on prevalence are limited.

Objective—To investigate the association between nutrient intakes and anemia prevalence, in relation to both incidence and persistence, in a longitudinal sample of post-menopausal women. We hypothesized that anemia prevalence, incidence and persistence would be greater among women reporting lower intake of B12, folate and iron.

Design—Prospective cohort analysis.

Correspondence to: Cynthia A. Thomson, cthomson@u.arizona.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Participants/setting—Observational Cohort of the Women’s Health Initiative(WHI-OS) including 93,676 postmenopausal women, age 50 to 79 years, were recruited across the United States at 40 clinical study sites. Women were enrolled between 1993 and 1998; data collection for these analyses continued through 2000.

Main outcome measures—Anemia was defined as a blood hemoglobin concentration of <12.0 mg/dL. Persistent anemia was defined as anemia present at each measurement time point. Diet was assessed by food frequency questionnaire for iron, folate, B12, red meat and cold breakfast cereal; inadequacies were based on dietary reference intakes for women over age 50 years.

Statistical analysis—Descriptive statistics (mean and standard deviation) were used to characterize the population demographics, anemia rates and diet. Unconditional logistic regression was used to investigate associations between diet and incident and persistent anemia. Associations are presented as odds ratio (OR) and 95% confidence intervals (CI).

Results—Anemia was identified in 3,979 women or 5.5% of the cohort. Inadequate intakes of multiple anemia-associated nutrients were less frequent in non-Hispanic whites (7.4%) than other race/ethnic groups (inadequacies demonstrated in 14.6 to 16.3% of sample). Age, body mass index and smoking were associated with anemia. Women with anemia reported lower intakes of energy, protein, folate, B12, iron, vitamin C and red meat. Multiple (more than a single nutrient) dietary deficiencies were associated with a 21% greater risk of persistent anemia (OR-1.21, 95% CI: 1.05–1.41) and three deficiencies resulted in a 44% increase in risk for persistent anemia (OR-1.44, 95% CI: 1.20–1.73).

Conclusion—Inadequate nutrient intake, a modifiable condition, is associated with greater risk for anemia in post-menopausal women participating in the WHI-OS. Efforts to identify and update incidence estimates for anemia-associated nutrient deficiencies in aging women should be undertaken.

Keywords

anemia; diet; aging

Introduction

Anemia is a relatively common health problem in the older population of the U.S. and is associated with increased mortality (1) and more frequent hospitalization (2). The World Health Organization (WHO) defines anemia as a hemoglobin concentration of less than 13.0 g/dL in men, and less than 12.0 g/dL in women (3). Using this criteria and the National Health and Nutrition Examination Survey (NHANES) III dataset, researchers found anemia to be present in over 10% of those aged 65 and over; this prevalence parallels that seen in other aging adults (1,4–7).

Nutritional anemia includes those associated with prolonged inadequate intake of folate, vitamin B12, iron, protein and vitamin C (8,9). Elderly women may be at an increased risk for inadequate micronutrient intake (10). Inadequate nutrient intake is commonly the result of health symptoms such as poor dentition (reduced red meat consumption resulting in reduced iron and vitamin B12 intake), reduced appetite (resulting in general reduction in all nutrients) and reduced tolerance of milk products (a primary source of vitamin B12). Additionally, a loss of intrinsic factor with aging is a well-characterized indirect cause of nutritional anemia.

Anemia of nutritional etiology is generally, but not always (as may be in the case in obese individuals with underlying inflammation and alterations in iron metabolism) (11), responsive to diet modification. Achieving adequacy in intake of anemia-associated

nutrients could either reduce risk for or correct prevalence and severity of nutritional anemia. Further, correction of anemia would in turn reduce or eliminate the associated morbidity (12,13).

In addition to anemia of nutritional etiology, non-nutritional anemia is common in aging women, including those associated with inflammation leading to elevated pro-inflammatory cytokines decreasing iron bioavailability inhibiting the hematopoietic response (14,15). Chronic illnesses, and anemia of unknown etiology (16,17) as well as blood loss (18) are also thought to be the primary causes of microcytic anemia in aging populations.

Having a better understanding of the dietary factors associated with anemia in an ethnically diverse population of post-menopausal women is an important first step for supporting such preventive and therapeutic approaches. The Women's Health Initiative Observational Study (WHI-OS) cohort affords an important opportunity to re-examine anemia in aging females with a focus on potential dietary etiologies for reduced hemoglobin concentrations. Further, the sample population provides specific information across diverse ethnicities and races. Baseline dietary intake data are now available for the 93,676 WHI-OS participants to evaluate these associations. The primary objective of this paper is to update available evidence investigating the association between self-reported nutrient intakes and anemia prevalence; secondarily, the paper prospectively investigates the risk of incident and persistent anemia associated with dietary intake. The hypothesis for this investigation was that the prevalence, incidence and persistence of anemia would be greater among postmenopausal women reporting low vitamin B12, folate or iron intake at time of study enrollment, relative to women with adequate intake of these nutrients (as defined by Dietary Reference Intakes), after controlling for confounding factors. Further, it was hypothesized that a greater number of dietary inadequacies of these nutrients would be associated with a greater risk for incident and persistent anemia at year 3.

Methods

Study Population

This study includes post-menopausal women enrolled in the WHI-OS. This observational cohort was recruited across 40 clinical research sites nationally from 1993–1997 (19). The majority of women were recruited through mail or newspaper advertisements; although a wide range of recruitment approaches were employed. Each recruitment site obtained Institutional Review Board approval and all participants provided written informed consent. The consent process was completed between the participant and local study coordinator during an on-site clinic visit. The WHI-OS population evolved from the screening cohort of women found to be either ineligible or uninterested in the clinical trials portion of WHI in addition to women recruited explicitly for the WHI-OS. At the time of enrollment women ranged in age from 50 to 79 years. Women were excluded from participation if they had a medical condition with less than 3 years life expectancy, if they were currently enrolled in another intervention trial, or if health habits or diagnosis diminished the women's ability to provide informed consent or study data (e.g. alcoholism, dementia, etc). Given the well-documented associations between non-nutrient-deficiency anemia and certain chronic diseases (20–24), participants with a history of myocardial infarction, congestive heart failure, rheumatoid arthritis, lupus, and any cancer were excluded from these analyses. Exclusion for renal disease was not necessary, as no participants were enrolled in the WHI with renal disease. Participants lacking baseline hemoglobin data were also excluded.

Participants without baseline dietary data and those reporting extreme energy intakes were identified and excluded using a two-part scheme: all extreme outliers (those reporting less than 300 kcal or more than 7000 kcal daily) were excluded; moderate outliers were excluded

when their reported energy intakes changed by more than 50% between baseline and year three (those reporting 300 to 499.99 kcal with >50% change, or 5000 to 7000 kcal with >50% change). Cutoff points were based on natural breaks in the data, as determined by visual inspection of energy distribution plots.

This prospective analysis required the availability of year-three hemoglobin data. To ensure consistency between analyses and simplicity of design, the same WHI-OS participants were included in the cross-sectional analysis of baseline dietary intake and anemia prevalence.

Anemia Assessment

In this study, anemia was defined according to the World Health Organization (WHO) definition as a baseline hemoglobin concentration less than 12.0 mg/dL (3). Incident anemia (newly diagnosed) was defined by a reduction in hemoglobin from values greater than or equal to 12.0 mg/dL at baseline to values less than 12.0 mg/dL at year three. Women were alternately categorized as having persistent anemia (hemoglobin less than 12.0 mg/dL at baseline and year 3) or resolved anemia in cases where hemoglobin increased to greater than 12.0 mg/dL from a baseline value below 12.0 mg/dL. Women could also be categorized as not having anemia (incident or persistent). Hemoglobin concentrations were assessed as part of the complete blood count (CBC) measured at baseline and again at the 3 year clinic visit from 12 hour fasting blood samples.

Dietary Intake

Dietary intake was collected from study participants at baseline and year 3 using a Food Frequency Questionnaire (FFQ) developed and validated by the Fred Hutchinson Cancer Research Center (FHCRC). This questionnaire was specifically designed for the Women's Health Initiative to include foods representative of the national food supply (25) and to capture regional and ethnic food selections. The questionnaire includes 122 food lines and asks respondents to indicate the portion consumed as well as the frequency of consumption using a Likert-type scale referencing food intake patterns over the previous 3-month period. Nutrient intake was derived from an FHCRC-designed specialty software program that utilizes the Nutrition Data System for Research (NDS-R) food and nutrient database (University of Minnesota Nutrition Coordinating Center) (26). A validation study suggested that the energy-adjusted anemia-associated nutrient intake correlations between FFQ and 8-day food records collected from WHI study participants were: 0.2 for vitamin B12, 0.5 for folate, 0.3 for vitamin C and 0.6 for iron (26). WHI-OS participants completed the FFQ at baseline during the screening period and each questionnaire was reviewed for completeness by the WHI site coordinator prior to being forwarded to the WHI Coordinating Center for analysis. It is important to note that the baseline food frequency questionnaire data were collected between 1993 and 1998 and reflect folic acid intake prior to the mandatory fortification date; data were analyzed using annual USDA database estimates for folic acid and folate as well as other nutrients.

Dietary exposure variables of interest included vitamin B12, folate, iron, vitamin C, animal protein intake, vegetable protein intake, red meat intake and fortified cereal intake (iron, folic acid). Dietary inadequacy was defined as below dietary reference intakes for individual nutrients based on intake recommendations for women over 50 years of age. Use of supplemental nutrients including multivitamins, B-complex vitamins and individual nutrient supplementation including iron, vitamin B12 and folic acid were also evaluated. In addition to evaluating individual nutrient intake, analyses were performed for combined nutrient inadequacies. In these models, each intake for an individual nutrient below the RDA was summed to determine the association between quantity of individual nutrient inadequacy and anemia.

Statistical Analysis

Baseline characteristics of the study participants are presented as mean (standard deviation (SD)) or frequencies (%) by anemia group. To compare women with baseline anemia versus those without baseline anemia, t-tests and chi-square tests were conducted for continuous and categorical variables, respectively. Adjusted and unadjusted mean nutrient intakes were calculated by anemia status; adjusted means were calculated using linear prediction, and means were adjusted for age category (50–59, 60–69, and 70–79), ethnicity, smoking status (non-smoker, former smoker, or current smoker) and income category (<\$10,000, \$10,000–19,999, \$20,000–34,999, \$35,000–49,999, \$50,000–74,999, \$75,000–99,999, \$100,000–149,999, \$150,000+).

Unconditional logistic regression was used to investigate the association of inadequate nutrient intake, servings of selected food groups and the number of anemia-associated dietary micronutrient inadequacies at baseline with both incident and persistent anemia. In investigating the association between the anemia-associated dietary micronutrient inadequacies, the data were assessed two ways: first, the number of inadequacies was treated as a categorical variable to allow for nonlinearity in the association with anemia; second, the number of inadequacies was treated as a continuous variable to test for trend if the categorical analysis suggested a possible linear association. Both crude and adjusted logistic regression models were conducted: adjusted models included indicator variables for category of age (50–59, 60–69, and 70–79), race/ethnicity (American Indian or Alaskan native, Asian or Pacific islander, Black or African-American, Hispanic/Latino, White (non-Hispanic), and other), smoking status (non-smoker, former smoker or current smoker) and income (<\$10,000, \$10,000–19,999, \$20,000–34,999, \$35,000–49,999, \$50,000–74,999, \$75,000–99,999, \$100,000–149,999, and \$150,000+) as well as body mass index (BMI) as a continuous variable. Confounders were selected *a priori* based on plausibility and previous literature and included BMI, age range, race/ethnicity, smoking and income.

For each of the five anemia-related nutrients (folate, iron, vitamin B12, vitamin C, and protein), multiplicative interaction terms between race/ethnicity and nutrient inadequacy were created to assess whether the association between anemia and a given nutrient varies by race/ethnicity. Specifically, we used the likelihood ratio test to determine if the addition of this interaction term significantly improved any of the adjusted models. Since this required testing for an interaction with ethnicity in five separate models (one for each of the anemia-related nutrients), the Bonferroni adjustment was used to account for multiple comparisons; thus, the level of significance required in any model was equal to α/n , where α was set to 0.05 and n equaled 5, the number of tests. All analyses were performed with Stata statistical software (version 10.1 and 11.1, Statacorp, College Station, TX).

Results

Of the 93,676 women enrolled in the WHI-OS; 18,733 were excluded from the analytical cohort due to previous diagnosis of chronic disease; 1,484 were excluded due to extreme energy reporting; 626 were excluded due to missing hemoglobin or baseline diet data. Of the 72,833 women in the analytical cohort, 3979 cases of baseline anemia were identified representing 5.5% of the WHI-OS study population (Table 1). Anemia was higher in women of advanced age (>63.5 years) and current smoking and inversely associated with education, income, and body weight (Table 1). Body mass index (BMI) showed a “u-shaped” association with anemia risk. Race/ethnicity also was associated with anemia with 16.3% of the 5413 African Americans having hemoglobin below 12.0 mg/dl and rates of 5.6%, 4.5%, and 4.2% in Hispanic/Latino, Non-Hispanic Whites, and Asians, respectively. Hemoglobin concentrations were normally distributed across the population.

Table 2 shows the average dietary intake at baseline among WHI-OS women categorized by anemia status. As shown, lower reported intake of most nutrients was demonstrated for women diagnosed with either incident or persistent anemia. Higher mean dietary intakes of energy, protein, folate, vitamin B12, iron and vitamin C as well as red meat were reported in women without anemia as compared to those with anemia. Supplemental iron and vitamin B12 were greater in women with anemia. Multivitamin use was not different by anemia group.

Given that poor diets may be associated with multiple nutrient deficiencies and that inadequacies in dietary intake may present differently across racial/ethnic groups, frequency of multiple deficiencies for iron, folate and vitamin B12 by race/ethnicity were also evaluated (Table 3). At baseline, the majority of WHI-OS women demonstrated adequate intake of the anemia-associated nutrients at levels meeting recommendations. However, Non-Hispanic whites were less likely to report multiple inadequacies compared to other racial/ethnic groups. Only 7.4% of non-Hispanic whites reported inadequacies in all three nutrients while 15.2% of Native Americans/Alaskans, 14.6 % Asian/Pacific Islanders, 15.3% of African Americans and 16.3% of Hispanic/Latinos reported all three nutrient inadequacies.

The odds ratios for incident and persistent anemia in relation to dietary inadequacies in intake of folate, iron, vitamin B12, vitamin C, and protein are presented in Table 4. Inadequate intakes of each of the evaluated nutrients at baseline were associated with increased risk of both incident and persistent anemia, with the associations tending to be stronger with persistent anemia than with incident anemia. An evaluation of anemia risk in relation to red meat (as an indicator of heme iron) versus fortified cold breakfast cereals (as a source of non-heme iron and/or folic acid) showed that greater red meat consumption was marginally associated with lower risk of persistent anemia (OR=0.89; 95% CI: 0.79–1.01) while fortified cold breakfast cereal was not (Table 5). These associations were attenuated when adjusted for covariates.

Modeling the association of the total number of inadequate nutrient intakes including iron, folate, and vitamin B12 at baseline with anemia suggests that a single nutrient inadequacy is associated with a significant increase in risk (Table 6). Tests for trends detected significantly increased risk of baseline, incident, and persistent anemia with additional baseline dietary deficiency. Specifically, any dietary deficiency at baseline was associated with an 18% greater risk of baseline anemia (Dietary deficiency OR=1.18; 95% CI: 1.08–1.29); a 10% (OR=1.10; 95% CI: 0.99–1.22) greater risk of incident anemia and a 21% (OR=1.21; 95% CI: 1.05–1.41) greater risk of persistent anemia, respectively. Total nutrient deficiencies were also associated with increased anemia risk, most substantially in relation to the demonstrated 34% increased risk for persistent anemia (OR=1.34; 95% CI: 1.14–1.56), a risk elevation that reached 56% when three deficiencies were present (OR=1.56; 95% CI: 1.25–1.95).

Discussion

This study of diet and anemia represents one of the largest assessments of anemia and dietary intake among postmenopausal women. The findings demonstrate that anemia associated nutrient inadequacies are not uncommon in postmenopausal women regardless of race/ethnicity and that select racial groups, such as African Americans, are at an elevated risk for diet-associated anemia as compared to non-Hispanic whites. These data also suggest that inadequacy in several nutrients, as compared to a single nutrient inadequacy, is associated with greater persistence of anemia in older women. Among eligible WHI-OS participants, 5.5% presented on enrollment with hemoglobin values < 12.0 mg/dL a

prevalence rate well below the 10.2% reported in a nationally representative sample of older women from NHANES III as well as rates reported in similar population samples including the Framingham cohort (4,27,28). One important difference across samples is that the WHI-OS sample included women as young as 50 years; however, when anemia prevalence was evaluated only for those women over 65 years of age, the estimate only increased to 5.64%. Nevertheless, WHI-OS is not a random sample of the aging female population in the U.S. The lower prevalence of anemia may partially reflect a nutritionally replete population of postmenopausal women. Baseline estimates of dietary intake in the WHI-OS sample averaged at the time of study enrollment 12 mg iron, 486 mcg folate (suggesting adequate intake even prior to mandate for folic acid fortification of grain products in 1998) and 5.6 mcg vitamin B12 daily; intakes above the Estimated Average Requirements of 6 mg, 320 mcg and 2.0 mcg daily, respectively for these anemia-associated nutrients. This suggests that fewer women demonstrated low hemoglobin concentrations than would have been expected based on cross-sectional diet reports at baseline.

An additional explanation for the lower rates of anemia in this sample may be that WHI-OS was a sample of free-living older women in contrast to other studies among institutionalized or home-bound elderly wherein anemia prevalence is reportedly as high as 39.6% (17,29,30). Women with renal disease were excluded from enrollment in WHI-OS and women with chronic diseases were excluded from the analytical cohort. There is the possibility that as the population aged, renal dysfunction may have contributed to new cases (31). Further, data regarding gastrointestinal diagnoses (e.g., irritable bowel syndrome) and/or medication use that might alter nutrient absorption (i.e., proton pump inhibitors, H2 blockers) are not available to evaluate as potential confounders. Consistent with data from other population samples, this analysis indicates that anemia prevalence is higher among women of lower income and smokers (32,33).

The association between body mass index and anemia is less frequently described in the literature, although a multiethnic cohort study among adults residing in New York City demonstrated an inverse association between body mass index (BMI) and circulating iron in Hispanic/Latino, but not non-Hispanic, women (34). These findings demonstrated a “u-shaped” association between BMI and low hemoglobin. This may suggest that nutrient inadequacies contribute to anemia rates among women with lower BMI (<25 kg/m²) while increased hepcidin concentrations that are more common in obese women, particularly in extreme obesity (11), could impair iron absorption resulting in anemia when BMI is > 30 kg/m². In support of this interpretation, a study of 50 obese women in Spain demonstrated a significant association between BMI and soluble transferrin receptor (35). Small sample size in these subgroups and lack of objective measures of iron regulatory proteins makes our interpretation of these findings somewhat speculative.

Few studies have explored racial/ethnic differences in anemia prevalence. This sample, which is enriched for minority post-menopausal women, found a higher prevalence of anemia in African Americans and, to a somewhat lesser extent, in Hispanic women. NHANES III reported a 28% prevalence of anemia among African American women, compared to only 8.7% in non-Hispanic whites (4). Additionally, anemia was found in 11.5% of older Hispanic women enrolled in the Massachusetts Hispanic Elders Study as compared to 7.3% in non-Hispanic white women (36). Our data also suggests that multiple anemia-associated nutrient inadequacies (3 nutrients or more) are more common in African American (15.3%) and Hispanic/Latinos (16.3%) than non-Hispanic White (7.4%) women suggesting that diet quality may be one factor contributing to risk. Earlier evidence suggests that lower dietary intake alone does not explain the totality of racial differences in hemoglobin levels (37). Genetic differences in sickle cell and thalassemia traits likely

account for some of the variance (38). This suggests that hemoglobin alone may have limited value in evaluating anemia in multi-racial/ethnic populations.

Inadequate intake of dietary iron, vitamin B12 and folate were each associated with approximately 10 to 20% elevated risk for incident anemia among WHI-OS study participants and the odds increased for persistent anemia to 21%. Persistent anemia was also associated with inadequate intake of vitamin C (OR 1.25, 95% CI 1.11–1.40) and protein (OR 1.22, 95% CI 1.08–1.36) suggesting that among women with persistent anemia a diet pattern of low overall nutrient density is likely. There was no indication that supplemental multivitamin or multivitamin-mineral use was protective against anemia. Anemia was diagnosed in 7.3% of women taking multi-nutrient supplements and 7.2% of those not taking these supplements. In addition, the levels of supplemental B12 or supplemental iron were higher in women demonstrating incident or persistent anemia as compared to women with no anemia. This finding suggests that nutritional supplementation for the treatment of anemia may have been a confounder in our analysis. As no specific data regarding clinical diagnosis and treatment of nutritional anemia is available for this cohort, it is only assumed that among women taking these supplements the demonstrated low hemoglobin concentrations reflect either a new diagnosis under treatment or a lack of response to supplementation. Also, women previously diagnosed with anemia may have been prescribed supplementation prior to enrolling in the WHI study. If supplementation corrected a previous inadequacy this would mask our ability to detect these associations using post-enrollment data. It is clear from this analysis (Table 6) that in WHI-OS the risk for persistent anemia is incrementally higher for people with an increased number of inadequate dietary intakes of the selected nutrients. These findings are important and suggest that individual nutrient inadequacies leading to nutritional anemia frequently do not occur in isolation. Therefore, reducing the likelihood and/or severity of diet-associated anemia should include evaluation of the overall diet. It is possible that among post-menopausal women presenting with hemoglobin concentrations below 12.0 mg/dL, enhanced access to nutrient-dense foods that target iron, B12, and/or folate intake may be required to correct nutritional anemia. This is supported by the findings that multivitamin-mineral use was not associated with lower rates of anemia.

This study does have some inherent limitations. First, anemia risk was evaluated in relation to self-reported dietary intake using a food frequency questionnaire. It is known that these instruments are associated with significant measurement error (39), although more so in relation to energy and macronutrient intake than the micronutrients evaluated here. The dietary data reflect a food supply prior to folic acid fortification and thus current anemia rates may be lower. Further, the definition of anemia was based on cut-off points for hemoglobin concentrations alone (3). While these values do correlate with other measures of anemia status, serum ferritin is the more generally accepted clinical measure of iron-deficiency anemia (40) and macrocytic anemias are better diagnosed using soluble or red blood cell folate and/or serum vitamin B12 levels. A more complete assessment of anemia also would include additional blood biomarkers, such as MCV, MCH, RBC, serum folate and serum vitamin B12 and ferritin as well as serum transferrin receptor and hepcidin. C-reactive protein and other inflammatory indices would also be included in order to rule out non-nutritional causes of anemia common in the elderly. These measures are currently not available for the entire WHI-OS study population. Further, current literature suggests that anemia cut-offs should be adjusted in older people and based on the association between blood biomarkers of anemia and co-morbidities such as falls, hospitalizations, etc (2,6). This study is also an observational study of a non-representative U.S. sample and as such has inherent limitations including lack of generalizability and the inability to determine whether improving the diet in relation to enhancement of intake of select micronutrients would result in correction of low hemoglobin concentrations.

There are several important strengths to this research. This is the first study of this size to investigate the associations between dietary intake and anemia in a prospective sample of aging, postmenopausal women. The study participants have provided extensive demographic and clinical data resulting in a robust characterization of enrollees and an opportunity to control for confounding factors. The multiethnic enrollment serves as a notable strength of this study in that few studies provide such diversity in describing anemia-diet associations. Repeated measures of hemoglobin allow for a prospective assessment of new on-set anemia and an opportunity to exclude acute anemia likely associated with non-dietary etiologies such as inflammation, new onset disease, extreme obesity or infection.

Conclusion

This study suggests that inadequate nutrient intakes are a significant risk factor for anemia in this population of older women and use of multivitamin-mineral supplements is not associated with lower rates of anemia. The study cannot ascertain whether nutrient repletion in the setting of non-nutritional causes of anemia (blood loss, inflammation, select medication use, etc) would alter risk given the lack of specific data in these areas. Yet, overall mortality is increased in relation to a diagnosis of anemia, (1,41) and anemia, particularly iron deficiency, has been associated with reduced capacity for physical work and physical inactivity (42–44), injury related to falls (45) and hospitalizations (2), making this an important health care concern in the aging. Efforts to identify anemia that may be responsive to modifiable factors such as diet to improve health outcomes are needed. Additional efforts to regularly evaluate postmenopausal women for anemia should be considered and should be accompanied by an assessment of dietary intake to determine adequacy of intake of anemia-associated nutrients including iron, vitamin B12 and folate. While anemia is often treated with supplemental nutrients, as designated by a more comprehensive biochemical assessment, nutritional therapy to improve overall nutrient-density and quality of the diet should also be a clinical focus.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Program Office: (National Heart, Lung, and Blood Institute, Bethesda, Maryland) Elizabeth Nabel, Jacques Rossouw, Shari Ludlam, Joan McGowan, Leslie Ford, and Nancy Geller.

Clinical Coordinating Center: (Fred Hutchinson Cancer Research Center, Seattle, WA)

Ross Prentice, Garnet Anderson, Andrea LaCroix, Charles L. Kooperberg, Ruth E. Patterson, Anne McTiernan; (Medical Research Labs, Highland Heights, KY) Evan Stein; (University of California at San Francisco, San Francisco, CA) Steven Cummings.

Clinical Centers: (Albert Einstein College of Medicine, Bronx, NY) Sylvia Wassertheil-Smolter; (Baylor College of Medicine, Houston, TX) Aleksandar Rajkovic; (Brigham and Women's Hospital, Harvard Medical School, Boston, MA) JoAnn E. Manson; (Brown University, Providence, RI) Charles B. Eaton; (Emory University, Atlanta, GA) Lawrence Phillips; (Fred Hutchinson Cancer Research Center, Seattle, WA) Shirley Beresford; (George Washington University Medical Center, Washington, DC) Lisa Martin; (Los Angeles Biomedical Research Institute at Harbor- UCLA Medical Center, Torrance, CA) Rowan Chlebowski; (Kaiser Permanente Center for Health Research, Portland, OR) Yvonne Michael; (Kaiser Permanente Division of Research, Oakland, CA) Bette Caan; (Medical College of Wisconsin, Milwaukee, WI) Jane Morley Kotchen; (MedStar Research Institute/Howard University, Washington, DC) Barbara V. Howard; (Northwestern University, Chicago/Evanston, IL) Linda Van Horn; (Rush Medical Center, Chicago, IL) Henry Black; (Stanford Prevention Research Center, Stanford, CA) Marcia L. Stefanick; (State University of New York at Stony Brook, Stony Brook, NY) Dorothy Lane; (The Ohio State University, Columbus, OH) Rebecca Jackson; (University of Alabama at Birmingham, Birmingham, AL) Cora E. Lewis; (University of Arizona, Tucson/Phoenix, AZ) Cynthia A Thomson; (University at Buffalo, Buffalo, NY) Jean Wactawski-Wende; (University of California at Davis, Sacramento, CA) John Robbins; (University of

California at Irvine, CA) F. Allan Hubbell; (University of California at Los Angeles, Los Angeles, CA) Lauren Nathan; (University of California at San Diego, LaJolla/Chula Vista, CA) Robert D. Langer; (University of Cincinnati, Cincinnati, OH) Margery Gass; (University of Florida, Gainesville/Jacksonville, FL) Marian Limacher; (University of Hawaii, Honolulu, HI) J. David Curb; (University of Iowa, Iowa City/Davenport, IA) Robert Wallace; (University of Massachusetts/Fallon Clinic, Worcester, MA) Judith Ockene; (University of Medicine and Dentistry of New Jersey, Newark, NJ) Norman Lasser; (University of Miami, Miami, FL) Mary Jo O'Sullivan; (University of Minnesota, Minneapolis, MN) Karen Margolis; (University of Nevada, Reno, NV) Robert Brunner; (University of North Carolina, Chapel Hill, NC) Gerardo Heiss; (University of Pittsburgh, Pittsburgh, PA) Lewis Kuller; (University of Tennessee Health Science Center, Memphis, TN) Karen C. Johnson; (University of Texas Health Science Center, San Antonio, TX) Robert Brzyski; (University of Wisconsin, Madison, WI) Gloria E. Sarto; (Wake Forest University School of Medicine, Winston-Salem, NC) Mara Vitolins; (Wayne State University School of Medicine/Hutzel Hospital, Detroit, MI) Michael Simon.

References

1. Semba RD, Ricks MO, Ferrucci L, Xue QL, Chaves P, Fried LP, Guralnik JM. Types of anemia and mortality among older disabled women living in the community: the Women's Health and Aging Study I. *Aging Clin Exp Res.* 2007; 19:259–264. [PubMed: 17726354]
2. Penninx BW, Pahor M, Woodman RC, Guralnik JM. Anemia in old age is associated with increased mortality and hospitalization. *J Gerontol A Biol Sci Med Sci.* 2006; 61:474–479. [PubMed: 16720744]
3. World Health Organization. Nutritional anemia: report of a World Health Organization Scientific Group. Geneva, Switzerland: World Health Organization; 1968.
4. Guralnik JM, Eisenstaedt RS, Ferrucci L, Klein HG, Woodman RC. Prevalence of anemia in persons 65 years and older in the United States: evidence for a high rate of unexplained anemia. *Blood.* 2004; 104:2263–2268. [PubMed: 15238427]
5. Culleton BF, Manns BJ, Zhang J, Tonelli M, Klarenbach S, Hemmelgarn BR. Impact of anemia on hospitalization and mortality in older adults. *Blood.* 2006; 107:3841–3846. [PubMed: 16403909]
6. Penninx BW, Pahor M, Cesari M, Corsi AM, Woodman RC, Bandinelli S, Guralnik JM, Ferrucci L. Anemia is associated with disability and decreased physical performance and muscle strength in the elderly. *J Am Geriatr Soc.* 2004; 52:719–724. [PubMed: 15086651]
7. Penninx BW, Pluijm SM, Lips P, Woodman R, Miedema K, Guralnik JM, Deeg DJ. Late-life anemia is associated with increased risk of recurrent falls. *J Am Geriatr Soc.* 2005; 53:2106–2111. [PubMed: 16398894]
8. Fishman SM, Christian P, West KP. The role of vitamins in the prevention and control of anaemia. *Public Health Nutr.* 2000; 3:125–150. [PubMed: 10948381]
9. Boccio JR, Iyengar V. Iron deficiency: causes, consequences, and strategies to overcome this nutritional problem. *Biol Trace Elem Res.* 2003; 94:1–32. [PubMed: 12907824]
10. Doyle W, Crawley H, Robert H, Bates CJ. Iron deficiency in older people: interactions between food and nutrient intakes with biochemical measures of iron; further analysis of the National Diet and Nutrition Survey of people aged 65 years and over. *Eur J Clin Nutr.* 1999; 53:552–559. [PubMed: 10452410]
11. McClung JP, Karl JP. Iron deficiency and obesity: the contribution of inflammation and diminished iron absorption. *Nutr Rev.* 2010; 67(2):100–104. [PubMed: 19178651]
12. Beghe C, Wilson A, Ershler WB. Prevalence and outcomes of anemia in geriatrics: a systematic review of the literature. *Am J Med.* 2004; 116 (Suppl 7A):3S–10S. [PubMed: 15050882]
13. Elwood PC. Evaluation of the clinical importance of anemia. *Am J Clin Nutr.* 1973; 26:958–964. [PubMed: 4580438]
14. Eisenstaedt R, Penninx BW, Woodman RC. Anemia in the elderly: current understanding and emerging concepts. *Blood Rev.* 2006; 20:213–226. [PubMed: 16472893]
15. Ferrucci L, Corsi A, Lauretani F, Bandinelli S, Bartali B, Taub DD, Guralnik JM, Longo DL. The origins of age-related proinflammatory state. *Blood.* 2005; 105:2294–2299. [PubMed: 15572589]
16. Shah M, Adams-Huet B, Kavanaugh A, Coyle Y, Lipsky P. Nutrient intake and diet quality in patients with systemic lupus erythematosus on a culturally sensitive cholesterol lowering dietary program. *J Rheumatol.* 2004; 31:71–75. [PubMed: 14705221]

17. Artz AS, Fergusson D, Drinka PJ, Gerald M, Bidenbender R, Lechich A, Silverstone F, McCamish MA, Dai J, Keller R, Ershler WB. Mechanisms of unexplained anemia in the nursing home. *J Am Geriatr Soc.* 2004; 52:423–427. [PubMed: 14962159]
18. Guralnik, JM.; Ershler, WB.; Schrier, SL.; Picozzi, VJ. Anemia in the elderly: a public health crisis in hematology. *Hematology: American Society of Hematology Special Symposium*; 2005. p. 528-532.
19. The Women's Health Initiative Study Group. Design of the Women's Health Initiative clinical trial and observational study. *Control Clin Trials.* 1998; 19:61–109. [PubMed: 9492970]
20. Knight EL, Rimm EB, Pai JK, Rexrode KM, Cannuscio CC, Manson JE, Stampfer MJ, Curhan GC. Kidney dysfunction, inflammation, and coronary events: a prospective study. *J Am Soc Nephrol.* 2004; 15:1897–1903. [PubMed: 15213279]
21. Mitchell JE. Emerging role of anemia in heart failure. *Am J Cardiol.* 2007; 99:15D–20D.
22. Wilson A, Yu HT, Goodnough LT, Nissenson AR. Prevalence and outcomes of anemia in rheumatoid arthritis: a systematic review of the literature. *Am J Med.* 2004; 116 (Suppl 7A):50S–57S. [PubMed: 15050886]
23. Wingard R. Increased risk of anemia in dialysis patients with comorbid diseases. *Nephrol Nurs J.* 2004; 31:211–214. [PubMed: 15114802]
24. Silverberg DS, Wexler D, Iaina A, Schwartz D. The interaction between heart failure and other heart diseases, renal failure, and anemia. *Semin Nephrol.* 2006; 26:296–306. [PubMed: 16949468]
25. Patterson RE, Kristal AR, Tinker LF, Carter RA, Bolton MP, Agurs-Collins T. Measurement characteristics of the Women's Health Initiative food frequency questionnaire. *Ann Epidemiol.* 1999; 9:178–187. [PubMed: 10192650]
26. Schakel SF, Sievert YA, Buzzard IM. Sources of data for developing and maintaining a nutrient database. *J Am Diet Assoc.* 1988; 88:1268–1271. [PubMed: 3171020]
27. Fleming DJ, Jacques PF, Tucker KL, Massaro JM, D'Agostino RB Sr, Wilson PW, Wood RJ. Iron status of the free-living, elderly Framingham Heart Study cohort: an iron-replete population with a high prevalence of elevated iron stores. *Am J Clin Nutr.* 2001; 73:638–646. [PubMed: 11237943]
28. Olivares M, Hertrampf E, Capurro MT, Wegner D. Prevalence of anemia in elderly subjects living at home: role of micronutrient deficiency and inflammation. *Eur J Clin Nutr.* 2000; 54:834–839. [PubMed: 11114677]
29. Argento V, Roylance J, Skudlarska B, Dainiak N, Amoateng-Adjepong Y. Anemia prevalence in a home visit geriatric population. *J Am Med Dir Assoc.* 2008; 9:422–426. [PubMed: 18585644]
30. Inelmen EM, D'Alessio M, Gatto MR, Baggio MB, Jimenez G, Bizzotto MG, Enzi G. Descriptive analysis of the prevalence of anemia in a randomly selected sample of elderly people living at home: some results of an Italian multicentric study. *Aging (Milan, Italy).* 1994; 6:81–89.
31. Lippi G, Franchini M, Salvagno GL, Montagnana M, Targher G, Guidi GC. Determinants of anaemia in the very elderly: a major contribution from impaired renal function. *Blood Transfus.* 2010; 8:44–48. [PubMed: 20104278]
32. Klesges LM, Pahor M, Shorr RI, Wan JY, Williamson JD, Guralnik JM. Financial difficulty in acquiring food among elderly disabled women: results from the Women's Health and Aging Study. *Am J Public Health.* 2001; 91:68–75. [PubMed: 11189828]
33. Pilch, S.; Senti, F. Assessment of iron nutritional status of the U.S. population based on data collected in the Second National Health and Nutrition Examination Survey, 1976–1980. Bethesda (MD): Federation of American Societies for Experimental Biology; 1984.
34. Chambers EC, Heshka S, Gallagher D, Wang J, Pi-Sunyer FX, Pierson RN Jr. Serum iron and body fat distribution in a multiethnic cohort of adults living in New York City. *J Am Diet Assoc.* 2006; 106:680–684. [PubMed: 16647325]
35. Lecube A, Carrera A, Losada E, Hernandez C, Simo R, Mesa J. Iron deficiency in obese postmenopausal women. *Obesity.* 2006; 14:1724–1730. [PubMed: 17062801]
36. Seaverson EL, Buell JS, Fleming DJ, Bermudez OI, Potischman N, Wood RJ, Chasan-Taber L, Tucker KL. Poor iron status is more prevalent in Hispanic than in non-Hispanic white older adults in Massachusetts. *J Nutr.* 2007; 137:414–420. [PubMed: 17237320]
37. Perry GS, Byers T, Yip R, Margen S. Iron nutrition does not account for the hemoglobin differences between blacks and whites. *J Nutr.* 1992; 122:1417–1424. [PubMed: 1619469]

38. Beutler E, West C. Hematological differences between African-American and whites: the roles of iron deficiency and α -thalassemia on hemoglobin levels and mean corpuscular volume. *Blood*. 2005; 106:740–745. [PubMed: 15790781]
39. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol*. 1985; 122:51–65. [PubMed: 4014201]
40. Guyatt GH, Oxman AD, Ali M, Willan A, McIlroy W, Patterson C. Laboratory diagnosis of iron-deficiency anemia: an overview. *J Gen Intern Med*. 1992; 7:145–153. [PubMed: 1487761]
41. Elwood PC, Waters WE, Benjamin IT, Sweetnam PM. Mortality and anaemia in women. *Lancet*. 1974; 1:891–894. [PubMed: 4133418]
42. Edgerton VR, Gardner GW, Ohira Y, Gunawardena KA, Senewiratne B. Iron-deficiency anaemia and its effect on worker productivity and activity patterns. *Br Med J*. 1979; 2:1546–1549. [PubMed: 534861]
43. Gardner GW, Edgerton VR, Senewiratne B, Barnard RJ, Ohira Y. Physical work capacity and metabolic stress in subjects with iron deficiency anemia. *Am J Clin Nutr*. 1977; 30:910–917. [PubMed: 868783]
44. Davies CT, Chukweumeka AC, Van Haaren JP. Iron-deficiency anaemia: its effect on maximum aerobic power and responses to exercise in African males aged 17–40 years. *Clin Sci*. 1973; 44:555–562. [PubMed: 4713663]
45. Duh MS, Mody SH, Lefebvre P, Woodman RC, Buteau S, Piech CT. Anaemia and the risk of injurious falls in a community-dwelling elderly population. *Drugs Aging*. 2008; 25:325–334. [PubMed: 18361542]

Table 1

Demographic characteristics of Women's Health Initiative (WHI) Observational Study women by anemia status (hemoglobin <12.0 mg/dL)

Variable	No Anemia		Anemia ^a	
	Number	%	Number	%
Total	68854	(94.5%)	3979	(5.5%)
Age at screening ^b	63.2	(7.3)	63.5	(7.6)
Education ^c				
Some high school or less (≤ 11 years)	3098	(4.5%)	265	(6.7%)
High school diploma or GED	11013	(16.1%)	672	(17.0%)
College or vocational school	54220	(79.3%)	3016	(76.3%)
Family income ^b				
Less than \$10,000	2336	(3.5%)	236	(6.2%)
\$10,000 to \$34,999	21447	(32.6%)	1296	(34.3%)
\$35,000 to \$99,999	32596	(49.5%)	1745	(46.1%)
More than \$100,000	7451	(11.3%)	369	(9.8%)
Don't know	2008	(3.0%)	137	(3.6%)
Ethnicity ^c				
American Indian or Alaskan native	270	(0.4%)	20	(0.5%)
Asian or pacific islander	2124	(3.1%)	94	(2.4%)
Black or African American	4533	(6.6%)	880	(22.2%)
Hispanic/Latino	2638	(3.8%)	156	(3.9%)
White (not of Hispanic origin)	58342	(85.0%)	2759	(69.6%)
Other	760	(1.1%)	56	(1.4%)
Body Mass Index (BMI)	27.1	(5.7)	27.0	(6.2)
BMI category ^c				
Underweight (< 18.5)	794	(1.2%)	54	(1.4%)
Normal (18.5 – 24.9)	27622	(40.6%)	1725	(43.8%)
Overweight (25.0 – 29.9)	23393	(34.4%)	1229	(31.2%)
Obesity I (30.0 – 34.9)	10398	(15.3%)	537	(13.6%)
Obesity II (35.0 – 39.9)	3687	(5.4%)	215	(5.5%)
Extreme obesity III (≥ 40)	2164	(3.2%)	178	(4.5%)

Variable	No Anemia		Anemia ^a	
	Number	%	Number	%
Height ^c	161.8	(6.7)	161.0	(6.8)
Weight ^d	71.3	(16.4)	70.4	(17.2)
Smoking status ^c				
Never smoked	34915	(51.4%)	2115	(54.1%)
Past smoker	28781	(42.4%)	1649	(42.1%)
Current smoker	4217	(6.2%)	149	(3.8%)
Hemoglobin (mg/dL) ^c mean (SD)	13.6	(0.9)	11.3	(1.0)
Hematocrit (%) ^c mean (SD)	40.3	(3.0)	34.5	(3.0)
White blood cell (kcell/ml) mean (SD)	6.2	(12.3)	5.9	(15.4)

^a Anemia as defined by World Health Organization (WHO) definition of hemoglobin < 12.0 mg/dL.

^b $p < 0.05$.

^c $p < 0.001$.

^d $p < 0.01$.

Table 2

Reported baseline nutrient intakes using Food Frequency Questionnaire (FFQ) data in WHI OS women according to anemia^a status; adjusted for BMI, age, ethnicity, smoking status and income

Dietary Variable	No Anemia		Incident Anemia ^b		Persistent Anemia ^c	
	Mean	(SD)	Mean	(SD)	Mean	(SD)
Dietary energy (kcal) ^d	1546.7	(580.4)	1517.0	(581.4)	1473.3	(584.8)
Dietary total carbohydrate (g) ^d	201.3	(77.0)	197.1	(77.2)	192.3	(77.6)
Dietary protein (g) ^d	65.3	(26.6)	63.7	(26.6)	61.2	(26.8)
Ratio of animal to vegetable protein ^e	2.4	(1.2)	2.4	(1.2)	2.4	(1.2)
Dietary folate equivalents (mcg) ^d	487.9	(206.2)	478.6	(206.6)	470.1	(207.8)
Dietary iron (mg) ^d	12.4	(5.3)	12.1	(5.3)	11.8	(5.4)
Dietary vitamin B12 (mcg) ^d	5.7	(3.3)	5.5	(3.3)	5.4	(3.3)
Dietary vitamin C (mg) ^d	104.6	(56.0)	101.8	(56.1)	98.3	(56.4)
Supplemental folic acid (mcg)	207.8	(259.2)	206.8	(259.7)	205.4	(261.2)
Supplemental iron (mg) ^d	8.6	(21.2)	11.4	(21.3)	11.9	(21.4)
Supplemental vitamin B12 (mcg) ^f	16.9	(75.5)	19.1	(75.6)	22.4	(76.1)
Supplemental vitamin C (mg)	330.8	(658.7)	310.6	(659.9)	324.4	(663.7)
Total dietary folate equivalents ^e	602.2	(252.4)	592.3	(252.9)	583.1	(254.3)
Total iron at baseline ^d	21.0	(21.9)	23.5	(21.9)	23.7	(22.0)
Total vitamin B12 at baseline ^e	22.5	(75.5)	24.6	(75.7)	27.7	(76.1)
Total vitamin C	435.4	(663.4)	412.4	(664.6)	422.6	(668.4)
Multivitamin without minerals						
Yes	2244	(4.2%)	116	(4.3%)	60	(4.2%)
No	51419	(95.8%)	2605	(95.7%)	1374	(95.8%)
Multivitamin with minerals						
Yes	20835	(38.8%)	1059	(38.9%)	522	(36.4%)
No	32828	(61.2%)	1662	(61.1%)	912	(63.6%)
Cold Cereal -- Fortified						
Yes	7022	(15.1%)	400	(14.7%)	193	(15.9%)

Dietary Variable	No Anemia		Incident Anemia ^b		Persistent Anemia ^c	
	Mean	(SD)	Mean	(SD)	Mean	(SD)
No	39516	(84.9%)	2321	(85.3%)	1016	(84.1%)
Cold Cereal -- medium servings/day	0.4	(0.4)	0.4	(0.4)	0.4	(0.4)
Red Meat -- medium servings/day ^d	0.6	(0.5)	0.6	(0.5)	0.6	(0.5)

^a Anemia as defined by WHO definition of hemoglobin < 12.0 mg/dL.

^b Incident anemia (newly diagnosed) is defined by a reduction in hemoglobin from values greater than or equal to 12.0 mg/dL at baseline to values less than 12.0 mg/dL at year three.

^c Persistent anemia is defined as hemoglobin less than 12.0 mg/dL at baseline and year 3.

^d $p<0.001$,

^e $p<0.05$,

^f $p<0.01$.

Table 3

Racial and ethnic differences in the number and percent of sample population reporting dietary nutrient inadequacies of folate, iron, and vitamin B12 in Women's Health Initiative Observational Study (n=72,632)

Number of nutrient inadequacies	American Indian or Native Alaska	Asian or Pacific Islander	Black or African American	Hispanic/Latino	White (Not Hispanic)	Other	Total
0	144 (49.79%)	1,169 (52.7%)	2,338 (43.2%)	1,324 (47.4%)	37653 (53.1%)	433 (53.1%)	43,061 (59.3%)
1	54 (18.6%)	345 (15.6%)	1,128 (20.8%)	506 (18.1%)	10701 (17.5%)	152 (18.6%)	12,886 (17.7%)
2	48 (16.6%)	380 (17.1%)	1,120 (20.7%)	508 (18.2%)	8,250 (13.5%)	124 (15.2%)	10,430 (14.4%)
3	44 (15.2%)	324 (14.6%)	827 (15.3%)	456 (16.3%)	4,497 (7.4%)	107 (13.1%)	6,255 (8.6%)
Total	290	2,218	5,413	2,784	61,101	816	72,632

Odds Ratios (OR) and 95% confidence interval (CI) for incident and persistent anemia, relative to no anemia^a (n=53,663), associated with inadequate of dietary intake of select nutrients as reported by food frequency questionnaire at baseline (Women's Health Initiative Observational Study (WHI OS))

Table 4

	Incident Anemia ^b (n= 2,721)			Persistent Anemia ^c (n= 1,434)		
	Crude	Adjusted ^d		Crude	Adjusted ^d	
	OR	95% CI	OR	95% CI	OR	95% CI
Dietary Folate Equivalents <400 mcg	1.20	(1.11, 1.30)	1.12	(1.03, 1.21)	1.46	(1.32, 1.63)
Dietary Iron <8 mg	1.23	(1.13, 1.35)	1.13	(1.02, 1.24)	1.56	(1.38, 1.75)
Dietary vitamin B12 <2.4 mcg	1.25	(1.12, 1.40)	1.16	(1.03, 1.30)	1.56	(1.35, 1.80)
Dietary Vitamin C <75 mg	1.08	(0.99, 1.17)	1.09	(1.00, 1.19)	1.24	(1.12, 1.39)
Dietary Protein <0.8 g/kg	1.18	(1.10, 1.28)	1.07	(0.98, 1.16)	1.43	(1.29, 1.59)
Dietary Protein <0.8 g/kg (energy adjusted)	1.12	(1.02, 1.24)	0.99	(0.89, 1.11)	1.21	(1.06, 1.39)

^a no anemia as defined by WHO definition of hemoglobin $\geq$ 12.0 mg/dL.

^b Incident anemia (newly diagnosed) is defined by a reduction in hemoglobin from values greater than or equal to 12.0 mg/dL at baseline to values less than 12.0 mg/dL at year three.

^c Persistent anemia is defined as hemoglobin less than 12.0 mg/dL at baseline and year 3.

^d Adjusted for BMI, age range, race/ethnicity, smoking status, and income.

Odds Ratios (OR) and 95% confidence interval (CI) for incident and persistent anemia, relative to no anemia^a (n=53,663), associated with each additional serving of selected food groups

Table 5

	Incident Anemia ^b (n= 2,721)			Persistent Anemia ^c (n= 1,434)		
	Crude		Adjusted ^d	Crude		Adjusted ^d
	OR	95% CI	OR	95% CI	OR	95% CI
Red Meat	0.95	(0.88, 1.03)	0.98	(0.90, 1.06)	0.82	(0.73, 0.92)
Fortified Cold Cereal	0.95	(0.85, 1.07)	0.97	(0.86, 1.09)	1.04	(0.89, 1.21)

^aNo anemia as defined by WHO definition of hemoglobin ≥ 12.0 mg/dL.

^bIncident anemia (newly diagnosed) is defined by a reduction in hemoglobin from values greater than or equal to 12.0 mg/dL at baseline to values less than 12.0 mg/dL at year three.

^cPersistent anemia is defined as hemoglobin less than 12.0 mg/dL at baseline and year 3.

^dAdjusted for BMI, age range, race/ethnicity, smoking status, and income.

Odds Ratios (OR)^a and 95% confidence interval (CI) for incident^b and persistent anemia^c, relative to no anemia^d, associated with the number of anemia-associated dietary micronutrient inadequacies at baseline (folate, vitamin B12, vitamin C, iron and protein) among women in the Women's Health Initiative Observational Study (WHI OS)

Table 6

		Baseline Anemia ^d		Incident Anemia ^b		Persistent Anemia ^c	
		OR	95% CI	OR	95% CI	OR	95% CI
Number of deficiencies of dietary intake	1	1.18	(1.08, 1.29)	1.10	(0.99, 1.22)	1.21	(1.05, 1.41)
	2	1.19	(1.09, 1.31)	1.09	(0.96, 1.22)	1.32	(1.13, 1.53)
	3	1.27	(1.13, 1.42)	1.24	(1.08, 1.43)	1.44	(1.20, 1.73)
Number of deficiencies of total intake	1	1.23	(1.12, 1.35)	1.09	(0.97, 1.23)	1.34	(1.14, 1.56)
	2	1.17	(1.04, 1.31)	1.09	(0.94, 1.25)	1.39	(1.16, 1.66)
	3	1.28	(1.11, 1.48)	1.14	(0.94, 1.37)	1.56	(1.25, 1.95)

^a Adjusted for BMI, age, income, smoking status, and race/ethnicity.

^b Incident anemia defined as hemoglobin less than 12 g/dL at baseline assessment;

^c Persistent anemia defined as hemoglobin less than 12.0 g/dL at baseline and year 3 assessments.

^d No anemia as defined by WHO definition of hemoglobin $\geq$ 12.0 mg/dL; anemia defined as hemoglobin $<$ 12.0 mg/dL.