ERRATUM

Mutually Exclusive Genotypes for Pyrazinamide and 5-Chloropyrazinamide Resistance Reveal a Potential Resistance-Proofing Strategy

Anthony D. Baughn, Jiaoyu Deng, Catherine Vilchèze, Angelica Riestra, John T. Welch, William R. Jacobs, Jr., and Oren Zimhony

Howard Hughes Medical Institute, Department of Microbiology and Immunology, Albert Einstein College of Medicine, Bronx, New York; Department of Chemistry, State University of New York at Albany, Albany, New York; and Infectious Diseases Division, Kaplan Medical Center, School of Medicine, Hebrew University and Hadassah Jerusalem, Rehovot 76100, Israel

Volume 54, no. 12, p. 5323–5328, 2010. Page 5324, Table 2: The table headings “pH 6” and “pH 6.8” were inadvertently switched twice during typesetting. Table 2 should appear as shown below.

TABLE 2. Effect of PncA and PzaA expression on PZA and 5-Cl PZA turnover and MIC in tuberculous bacilli

Strain	Characteristic ^a	5-Cl PZA			PZA			Reference
		Turnover (nmol/min/ml of cells [OD ₆₀₀])	MIC µg/ml		Turnover (nmol/min/ml of cells [OD ₆₀₀])	MIC µg/ml		
			pH 6.8	pH 6		pH 6.8	pH 6	
<i>M. tuberculosis</i> H37Ra	Attenuated mutant of H37Rv	0.15 ± 0.01	25	25	0.23 ± 0.01	>1,000	50	
mc ² 7092	H37Ra <i>attB</i> _{L5} ::P _{Tc} :: <i>pzaA</i> _{Msmeg}	27 ± 1	200	100	20 ± 1	62.5	25	This work
mc ² 7093	H37Ra <i>attB</i> _{L5} ::P _{Tc} :: <i>pncA</i> _{Mtb}	0.096 ± 0.002	25	25	2.9 ± 1	62.5	50	This work
<i>M. bovis</i> BCG-Pasteur	Attenuated mutant of <i>M. bovis</i>	0.11 ± 0.01	12.5	12.5	0.015 ± 0.001	>1,000	>1,000	
mc ² 7091	BCG-Pasteur <i>attB</i> _{L5} ::P _{Tc} :: <i>pzaA</i> _{Msmeg}	24 ± 1	200	50	16 ± 1	62.5	12.5	This work
mc ² 7099	BCG-Pasteur <i>attB</i> _{L5} ::P _{Tc} :: <i>pncA</i> _{Mtb}	0.080 ± 0.01	12.5	12.5	3.5 ± 0.1	62.5	12.5	This work

^a Msmeg, *M. smegmatis*; Mtb, *M. tuberculosis*.