Draft Genome Sequence of *Turicibacter sanguinis* PC909, Isolated from Human Feces[▽]

Páraic Ó Cuív,¹ Eline S. Klaassens,¹ A. Scott Durkin,² Derek M. Harkins,² Les Foster,²
Jamison McCarrison,² Manolito Torralba,² Karen E. Nelson,² and Mark Morrison^{1,3*}

CSIRO Preventative Health Flagship Research Program and Division of Livestock Industries, Queensland Biosciences Precinct,
306 Carmody Road, Queensland 4067, Australia¹; J. Craig Venter Institute, Rockville, Maryland 20850²; and
The Ohio State University, Columbus, Ohio³

Received 2 November 2010/Accepted 13 December 2010

While the microbiota resident in the human gut is now known to provide a range of functions relevant to host health, many of the microbial members of the community have not yet been cultured or are represented by a limited number of isolates. We describe here the draft genome sequence of *Turicibacter sanguinis* PC909, isolated from a pooled healthy human fecal sample as part of the Australian Human Gut Microbiome Project.

Turicibacter spp. belong to a deeply branching, low-DNA G+C class of the *Firmicutes* termed the *Erysipelotrichia* that is comprised of the single order *Erysipelotrichiales* and the family *Erysipelotrichaceae* (3). *Turicibacter* spp. have been detected in the gastrointestinal tracts of several animals, including humans (e.g., references 4 and 7), suggesting that they may be important members of the gut microbiota. Consistent with this, *Turicibacter* spp. have been found to constitute part of the core measurable microbiota in mice, where they have also been shown to vary quantitatively in association with the host genotype (1). To date, only one axenic isolate has been described, *Turicibacter sanguinis* MOL361, a strain that was recovered from the bloodstream of a patient with acute appendicitis (2). However, it has been suggested that an uncultured murine phylotype, *Turicibacter* 458, may possess putative immunomodulatory and invasive properties (9) and that *Turicibacter* spp. may cause subclinical infections in piglets (10).

The *rrs* gene of *T. sanguinis* PC909 displays 100% sequence identity to the corresponding region of *T. sanguinis* MOL361 (NR_028816) and that of *Turicibacter* 458 (EU375462). In order to better understand the interaction of *T. sanguinis* with the host, we performed whole-genome sequencing using a 454 Life Sciences GS FLX system at the J. Craig Venter Institute (JCVI). The sequence data consists of 2,953,411 bp of DNA sequence at 70.6× coverage. Contig assembly was performed using the Newbler Assembler v. 2.3. The sequences assembled into 125 individual contigs, with a contig N50 of approximately 40.7 kb and with the largest contig approximately 146.6 kb. The DNA sequences were annotated using JCVI's prokaryotic annotation pipeline.

The draft genome has a G+C content of 34% and contains 2,851 genes with 2,781 protein coding genes and 70 structural RNAs. The annotated genome sequence revealed the presence of three internalin related proteins, including a protein with

sequence similarity to internalin A from *Listeria monocytogenes*, which has been shown to be necessary for gastrointestinal (6) and intracellular invasion (5). Comparison with the Virulence Factor Database (11) revealed the presence of proteins with suspected roles in binding to host structural factors (e.g., fibronectin, laminin, HSP60), the production of a collagen binding pilus, and capsule biosynthesis. Interestingly, genes related to sporulation and tolerance to oxygen species were also identified in the *T. sanguinis* PC909 genome sequence, which contrasts with the original phenotypic description for the genus and species (2).

Two other *Turicibacter* sp. genomes are currently being sequenced, *T. sanguinis* MOL361 and *Turicibacter* sp. HGF1, and in addition to that of *T. sanguinis* PC909, these genome sequences will provide a greater insight into the functional repertoire and potential pangenome of *Turicibacter* spp.

Nucleotide sequence accession numbers. The Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number ADMN000000000. The version described here is the first version, accession number ADMN01000000. The genome project data are also available at GenBank under the genome project ID 42765. The *rrs* sequence of *T. sanguinis* PC909 has been deposited in GenBank under accession number HQ428099.

This research was performed as part of CSIRO's contribution to the International Human Microbiome Consortium. The research was funded by CSIRO's Transformational Biology Capability Platform, in support of the Preventative Health Flagship Research Program. We also gratefully acknowledge the support provided by CSIRO's OCE Science Leader scheme (to M.M.) and the OCE Postdoctoral Fellow program (to E.S.K.).

We thank Stuart E. Denman and Carly P. Rosewarne for providing critical reading and suggestions to improve the manuscript.

REFERENCES

1. Benson, A. K., et al. 2010. Individuality in gut microbiota composition is a complex polygenic trait shaped by multiple environmental and host genetic factors. *Proc. Natl. Acad. Sci. U. S. A.* **107**:18933–18938.
2. Bosshard, P. P., R. Zbinden, and M. Altwegg. 2002. *Turicibacter sanguinis* gen. nov., sp. nov., a novel anaerobic, Gram-positive bacterium. *Int. J. Syst. Evol. Microbiol.* **52**:1263–1266.
3. Bosshard, P. P. 2009. Genus VIII. *Turicibacter* Bosshard, Zbinden and Altwegg 2002, 1266^{VP}, p. 1314–1315. In P. de Vos et al. (ed.), *Bergey's*

* Corresponding author. Mailing address: CSIRO Livestock Industries, Queensland Biosciences Precinct, 306 Carmody Road, Queensland 4067, Australia. Phone: 61 (0)7 3214 2216. Fax: 61 (0)7 3214 2900. E-mail: mark.morrison@CSIRO.au.

[▽] Published ahead of print on 23 December 2010.

- manual of systematic bacteriology, 2nd ed., vol. 3. The Firmicutes. Springer-Verlag, New York, NY.
4. **Falk, A., et al.** 2007. Ileal pelvic pouch microbiota from two former ulcerative colitis patients, analysed by DNA-based methods, were unstable over time and showed the presence of *Clostridium perfringens*. *Scand. J. Gastroenterol.* **42**:973–985.
 5. **Gaillard, J. L., P. Berche, C. Frehel, E. Gouln, and P. Cossart.** 1991. Entry of *L. monocytogenes* into cells is mediated by internalin, a repeat protein reminiscent of surface antigens from gram-positive cocci. *Cell* **65**:1127.
 6. **Lecuit, M., et al.** 2001. A transgenic model for listeriosis: role of internalin in crossing the intestinal barrier. *Science* **292**:1722–1725.
 7. **Licht, T. R., B. Madsen, and A. Wilks.** 2007. Selection of bacteria originating from a human intestinal microbiota in the gut of previously germ-free rats. *FEMS Microbiol. Lett.* **277**:205–209.
 8. Reference deleted.
 9. **Presley, L. L., B. Wei, J. Braun, and J. Borneman.** 2010. Bacteria associated with immunoregulatory cells in mice. *Appl. Environ. Microbiol.* **76**:936–941.
 10. **Rettedal, E., et al.** 2009. Alteration of the ileal microbiota of weanling piglets by the growth-promoting antibiotic chlortetracycline. *Appl. Environ. Microbiol.* **75**:5489–5495.
 11. **Yang, J., L. Chen, L. Sun, J. Yu, and Q. Jin.** 2008. VFDB 2008 release: an enhanced web-based resource for comparative pathogenomics. *Nucleic Acids Res.* **36**:D539–D542.