CORRESPONDENCE

Eight Years of Colonoscopic Bowel Cancer Screening in Germany: Initial Findings and Projections

by Prof. Dr. med. Hermann Brenner, Dr. rer. soc. Lutz Altenhofen,
Dr. sc. hum. Michael Hoffmeister in volume 43/2010

„Number Needed to Screen“

The initial findings of the colonoscopic bowel cancer screening initiative in Germany are important and could give direction to the use of screening programs.

However, it seems sensible to present the results and validity of this screening program by using the terms usually applied in prevention—positive predictive value and numbers needed to screen.

In 2.82 million performed (or rather, projected) screening colonoscopies in members of statutory health insurance schemes aged 55–84, 98 734 precursor lesions and 47 168 preclinical (asymptomatic) cases of bowel cancer were treated successfully (projected).

The positive predictive value is the measure for a true positive test result (true positives/negatives). If colonoscopy screening has 90% sensitivity and specificity, the positive predictive value is 3%.

The number needed to screen is the number of screening rounds needed to prevent one death or adverse event. On the basis of the available data, the number needed to screen is 19.

If the results are even only approximately correctly interpreted and if the number needed to screen is compared with mammography screening, for example, the authors' demand for improved uptake of the screening program should be urgently supported.

DOI: 10.3238/arztebl.2011.0170a

REFERENCES

1. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Dr. med. Michael Rothe
Lehrbereich Allgemeinmedizin
Universität Bonn
Sigmund-Freud-Str. 25, Haus 33
53127 Bonn, Germany
Michael.Rothe@ukb.uni-bonn.de

Picture Is Distorted

The article rightly mentions the low uptake rate among persons entitled to have colonoscopy screening. However, since this is more or less the same group of people who had a once-only screening examination in 2003–2010, the total proportion of participants among those entitled to be screened is rather higher. The exception is the group of those who, having reached the age of 55, are newly added to the pool of those entitled, minus those who have died.

I am no mathematician, but I would calculate a value of some 18%, rather than 2.6%, as reported. Reporting only the annual participation rate distorts the picture. Obviously, an even higher rate would be extremely desirable.

DOI: 10.3238/arztebl.2011.0170b

REFERENCES

1. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Norbert Schuster

Krankenhaus Bethel Berlin gGmbH
Promenadenstraße 3–5, 12207 Berlin
KHBE@BethelNet.de

Only Screening? Really?

The colonoscopy uptake rate of just 2.6% compared with those entitled to be screened seems too small to be a reliable basis for projections.

In this context, it would be interesting to know how many of these 2.6% were actual screening colonoscopies and how many indicated colonoscopies are included in that rate. The effect of the measure may be substantially overestimated as a result.

In all experience, patients are far more easily persuaded to undergo colonoscopy screening if they are given indications of increased risk (irregular bowel movements, family history, or a positive test for fecal occult blood in the stools). Might reimbursement considerations be a contributing factor too? Early detection colonoscopy is reimbursed outside the regular budget.

Colonoscopy screening may therefore make an only seemingly large contribution to the prevention and early detection of colorectal cancers in Germany, because of—rather than in spite of—low uptake rates.

DOI: 10.3238/arztebl.2011.0170c

REFERENCES

1. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Dr. med. Markus Hedemann

Am Posthof 1, 31785 Hameln
praxis-posthof@t-online.de

Would Digital Rectal Examination in Over-45s Make Sense?

In the context of gynecological cancer screening, digital examination of the rectum is a standard procedure in the over=45s. This has been the law since 1977. Women have a far higher awareness of screening than men. 34% of annual participants since 1990 are women; men only account for 14% of participants. Does this result in a sex-specific difference in the prognosis and survival after treatment for rectal cancer owing to early detection? Rectal cancers account for 20–25% of colorectal cancers, which can be diagnosed by means of preventive colonoscopy—and this includes precursor lesions.

This would have to translate into a benefit in women, whose uptake of statutory cancer screening is double than it men.

However, the early findings of colonoscopy screening for bowel cancer do not support this assumption. As many cancers were detected in women aged 55–59 as in men (13% vs 12.2%). The same is true for adenomas (21% vs 19.3%). The age group of 55–59 year olds was chosen because these women are most likely to attend gynecological cancer screening.

Colonoscopy screenings after the 55th year of life (90% of incident cases are diagnosed after this age) are attended by only 2.6% of persons (aged 55–84) who are legally entitled to participate.

The estimated number of cancers for this cohort, for early-stage cancers and advanced adenomas, was slightly higher in men than in women. Half of these findings apply to persons aged 60–70 years. Very few women attend gynecological screenings in this decade of life.

This raises the question of whether rectal-digital examinations in the context of gynecological cancer screening should be re-evaluated 33 years after introduction.

By contrast, it is certainly of note that preventive colonoscopy is being evaluated as early as 8 years after its introduction.

DOI: 10.3238/arztebl.2011.0171a

REFERENCES

1. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Prof. Dr. med. J. Matthias Wenderlein
Universität Ulm, Eyth-Str. 14
89075 Ulm, Germany
wenderlein@gmx.de

Complete and Comprehensible Data Are Required

In order to evaluate the data regarding projections in the context of colonoscopy screening, critical appraisal and comparison with other preventive measures is needed. Such data can be calculated from the authors’

models with very little extra effort, as a similar publication by the same authors shows (1). It would therefore be helpful if the authors could provide the following information:

- In addition to the number of estimated prevented cancers, or those detected early, what is needed as a reference value is the total number of cases of bowel cancer for the individual age groups and time periods.
- Mention should be made of the estimated data that would apply to a situation without colonoscopy screening in Germany.
- The data should be presented with constant reference values and time periods. Thus, it should be reported how many persons the prevented cancer cases correspond to for each time period.
- The data should be presented with comprehensible reference values. For example: “As a result of colonoscopy screening, every year (over a period of 10 years) XX persons per 1000 population who are entitled to be screened receive a diagnosis of bowel cancer and XX die from their cancer. Without colonoscopy screening, XX per 1000 population would die each year (every 10 years).”
- In order to enable easier classification, the total number of cancer cases in each category should be listed.
- It should be discussed why no projections can be made regarding possible complications.

In general I would suggest that *Deutsches Ärzteblatt* makes it a rule to present study results within a context that rules out misunderstandings and misinterpretations and that enables readers to appraise data easily. Standards are available for communicating the results of cancer screening programs, and these can be adapted for scientific studies (2, 3).

DOI: 10.3238/arztebl.2011.0171b

REFERENCES

1. Brenner H, Hoffmeister M, Brenner G, Altenhofen L, Haug U: Expected reduction of colorectal cancer incidence within 8 years after introduction of the German screening colonoscopy programme: Estimates based on 1 875 708 screening colonoscopies. *Eur J Cancer* 2009; 45: 2027–33.
2. Koch K, Mühlhauser I, für den Fachbereich Patienteninformation des Deutschen Netzwerks für Evidenzbasierte Medizin: Kriterien zur Erstellung von Patienteninformationen zu Krebsfrüherkennungsuntersuchungen, 2008. www.ebm-netzwerk.de.
3. Gigerenzer G, Gaissmaier W, Kurz-Milcke EK, Schwartz LM, Woloshin S: Helping doctors and patients make sense of health statistics. *Psychol Sci Public Interest* 2007; 8(2): 53–96.
4. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Prof. Dr. med. Ingrid Mühlhauser
Universität Hamburg, MIN Fakultät
Gesundheitswissenschaften
Martin-Luther-King Platz 6
20146 Hamburg, Germany
Ingrid_Muehlhauser@uni-hamburg.de

In Reply:

We thank our correspondents for their interest in our work and for their comments.

We agree with Rohde that the positive predictive value and the numbers needed to screen can be additional useful measures and that they underline the importance of screening colonoscopy we described in our article.

We correctly presented uptake rates as annual participation rates. On the basis of these it is obviously possible to calculate approximate projections for longer time periods. However, these are subject to uncertainties since, epidemiologically speaking, what's under discussion is not a closed cohort. Whether the projection for 8 years as suggested by Schuster is the most sensible variant requires further discussion. In view of the 10-year screening intervals that are foreseen for screening colonoscopy, a 10-year projection would better reflect total uptake rates.

Our large studies, in which we collected detailed information on screening colonoscopy participants' medical histories and reasons for participating (1, 2) showed that the reasons for conducting screening colonoscopy mentioned by Hedemann may apply indeed, but only in a small minority of cases. And even if the offer of screening colonoscopy would only result in additional "indicated" colonoscopies being conducted, then this would also be a "positive side effect" associated with the screening offer.

Wenderlein points out an important issue—namely, the role of gynecologists in bowel cancer screening. A large proportion of tests for fecal occult blood—for whose effectiveness there is substantially more evidence (3) than for the rectal-digital examinations he cites—are being conducted in the setting of gynecological screenings.

Mühlhauser points to her own work in her letter, where she discusses criteria for compiling patient information on cancer screening examinations. She claims these to be the standard for communicating the results of cancer screening examinations and that they

could be adapted for scientific studies. We thank the author for pointing this out; however, we intentionally based the presentation of our results on clearly defined study objectives and internationally established standards for scientific publications. The additional information that Mühlhauser is asking for was not the focus of our study. Furthermore, it would be possible to estimate the requested data only on the basis of far reaching additional assumptions, whose detailed presentation and discussion would not have made sense, nor even been possible, within the word limit imposed by Deutsches Ärzteblatt. With regard to serious complications we'd like to draw attention to page 757 of our article.

DOI: 10.3238/arztebl.2011.0172

REFERENCES

1. Hundt S, Haug U, Brenner H: Comparative evaluation of immunochemical fecal occult blood tests for colorectal adenoma detection. *Ann Intern Med* 2009;150:162–9.
2. Brenner H, Tao S, Haug U: Low-dose aspirin use and performance of immunochemical fecal occult blood tests. *JAMA* 2010; 304: 2513–20.
3. Hewitson P, Glasziou P, Watson E, Towler B, Irwig L: Cochrane systematic review of colorectal cancer screening using the fecal occult blood test (hemoccult): an update. *Am J Gastroenterol*. 2008; 103: 1541–9.
4. Brenner H, Altenhofen L, Hoffmeister M: Eight years of colonoscopic bowel cancer screening in Germany: Initial findings and projections, *Dtsch Arztebl Int* 2010; 107(43): 753–60.

Prof. Dr. med. Hermann Brenner*
Dr. rer. soc. Lutz Altenhofen
Dr. sc. hum. Michael Hoffmeister

*Abteilung Klinische Epidemiologie und Altersforschung
 Deutsches Krebsforschungszentrum
 Bergheimer Str. 20
 69115 Heidelberg, Germany
 h.brenner@dkfz-heidelberg.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.