Excision of specific DNA-sequences from integrated retroviral vectors via site-specific recombination

Jörg Bergemann, Klaus Kühlcke, Boris Fehse, Ilka Ratz, Wolfram Ostertag and Heinz Lothar*

Heinrich-Pette-Institut für Experimentelle Virologie und Immunologie an der Universität Hamburg, Martinstraße 52, 20251 Hamburg, Germany

Received July 3, 1995; Revised and Accepted October 2, 1995

ABSTRACT

Vectors for gene transfer and gene therapy were developed which combine the advantages of the integrase and recombinase systems. This was achieved by inserting two *loxP* sites for specific DNA excision into an MESV based retroviral vector. We show that this 'retroviral *lox* system' allows the infection of cells and the expression of transferred genes. In addition, we constructed an efficient retrovirus-based expression system for a modified Cre recombinase. Functional tests for DNA excision from integrated retroviral *lox* vectors were performed by the use of a negative selectable marker gene (thymidine kinase). Cre expression in cells infected with retroviral *lox* vectors and subsequent BrdU selection for cells in which site-specific recombination has occurred results in large numbers of independent cell clones. These results were confirmed by detailed molecular analysis. In addition we developed retroviral suicide vectors in which the enhancer/promoter elements of both LTRs were replaced by *lox* sequences. We show that *lox*-sequences located in the LTRs of retroviral vectors are stable during retroviral replication. Potential applications of this system would be the establishment of revertants of retrovirus-infected cells by controlled excision of nearly the complete proviral DNA.

INTRODUCTION

Retroviral vectors are commonly used for gene transfer and gene therapy (1–3). They are derivatives of retroviruses which integrate DNA into host genomes as an obligatory step during their life cycle (4). Retroviral vectors infect a wide range of mitotically active cells by binding to specific receptors. The viral RNA is reverse transcribed after infection and the proviral DNA is integrated into the host genome. The virus encoded DNA endonuclease, the integrase, catalyzes this process (5,6). At present no consensus sequence for integration has been observed, however a preference for DNase I hypersensitive, transcriptionally active regions has been described (7,8). Depending on the

multiplicity of infection, one or multiple copies of the retroviral vector are integrated into the host genome (9–11). In comparison to other DNA transfer methods, the specific enzymatic reaction decreases the risk of cell transformation (12) and leads to stable integration (13). Due to these advantages, retroviral vectors are the most widely used tool for gene transfer and gene therapeutic applications.

Systems for site-specific recombination have been used for genomic engineering of yeast (14–15), plant (16–22) and animal cells (23–35). These systems include Cre-*lox* from bacteriophage P1, FLP-*FRT* from *Saccharomyces cerevisiae* and R-*RS* from *Xygosaccharomyces rouxii* (reviewed in 36). The 38 kDa Cre recombinase from bacteriophage P1 catalyzes site-specific recombination between 34 bp repeats termed *loxP* (37,38). These *lox* sites contain two 13 bp inverted repeats separated by an 8 bp asymmetrical core region. Upon binding to the inverted repeats, Cre synapses with a second *lox* site and cleaves the DNA in the spacer region to initiate strand exchange with the synapsed *lox* partner. Site-specific recombination results in excision or inversion of intervening DNA. This reaction depends on the orientation of the two *lox* sites (head to tail or head to head). No additional factors are required in the recombination reaction (39,40).

In this article, we describe a system for the specific excision of DNA fragments from integrated retroviral vectors. This system allows the complete silencing of retrovirally transferred genes by the excision of their ORFs. We replaced the enhancer/promoter elements of retroviral LTRs by *lox* sites and showed that these *lox* sites are stable during retroviral replication. We show that negative selectable marker genes allow the exclusive selection of cells in which excision has occurred. Potential applications of the system described here are discussed.

MATERIALS AND METHODS

Recombinant DNA

The plasmids pJBlox7 and pJBCre31 were constructed using standard methods (41). The relevant structures are depicted in Figure 1. A *Bam*HI-*Kpn*I *lox* fragment from plasmid pED32A (D. Ow) was used for the construction of pDB1 and inserted into pUC19. An additional *lox* site (pED32A) was added in a head to tail orientation as a *Bam*HI-*Sal*I *lox* fragment to create pDB5. The

* To whom correspondence should be addressed

Figure 1. Experimental design for a Cre mediated excision of DNA from integrated retroviral vectors. Schematic representation of an integrated retroviral vector with the neomycin resistance gene (neo) as a positive selectable marker and the herpes simplex virus type 1 thymidine kinase gene (HSV-1 TK) as a negative selectable marker. The HSV-1 TK gene is flanked by two *lox* sites in a head to tail orientation (A). The recombination event is drawn schematically in (B). The modified ORF for the Cre recombinase in the expression plasmid pJBCre31 is indicated by different rectangles. The integrated retroviral vector after site-specific excision of DNA is shown in (C).

HSV-TK expression cassette from plasmid pIC19R/MC1-TK (42) was inserted as a blunted *XhoI*-*HindIII* fragment between the two *lox* sites (*Bam*HI/blunt) of pDB5. This expression cassette contains the herpes simplex virus thymidine kinase gene driven by the promoter and a duplicated enhancer of a polyoma virus. The *lox-tk-lox* fragment from pDB6 was isolated as a *PvuII* fragment and inserted into the blunted *Bam*HI-*Eco*RI sites of the retroviral vector p50-Mneo (C. Stocking). This newly created retroviral vector pJBlox7 contains the ORF for the thymidine kinase gene in an opposite orientation. Deletions in *lox* sites occurred frequently during construction of these plasmids. This necessitated the performance of transformations in an *E. coli* SURE strain (Stratagene) and the incubation of cells at 28°C.

To create pJBCre31, the sequence for a modified ORF of the Cre recombinase was amplified by PCR using pMM23 (D. Ow) as template. Primers used for PCR were: DB14 5'-ATATTGGTACCTACCTACCATGGTGCCCAAGAA-GAAGAGGAAGGTGTCCAATTTACTGACCG-3' and DB2myc 5'-CATGCTCTAGATCAGTTCAAGTCTTCTTCA-GAAATAAGCTTTTGTTCGTTATCGCCATCTTCCAGCAG-GC-3'. PCR amplification yielded a modified ORF for the Cre recombinase with an N-terminal added Kozak sequence (43) and a nucleus localization sequence (44). A 9E10 epitope for immunological detection (45) was added at the C-terminus. The amplified DNA was cloned as a *KpnI*-*XbaI* fragment into pSP72 to create pSI1#13. A *KpnI*-*XhoI* fragment of pSI1#13 was inserted into pHB40p to create pSI1#13D. This plasmid was used for high-level expression in the T7-system. An *Eco*RI-*Hind*III fragment of pSI1#13 was inserted into pUC-M5neo (46) replacing the neo^R gene. The resulting vector pJBCre31 was used for expression in animal cell lines.

To generate the self-inactivating retroviral vector pJBVE6, the promoter/enhancer elements of the retroviral 3'LTR were replaced by a *lox* site. The promoter/enhancer elements in the U3 region of the 3'LTR of plasmid p5N-M (C. Stocking) were replaced by a *Bam*HI-*SstI* *lox* fragment from plasmid pED32a (D. Ow). The hyg/tyk expression cassette from plasmid tgCMV/hygtk was then inserted as a *NotI* fragment. The opposite orientation of the ORF for the hyg/tyk fusion protein was determined by restriction analysis. The resulting vector, pJBVE6, contains a wild-type (wt) 5'LTR and a 3'LTR with a *lox* site instead of promoter/enhancer elements. The promoter/enhancer elements in the wt 5'LTR were replaced during retroviral replication by the modified U3 region of the 3'LTR. This leads to an inactivated provirus without promoter/enhancer elements in both LTR's.

Culture, electroporation and retroviral infection of animal cells

GP-envAm12 helper cells (47) and RAT-2 fibroblasts (48), were maintained in DMEM containing 10% FCS supplemented with glutamine (2 mM). Electroporation was performed as described (49). GP-envAm12 cells and RAT-2 cells (2×10^6) were electroporated with 10 µg DNA (250 V and 1050 µF). Neo^R cells were selected with 400 µg/ml G418 for ≥ 2 weeks. BrdU^R cells were selected with 100 µg/ml BrdU for ≥ 2 weeks. Retroviral infection (50) was performed with different dilutions of supernatants from helper cell lines in the presence of 8 µg/ml polybrene. DNA isolation was performed by incubating 10^7 cells over night in 5 ml lysis buffer (containing 0.5% SDS, 300 µg/ml proteinase K in TE-buffer) and subsequent isopropanol precipitation.

Molecular analysis

In vitro DNA amplification was performed using a Biometra Temperature Cycler with a heated cover (Biometra). *Taq* polymerase was added after 10 min of denaturation followed by 40 cycles of denaturation (94°C, 1 min), annealing (62°C, 1 min) and extension (72°C, 2 min). The reaction (0.5 µg genomic template) was performed using standard conditions recommended by the polymerase supplier (Boehringer). Primers used for PCR analysis were:

DB3, 5'-GAGTCAAAACTAGAGCC-3';
DB10, 5'-GATCCTCCAGCGCGGGG-3';
DB20, GCCATTTTGCAAGGCATGG and
DB21, CTAGTTGTGGTTTGTC.

PCR fragments were cloned into the *XcmI* sites of vector pXcmKn12 (51). Southern analysis was performed by blotting fractionated DNA onto nylon membranes (Biodyne B). Filters were hybridized and washed using conditions recommended by the membrane supplier. Radioactive labelling of DNA fragments was performed by primer extension as described previously (52). Nucleotide sequences were determined by the dideoxynucleotide chain termination method of Sanger *et al.* (53). Alkali-denatured plasmid DNA was sequenced using T7 DNA polymerase (T7 sequencing Kit, Pharmacia).

RESULTS

Functional test of pJBlox7 and pJBCre31

A retrovirus based Cre-*lox* excision system was designed that was made up of two components (Fig. 1). A DNA excision cassette, including a gene for expression flanked by two *loxP* sites, was

Figure 2. Cre mediated BrdU resistance of pJB7#7 cells. RAT pJB7#7 cells were electroporated with plasmid pJBCre31 (+CRE) or the control plasmid pUC-M5neo (-Cre) and plated in 1:200 (A), 1:400 (B) and 1:800 (C) dilutions. BrdU selection was started 3 days after electroporation, methylene blue staining was carried out 2 weeks after electroporation.

inserted into an MESV based retroviral vector (pJBlox7). The second component (pJBCre31) was a retrovirus based expression vector for a modified Cre recombinase (see Materials and Methods). The function of the Cre-*lox* excision system was checked in prokaryotic cells and *in vitro*. This was necessary as we often obtained mutated *lox* sites during cloning. The modifications of the ORF for the Cre recombinase made it also necessary to examine the function of the enzyme.

The Cre expression strain BNN132 (Clontech) was transformed with plasmid pJBlox7. The plasmid DNA was isolated and checked by restriction analysis for recombination (54). We could demonstrate with DNA restriction analysis that in eight of 10 clones a complete excision of the tk-gene has occurred. In two of the 10 clones the excision was nearly complete (data not shown). Sequence analysis showed no base exchanges in the *lox* sites after recombination. This indicated that the pJBlox7 vector contained a functional *lox* system for DNA excision.

The modified Cre recombinase (pJBCre31) was functionally tested after expression by the T7-system. Plasmid pSI1#13D (see Materials and Methods) was transformed into the BL21DE3 strain (Cre⁻) and the modified Cre recombinase was expressed (55,56). High-level expression was checked by PAGE and immunological detection (45) using the monoclonal 9E10-antibody (Dianova). Expressed recombinase was prepared by electroelution and functionally tested *in vitro* with pOCUS-1 plasmids using conditions recommended by the supplier (AGS-company). This yielded 2×10^6 transformants per μg DNA, showing that the modified Cre recombinase works with high efficiency.

Selection for clones after site-specific recombination

A vector with a positive and a negative selectable marker (pJBlox7) was used to show the excision of specific DNA fragments from retroviral vectors. The strategy to detect the specific DNA excision is shown in Figure 1. The neomycin resistance gene enables selection for integration of the retroviral DNA into animal cell lines. Negative selection against thymidine kinase allows the establishment of cell clones in which site-specific recombination has occurred.

RAT-2 target cells (which do not express endogenous Tk) were infected with retroviral vectors produced by GP-envAm12 helper

Figure 3. Cre mediated BrdU resistance as a function of time between electroporation and start of selection. Cells were electroporated at day 0 and BrdU selection was started on various days after electroporation. Methylene blue staining was carried out 2 weeks after electroporation. y-axis shows percentage of clones relative to number of clones obtained without BrdU selection (cloning efficiency was 10%).

cells. Neomycin resistant clones were isolated by G418 selection. The function of the thymidine kinase gene was proven by BrdU sensitivity. These results indicate that the retroviral *lox* vector allows infection of cells and expression of the transferred genes. The resulting cell lines RAT pJB7#2 and RAT pJB7#7 were electroporated with pJBCre31 or the control plasmid pUC-M5neo, and cultivated in serial dilutions. The BrdU selection was started three days after electroporation. This period of time was chosen to ensure removal of thymidine kinase polypeptides from cells in which site-specific excision had occurred. The best time point to start BrdU selection was also determined (see below). Colonies were stained with methylene blue and counted after 2 weeks of BrdU selection. A typical result of this approach is shown in Figure 2. We were able to isolate a sufficient number of BrdU resistant clones using this approach. The electroporation with pJBCre31 resulted in 20-fold more BrdU resistant clones than transformation with the control plasmid pUC-M5neo. This ratio is sufficient to use this method to excise specific DNA fragments from retroviral vectors. The BrdU resistant clones obtained in the control could be explained by silencing of the retroviral vector or mutations in the thymidine kinase gene.

The efficiency of the retrovirus based Cre expression plasmid pJBCre31 was compared to a standard SV40 expression vector using the same conditions for electroporation and selection of BrdU resistant clones. In these tests we obtained significantly increased (10-fold higher) recombination efficiency by using vector pJBCre31. This shows that the plasmid pJBCre31 works with high efficiency as a Cre recombinase supplier in animal cells.

Efficiency of recombination in the pJBlox7/pJBCre31-system

The frequency of specific DNA excision by Cre as a function of the time before BrdU selection was tested. To do this we

Figure 4. (A) Autoradiogram of a Southern blot analysis of PCR amplifications. The primer pair DB3/DB10 was used for amplification on different templates. Lane 1, no template/PCR control; lane 2, pJB7 DNA isolated from a prokaryotic Cre producer (size marker for the expected DNA fragment after excision); lane 3, pJB7 DNA; lane 4–12, genomic DNA isolated from RAT pJB7#7 Cre1–Cre9 cells; lane 13–16 genomic DNA isolated from RAT pJB7#7 Co1–Co4 cells. (B) Location of PCR fragments and Southern probe relative to recombinant and non recombinant pJB7 DNA.

determined the number of BrdU resistant clones obtained after BrdU addition on different days after transformation with the Cre expression plasmid. As shown in Figure 3, the number of BrdU resistant clones increased with time between transformation and start of selection. We obtained a maximum of >20% BrdU resistant clones, relative to the total amount of unselected clones when BrdU selection was started 4 days after electroporation. This showed that in one out of five clonogenic cells, site-specific recombination had occurred. The efficiency of electroporation in RAT-2 fibroblasts was tested in an X-gal assay (data not shown) using 10 μ g of pCMV β (Clontech). In this assay we obtained 5–10% blue-stained cells which indicates a total electroporation efficiency for RAT-2 cells of ~20% (49). The correlation between efficiency of electroporation and efficiency of recombination suggests that in the majority of the cells containing the Cre-plasmid recombination has occurred.

Molecular analysis for specific excision of DNA from retroviral *lox* vectors

Thirteen independent cell lines were isolated by BrdU selection as described and tested for excision at the DNA level. Nine cell lines (RAT pJB7#7 Cre1–Cre9) were derived from an electroporation with the Cre expression plasmid pJB Cre31. Four cell lines (RAT pJB7#7 Co1–Co4) were derived from an electroporation with the control plasmid pUC-M5neo. PCR analysis was performed to detect the recombination event in DNAs isolated from cell lines RAT pJB7#7 Cre1–Cre9. The use of the primers DB3 and DB10, which bind 5' and 3' of the *lox*-thymidine kinase-*lox* fragment lead to amplification of 0.5 kb fragments if site-specific recombination has occurred (Fig. 4A). The specificity of the amplified signal was determined by Southern analysis. As shown in Figure 4A the expected signal was obtained in all

nine cell lines. This indicates that BrdU resistance is caused by specific Cre-mediated DNA excision. These 0.5 kb fragments were also isolated, cloned and sequenced and found to contain the expected sequence with an intact *lox* site. No signals for the recombination event were obtained in DNAs derived from cell lines RAT pJB7#7 Co1–Co4 (Fig. 4A). The presence of the tk-gene in DNAs derived from cell lines RAT pJB7#7 Co1–Co4 was determined by PCR analysis of the C-terminal part of the tk-gene (data not shown). Additional genomic Southern blot analysis was performed to test three cell lines electrotransformed with the Cre expression plasmid for complete DNA excision (Fig. 5). In all three BrdU resistant clones only the recombinant DNA's without signals at the size corresponding to non-recombined DNA were found. These data show that the analyzed cell lines underwent complete and precise recombination.

Generation of retroviral suicide vectors containing *lox* sites in both LTR's

The retroviral vector pJBVE6 contains a wt 5'LTR, a hyg/tk expression cassette and a 3'LTR in which a *lox* site replaces the promoter/enhancer elements. This vector was used for electroporation of GP-envAM12 helper cells. The expression of the retroviral vector in the helper cell line was controlled by the 5' wt LTR. Hyg^R cells were selected with 200 μ g/ml hygromycin for at least two weeks. We used supernatants from two independent cell lines (GPE-JBVE6#2 and GPE-JBVE6#4) for infection of RAT-2 target cells. Selection for infected cells was performed with 800 μ g/ml hygromycin. During infection of the target cells the retroviral replication leads to the replacement of the wt 5'LTR sequences by the modified 3'LTR sequences. This results in a provirus without promoter/enhancer elements in both LTR's, so called 'self inactivating vector' (57). As shown in Figure 6, two

Figure 5. Autoradiogram of a Southern blot analysis. *SacI* digest of genomic DNA separated on a 0.8% agarose gel. Probed with a radioactive labeled 1.9 kb *SacI* fragment from pJBlox 7 DNA isolated from a prokaryotic Cre producer. Lane 1, DNA from RAT pJBlox7#7 cells, lane 2–4, DNA from RAT pJBlox7#7-Cre1, -Cre2 and -Cre3 cells. The expected sizes for DNA fragments in which excision has/has not occurred are indicated by arrows (rec./no rec.)

Figure 6. (A) Organisation of retroviral vector pJBVE6 before and after replication. The binding sites for primer DB20 and DB21 and the expected sizes for the amplified products are indicated. (B) Autoradiogram of a Southern blot analysis of PCR amplifications probed with a radioactively labelled wt LTR sequence. The primer pair DB20/DB21 was used for amplification on different templates: lane 1, genomic DNA isolated from cell line RAT-JBVE6#2; lane 2, genomic DNA isolated from cell line RAT-JBVE6#4; lane 3 and 4, DNA from plasmid pJBVE6; lane 5, PCR control without template DNA. The obtained sizes for the amplifications on modified LTR sequences (1.1 kb) and wild type LTR sequences (1.4 kb) are indicated by arrows.

infected cell lines (RAT-JBVE6#2 and RAT-JBVE6#4) were used for molecular analysis. PCR analysis was performed to answer the question, whether the U3 region of the 5'LTR is completely replaced by the modified U3 region of the 3'LTR. For this analysis one primer inside the LTR (DB20) and one primer outside the LTR (DB21) was used. The replacement of the wt U3 region by the modified U3 region leads to a shortened LTR sequence. By using the primer pair DB20 and DB21 a wt LTR template DNA

leads to amplification of 1.4 kb fragments and the modified LTR template DNA leads to amplification of 1.1 kb fragments. In both infected cell lines we detected the expected signals for a modified 5'LTR sequence. The amplified DNA fragments were isolated from cell line RAT-JBVE6#4 and inserted into the *XcmI* site of vector pXcmKn12. Sequence analysis showed the expected sequences for the modified LTR with an intact *lox* site. These results show that the promoter/enhancer elements of a retroviral vector can be replaced by *lox* sites. After excision of the self inactivating *lox* vector by Cre expression, only a few LTR sequences should remain in the host genome.

DISCUSSION

A DNA excision cassette including a gene for expression flanked by two *loxP* sites was inserted into an MESV-based retroviral vector. We have shown that this retroviral *lox* vector allows infection of cells and expression of the transferred genes. This system can thus be used for retroviral vector-mediated gene transfer. It should only be limited by the maximum of 8 kb DNA accepted by retroviral vectors (58). We have also shown that Cre-mediated excision allows selection of cell clones in which excision has completely occurred. This complete excision will be important in cell lines used for quantitative assays or gene therapy (see below). The Cre expression plasmid pJBCre31 works highly efficiently. We have calculated that in most if not all cells electroporated with pJBCre31 excision must have occurred. This allows selection for the site-specific recombination event by controlled Cre expression.

A general application of retroviral *lox* vectors is the use in temporary and regulated gene expression. This combines the advantages of the integrase and recombinase systems: a safe and stable integration of foreign DNA with the possibility to completely down regulate the gene of interest. The retroviral *lox* vector pJBlox7 could be used for genes with an easily detectable expression (e.g. receptors). This system could also be used in gene therapy for complete excision of the marker genes in proviral DNA (LTR; gene for therapy; *lox-hyg/tk-lox*; LTR.). By using both negative and positive selectable marker genes (59), the integration of the retroviral vector as well as the excision of the marker gene can be detected. This leads to the selection of cells only expressing the gene of interest without additional foreign gene products which could elicit an immune response. For a selectable excision of retrovirally transferred genes the system can be modified by flanking the gene of interest and the selectable marker gene by *loxP* sites. These vectors could be used in hit-and-run experiments (60/61) to prove if the expression of a regulatory protein is necessary for the maintenance of a process initiated by that protein.

We have shown that *lox* sites located in the LTRs of retroviral vectors are stable during retroviral replication. By using the self inactivating vector pJBVE6, the enhancer/promoter elements of both LTRs were replaced by *lox* sequences. Due to the loss of LTR enhancer sequences the risk for insertional mutagenesis is minimized. We predict that after site specific recombination only a single *lox* site flanked by a few LTR sequences remains in the host genome. This *lox* site is safely and stably integrated via retroviral transfer and can be used for additional recombination events. In addition this method should allow the generation of revertants of retrovirus-infected cells by controlled excision of

nearly the complete proviral DNA. Further applications in gene therapy and studies for gene regulation are thus numerous.

ACKNOWLEDGEMENTS

We gratefully acknowledge Carol Stocking and David Ow for helpful comments and providing plasmids. We thank Andy Scutt for critical reading of the manuscript. This work was supported by grants from the BMFT (SUZ 0310718) and Deutsche Krebshilfe (W24/92/054). The Heinrich-Pette-Institut is financially supported by Freie und Hansestadt Hamburg and Bundesministerium für Gesundheit.

REFERENCES

- 1 Miller, A.D. (1992) *Nature*, **357**, 455–460.
- 2 Mulligan, R.C. (1993) *Science*, **260**, 926–923.
- 3 Vile, R. and Russell, S.J. (1994) *Gene Therapy*, **1**, 88–98.
- 4 McLachlin, J.R., Cornetta, K., Eglitis, M.A. and Anderson, W.F. (1990) *Prog. Nucleic Acids Res. Mol. Biol.*, **38**, 91–135.
- 5 Panganiban, A.T. and Temin, H.M. (1984) *Proc. Natl. Acad. Sci. USA*, **81**, 7885–7889.
- 6 Whitcomb, J.M. and Hughes, S.M. (1992) *Annu. Rev. Cell Biol.*, **8**, 275–306.
- 7 Rohdewohld, H., Weiher, H., Reik, W., Jaenisch, R. and Breindl, M. (1987) *J. Virol.*, **61**, 336–343.
- 8 Mooslehner, K., Karls, U. and Harbers, K. (1990) *J. Virol.*, **64**, 3056–3058.
- 9 Hwang, L.-H.S. and Gilboa, E. (1984) *J. Virol.*, **50**, 417–424.
- 10 Grez, M., Zörnig, M., Nowock, J. and Ziegler, M. (1991) *J. Virol.*, **65**, 4691–4698.
- 11 Beck-Engeser, G., Stocking, C., Just, U., Albritton, L., Dexter, M., Spooner, E. and Ostertag, W. (1991) *Hum. Gene Ther.*, **2**, 61–70.
- 12 Moolten, F.L. and Cupples, L.A. (1992) *Hum. Gene Ther.*, **3**, 479–486.
- 13 Miller, D.G., Adam, M.A. and Miller, A.D. (1990) *Mol. Cell. Biol.*, **10**, 4239–4242.
- 14 Sauer, B. (1987) *Mol. Cell. Biol.*, **7**, 2087–2096.
- 15 Cregg, J.M. and Madden, K.R. (1989) *Mol. Gen. Genet.*, **219**, 320–323.
- 16 Dale, E.C. and Ow, D.W. (1990) *Gene*, **91**, 79–85.
- 17 Odell, J., Caimi, P., Sauer, B. and Russell, S. (1990) *Mol. Gen. Genet.*, **223**, 369–378.
- 18 Maeser, S. and Kahmann, R. (1991) *Mol. Gen. Genet.*, **230**, 170–176.
- 19 Onouchi, H., Yokoi, K., Machida, C., Matsuzaki, H., Oshima, Y., Matsuoka, K., Nakamura, K. and Machida, Y. (1991) *Nucleic Acids Res.*, **19**, 6373–6378.
- 20 Lyznik, L.A., Mitchell, J.C., Hirayama, L. and Hodges, T.K. (1993) *Nucleic Acids Res.*, **21**, 969–975.
- 21 Lloyd, A.M. and Davis, R.W. (1994) *Mol. Gen. Genet.*, **242**, 653–657.
- 22 Albert, H., Dale, E.C., Lee, E. and Ow, D.W. (1995) *Plant J.*, **7**(4), 649–659.
- 23 Sauer, B. and Henderson, N. (1989) *Nucleic Acids Res.*, **17**, 147–161.
- 24 Golic, K.G. and Lindquist, S. (1989) *Cell*, **59**, 499–509.
- 25 O'Gorman, S., Fox, D.T. and Wahl, G.M. (1991) *Science*, **251**, 1351–1355.
- 26 Morris, A.C., Schaub, T.L. and James, A.A. (1991) *Nucleic Acids Res.*, **19**, 5895–5900.
- 27 Konsolaki, M., Sanicola, M., Kozlova, T., Liu, V., Arpa, B., Savakis, C., Gelbhart, W.M. and Kafatos, F.C. (1992) *New Biol.*, **4**, 551–557.
- 28 Orban, P.C., Chui, D. and Marth, J.D. (1992) *Proc. Natl. Acad. Sci. USA*, **89**, 6861–6865.
- 29 Lakso, M., Sauer, B., Mosinger, A., Jr., Lee, E.C., Fanning, R.W., Yu, S.-H., Mulder, K.L. and Westphal, H. (1993) *Proc. Natl. Acad. Sci. USA*, **89**, 6232–6236.
- 30 Gu, H., Zou, Y.-R. and Rajewsky, K. (1993) *Cell*, **73**, 1155–1164.
- 31 Fiering, S., Kim, C.G., Epner, E.M. and Goodwin, M. (1993) *Proc. Natl. Acad. Sci. USA*, **90**, 8469–8473.
- 32 Struhl, G. and Basler, K. (1993) *Cell*, **72**, 327–340.
- 33 Baubonis, W. and Sauer, B. (1993) *Nucleic Acids Res.*, **21**, 2025–2029.
- 34 Gu, H., Marth, J.D., Orban, C., Mossmann, H. and Rajewsky, K. (1994) *Science*, **265**, 103–106.
- 35 Disanto, J.P., Müller, W., Guy-Grand, D., Fischer, A. and Rajewsky, K. (1995) *Proc. Natl. Acad. Sci. USA*, **92**, 377–381.
- 36 Kirby, N.J., Snaith, M.R. and Murray, A.H. (1993) *Trends Genet.*, **9**, 413–421.
- 37 Sternberg, N. and Hamilton, D. (1981) *J. Mol. Biol.*, **150**, 467–468.
- 38 Sauer, B. and Henderson, N. (1988) *Proc. Natl. Acad. Sci. USA*, **85**, 5166–5179.
- 39 Abremski, K., Hoess, R. and Sternberg, N. (1983) *Cell*, **32**, 1301–1311.
- 40 Abremski, K. and Hoess, R. (1984) *J. Biol. Chem.*, **259**, 1509–1514.
- 41 Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- 42 Mansour, S.L., Thomas, K.R., and Capecchi, M.R. (1988) *Nature*, **336**, 348–352.
- 43 Kozak, M. (1986) *Cell*, **44**, 283–292.
- 44 Moore, M.S. and Blobel, G. (1992) *Cell*, **69**, 939–950.
- 45 Kolodziej, P.A. and Young, R.A. (1991) *Methods Enzymol.*, **194**, 508–519.
- 46 Laker, C., Stocking, C., Bergholz, U., Hess, N., De Lamarier, J.F. and Ostertag, W. (1987) *Proc. Natl. Acad. Sci. USA*, **84**, 8458–8462.
- 47 Markowitz, D., Goff, S. and Bank, A. (1988) *Virology*, **167**, 400–406.
- 48 Topp, W.C. (1981) *Virology*, **113**, 408–411.
- 49 Baum, Ch., Forster, P., Hegewisch-Becker, S. and Harbers, K. (1994) *BioTechniques*, **6**, 1058–1062.
- 50 Beck-Engeser, G., Stocking, C., Just, U., Albritton, L., Dexter, M., Spooner, E. and Ostertag, W. (1991) *Hum. Gene Ther.*, **2**, 61–70.
- 51 Cha, J., Bishai, W. and Chandrasekaran, S. (1993) *Gene*, **136**, 369–370.
- 52 Feinberg, A.P. and Vogelstein, B. (1983) *Anal. Biochem.*, **132**, 6–13.
- 53 Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA*, **74**, 5463–5467.
- 54 Elledge, S.J., Mulligan, J.T., Rutter, S.W., Spillwood, M. and Davis, R.W. (1991) *Proc. Natl. Acad. Sci. USA*, **88**, 1731–1735.
- 55 Studier, F.W. and Moffatt, B.A. (1986) *J. Mol. Biol.*, **189**, 113–130.
- 56 Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) *Methods Enzymol.*, **185**, 60–89.
- 57 Yu, S.F., Rüden, T., Kanthoff, P.W., Garber, C., Seiberg, M., Rüther, U., French Anderson, W., Wagner, E.F. and Gilboa, E. (1986) *Proc. Natl. Acad. Sci. USA*, **83**, 3194–3198.
- 58 Miller, A.D. (1992) *Curr. Top. Microbiol. Immunot.*, **158**, 60–89.
- 59 Lupton, S.D., Brunton, L.L., Kalberg, V.A. and Overell, R.A. (1991) *Mol. Cell. Biol.*, **11**, 3347–3378.
- 60 Geldas, R., Roux, J., Pictet, R. and Grange, T. (1991) *Cell*, **67**, 977–986.
- 61 Zhang, C.C. and Bienz, M. (1992) *Proc. Natl. Acad. Sci. USA*, **89**, 7511–7515.