Published in final edited form as:

Mol Oral Microbiol. 2011 April ; 26(2): 117–126. doi:10.1111/j.2041-1014.2010.00602.x.

Characterization of competence and biofilm development of a *Streptococcus sanguinis* endocarditis isolate

Lin Zhu¹, Yongshu Zhang², Jingyuan Fan², Mark C. Herzberg^{2,3}, and Jens Kreth^{4,*}

¹ College of Dentistry, University of Oklahoma Health Sciences Center, Oklahoma City, OK 73104, USA

² Department of Diagnostic and Biological Sciences, School of Dentistry, University of Minnesota, Minneapolis, MN 55455, USA

³ Mucosal and Vaccine Research Center, Minneapolis Veterans Affairs Medical Center, Minneapolis, MN 55417, USA

⁴ Department of Microbiology and Immunology, University of Oklahoma Health Sciences Center, Oklahoma City, OK 73104, USA

Abstract

Streptococcus sanguinis is an oral commensal bacterium and endogenous pathogen in the blood, which generally is naturally competent to take up extracellular DNA. Regarded as a stress response, competence development enables *S. sanguinis* to acquire new genetic material. The sequenced reference strain SK36 encodes and expresses the genes required for competence (*com*) and uptake of DNA. Isolated from blood cultures of a confirmed case of infective endocarditis, strain 133–79 encodes all necessary *com* genes but is not transformable under conditions permissive for competence development in SK36. Using synthetic competence-stimulating peptides (sCSP) based on sequences of SK36 and 133–79 *comC*, both strains developed competence at similar frequencies in cross-transformation experiments. Furthermore, downstream response pathways are similar in strains SK36 and 133–79 since platelet aggregation and biofilm formation appeared unaffected by CSP. Collectively, the data indicate that strains SK36 and 133–79 respond to CSP similarly, strongly suggesting that endogenous production or release of CSP from 133–79 is impaired.

Introduction

Oral streptococci are constantly challenged by environmental stress. Saliva flow and food intake cause fluctuations in nutrient supply, pH, temperature and shear stress. The biofilm community adapts to dynamic conditions, therefore, by growth to high cell density and species diversity (Kreth *et al.*, 2009). Under biofilm growth conditions, competence development is an important mechanism of stress adaptation (Li *et al.*, 2002). Competence in streptococci seems to be a transient physiological state during which DNA is taken up from the environment (Morrison, 1997). DNA uptake as a consequence of competence development leads to horizontal gene transfer, genomic plasticity and the acquisition of new genetic traits.

*Corresponding author address: Department of Microbiology and Immunology, University of Oklahoma Health Sciences Center, BMSB 907, 940 Stanton L. Young Blvd., Oklahoma City, OK 73104. Phone: (405) 271-1202. Fax: (405) 271-3117. JKreth@ouhsc.edu.

The molecular details of competence development, including the identification of competence regulating genes, were initially analyzed in *Streptococcus pneumoniae* (for a recent review, see (Johnsborg & Havarstein, 2009)). In *S. pneumoniae*, competence development requires the production of the species-specific competence stimulating peptide (CSP) (gene *comC*), which is secreted into the environment by the ComAB transport proteins (Hui & Morrison, 1991). Environmental CSP is sensed by the two-component system ComDE; at a critical CSP threshold concentration a cellular response is triggered (Havarstein *et al.*, 1995, Pestova *et al.*, 1996). Accumulation of CSP results in quorum sensing, whereby early competence genes, including *comCDE*, *comAB* and *comX*, are transcribed and amplified in an auto-regulatory mechanism (Morrison & Lee, 2000). ComX, an alternative sigma subunit of the DNA polymerase, then initiates transcription of delayed competence genes required for DNA uptake, processing and chromosomal integration (Lee & Morrison, 1999, Luo *et al.*, 2003, Luo & Morrison, 2003). Associated with competence development in streptococci, CSP controls the expression of other genes not involved in horizontal gene transfer, including bacteriocin production (Kreth *et al.*, 2005) and biofilm development (Li *et al.*, 2002). The competence state has been renamed the X-state, reflecting the general stress adaptation mechanisms responding to CSP accumulation and high cell density (Claverys *et al.*, 2006).

In this report, we investigate competence development leading to DNA uptake in two different strains of *Streptococcus sanguinis*. *S. sanguinis* is an oral commensal bacterium and pioneer colonizer during oral biofilm formation (Xu *et al.*, 2007). Associated with oral health, *S. sanguinis* is considered harmless in the oral cavity; healthy individuals have significantly higher numbers than individuals with caries who have virtually no detectable *S. sanguinis* (Becker *et al.*, 2002). Indeed, *S. sanguinis* and cariogenic *S. mutans* are mutually antagonistic; acquisition of *S. mutans* is delayed when *S. sanguinis* is present in high abundance in human subjects (Caufield *et al.*, 2000). Upon entry into circulating blood, *S. sanguinis* bacteremia can cause severe infections in at-risk patients, such as infective endocarditis (Moreillon & Que, 2004). Infective endocarditis caused by oral streptococci remains associated with high morbidity and considerable mortality despite the use of antibiotics (Westphal *et al.*, 2009). A non-pathogenic commensal in the oral cavity and pathogen in the blood, the confounding lifestyles of *S. sanguinis* may be explained by environmental factors that elicit strain-specific differences in cell-to-cell signaling and related phenotypic characteristics.

In general, *S. sanguinis* is naturally competent to take up extracellular DNA (Turner *et al.*, 2009). The sequenced strain SK36, expresses the *com* genes required for competence development (*comCDE*) and uptake of DNA (*comYA comYD*) (Xu *et al.*, 2007). In contrast, strain 133-79, isolated from blood cultures of a confirmed case of infective endocarditis (Herzberg *et al.*, 1990), does not become competent in conditions permissive for competence development in SK36. In this study, we compared both strains for competence gene expression and DNA uptake in response to CSP administration. We provide experimental evidence that 133-79 encodes the *com*-genes and responds to exogenous CSP, but might not produce sufficient CSP by itself to signal for competence.

Materials and Methods

Bacterial Strains, growth conditions and chemicals

S. sanguinis strains SK36 (Xu *et al.*, 2007) and 133-79 (Herzberg *et al.*, 1990) were routinely grown in Bacto™ Todd-Hewitt (TH) broth (BD; Sparks, MD). Liquid cultures were incubated without agitation at 37°C in ambient air, agar plates were incubated in candle jars at 37°C. Trypticase™ soy agar plates with 5% sheep blood (TSA II) were purchased from BD and incubated in candle jars at 37°C. For selective cultures, kanamycin

was added to a final concentration of 400 µg/ml. Synthetic competence stimulating peptides, sCSPs, (sequence see Fig. 1) were custom-synthesized by ChemPep Inc. (Miami, FL). As assessed with HPLC, peptide purity was determined to be 98% for sCSP_{SK36} and 98.8% for sCSP₁₃₃₋₇₉. The peptides were dissolved in molecular grade water (1 mg/ml), aliquoted and stored at -20°C until further use.

Transformation assays

Transformation was done essentially as described previously (Paik *et al.*, 2005) with modifications. Briefly, overnight cultures were grown in TH broth containing 2.5% heat-inactivated horse serum (PAA Laboratories; Dartmouth, MA). The next day, cells were freshly inoculated in the same medium (1:25) and incubated until the cell density reached an absorption at 600 nm ($A_{600\text{nm}}$) of 0.08. The cultures were aliquoted into 1 ml portions and 8 µg/ml chromosomal DNA carrying a kanamycin resistance determinant (gene *aphA*) was added. The chromosomal DNA was isolated from a previously constructed SK36 strain (genotype *pKnB::aphA*, Kan^R; unpublished). Assay for competence monitored uptake of chromosomal DNA and chromosomal integration via homologous recombination. DNA concentration was measured using a NanoDrop spectrophotometer. At indicated times, sCSP was added to a final concentration of 0.5 µg/ml. Incubation continued for two hours at 37°C and cells were plated on selective TH plates. Transformation efficiency was calculated as the number of kanamycin-resistant CFUs relative to the number of CFUs on non-selective Todd-Hewitt agar.

RNA isolation, cDNA synthesis and real-time PCR

To isolate RNA, overnight cultures (1.6 ml) grown in TH medium were inoculated into 40 ml of fresh TH medium in a 50-ml conical tube. Growth was monitored spectrophotometrically at $A_{600\text{nm}}$. Mid-log phase cells ($A_{600\text{nm}} \sim 0.4$) were induced with homologous or heterologous sCSP (0.5 µg/ml final concentration) and incubated for 2 h to ensure a high yield of RNA for downstream application. Cells were collected by centrifugation at $11,000 \times g$ for 15 min at 4°C and resuspended in 778 µl RLT buffer (QIAGEN RNeasy Kit) and 222 µl nuclease free water. Cells were immediately disrupted twice for 1 minute each using Lysing Matrix B (MP Biomedicals; Solon, OH) and a FastPrep FP210 Homogenizer (Thermo Scientific). RNA was purified using a QIAGEN RNeasy Kit (QIAGEN; Valencia, CA) following the manufacturer's protocol. Isolated RNA was treated with DNase I (Invitrogen; Carlsbad, CA) to remove traces of chromosomal DNA. After the treatment, RNA samples were cleaned with the QIAGEN RNeasy Kit. The concentration and quality of RNA samples were confirmed using NanoDrop spectrophotometer measurements and gel electrophoresis.

cDNA was synthesized from 2 µg RNA using the SuperScript IITM Reverse Transcriptase and Random Primers (Invitrogen) according to manufacturer's instructions.

Specific transcripts were quantified with the comparative threshold cycle (C_t) method using the Bio-Rad MyiQ real-time PCR (RT-PCR) detection system (Hercules, CA). The RT-PCR protocol included one cycle of 95°C for 90 seconds, followed by 40 cycles of 95°C for 15 seconds and 60°C for 1 min. For Real-Time RT-PCR, oligonucleotide primers were designed with Beacon Designer 7 software (PREMIER Biosoft), which selects primers optimized for SYBR[®] green based PCR analysis. Primer sequences are listed in Table 1. Note that all primers could be used in both strains with equal efficiency, except for *comB* and *comC*, which required specific primers for each strain. Real-time PCR was performed using 1 x SYBR green master mix (Bio-Rad) according to the manufacturer's instructions. Changes of gene expression were calculated using the C_t method as follows: $C_t = C_t$ (target)

– C_t (housekeeping gene). The fold changes are calculated as 2^{-C_t} . The *gyrA* gene was used as the housekeeping reference gene.

Biofilm formation and quantification

Biofilm formation was measured using a modification of the crystal violet (CV) microtiter assay as reported (Ashby *et al.*, 2009).

Determination of transcription start sites

The FirstChoice RLM random amplification of cDNA ends (RACE) kit (Ambion) was used to determine transcription start sites essentially as described earlier (Okinaga *et al.*). RACE specific primers are listed in Table 1.

Sequencing

GenBank sequence accession numbers: GU943505, GU943506, GU943507, GU943508, GU943509, GU943510.

Statistical methods

Statistical analysis of two datasets was performed with QuickCalcs online calculators (<http://www.graphpad.com/quickcalcs/index.cfm>) using the *t*-test software to calculate mean and SD. Statistically significant differences were set at a *P* value of <0.05.

Results

Competence phenotype of *S. sanguinis* strain 133–79

Strain SK36 contains all genes required for competence development, uptake and incorporation of DNA into the chromosome (Xu *et al.*, 2007). When tested for competence with chromosomal DNA encoding a kanamycin resistance cassette, the natural transformation efficiency of strain SK36 was about 10^{-7} . In contrast, strain 133–79 was not transformable when tested without the addition of sCSP, suggesting that strain 133–79 is unable to develop natural competence under laboratory conditions.

Comparison of competence genes in *S. sanguinis* strains SK36 and 133–79

To determine whether strain 133–79 contains the genes required for competence development and DNA uptake, the chromosomal loci for *comCDE*, *comX* and *comYA-comYD* were sequenced and compared to the annotated sequence of strain SK36 (Xu *et al.*, 2007). The *comCDE*, *comX* operon, and the *comCDE* operon are present in 133–79. Each SK36 and 133–79 gene was more than 95% identical at the nucleotide level, with the exception of *comC*, showing only 85% identity (data not shown). By aligning the full-length peptide sequences, SK36 and 133–79 CSPs showed five exchanges. Three conservative replacements of leucine and valine with isoleucine as well as leucine with alanine, a polar to non-polar exchange from serine to leucine and a non-polar to polar exchange from glycine to aspartate (Fig. 1A). Three of the exchanges occurred in the peptide sequence defined to be the mature form.

The *comCDE* promoter sequences of SK36 and 133–79 are identical. To better characterize the promoter region of the *comCDE* operon (Fig. 1B), the transcriptional start site was determined by RACE PCR in both strains. The start was determined to be 105 bp upstream of the start-codon in both strains. A putative extended -10 region is located 8 bp upstream of the transcription start. No obvious -35 region could be located (Fig. 1C). When compared to SK36, the 133–79 competence gene sequence similarities suggest the ability to develop competence.

Strain-specific competence stimulating peptide (CSP) cross-induces competence development

To learn whether the *comC* gene of strain 133–79 encodes a non-functional CSP, CSPs of SK36 and 133–79 were synthesized as sCSP_{SK36} and sCSP_{133–79} using the published peptide sequence as reference (Gaustad & Havardstein, 1997), respectively. Addition of strain-specific sCSPs induced competence in the homologous strains. sCSP_{SK36} increased strain SK36 transformation efficiency about 4-fold (Fig. 2). When compared to SK36 induced with sCSP_{SK36}, addition of sCSP_{133–79} to strain 133–79 increased transformation efficiency 10-fold more. Using the heterologous sCSPs, cross-induction resulted in similar transformation efficiency suggesting that sCSP_{SK36} and sCSP_{133–79} are equally effective for SK36 or 133–79 (Fig. 2).

Effect of sCSP concentration and growth phase on competence

To test for concentration dependence, strain 133–79 and SK36 were incubated with increasing sCSP concentrations and transformability was tested. Both strains formed similar numbers of transformants per µg DNA when increasing concentrations of strain specific sCSP was added to the medium (Fig. 3A).

Next, we tested whether competence development in 133–79 and SK36 differs as a function of stage of growth and cell density. SK36 is naturally competent during early stage of growth (Fig. 3B), consistent with previous reports for *S. pneumoniae* (Morrison, 1997). Strain 133–79 did not develop competence during growth at any time or cell density.

Cross-induction of competence gene expression

SK36 responded to homologous sCSP by upregulating expression of *comX* about 10-fold, *comYA* about 7-fold, and about 20-fold when compared the non-induced controls (Fig. 4A to C, respectively). Similar to strain SK36, 133–79, responded to homologous sCSP by inducing expression of *comX* about 100-fold, *comYA* about 500-fold, and *comY* about 20-fold (Fig. 4D to F, respectively). After addition of heterologous sCSPs, expression of competence genes also increased in both strains (Fig. 4A to C for SK36 and D to F for 133–79, respectively). The transcriptional responses of SK36 and 133–79 to sCSPs were consistent with the results obtained for transformation efficiency (Fig. 2).

Comparison of competence gene expression in SK36 and 133–79

To compare the two strains for the expression levels of the competence genes, we first determined that the Real-Time RT PCR primers amplify the selected genes in both strains with equal efficiency. The *C_t* values of *gyrA* in both strains were similar. When compared, SK36 and 133–79 showed similar ratios of competence genes to *gyrA* (Table 2). Non-induced SK36 expressed *comC*, *comD*, *comX* and *comYA* at higher levels than non-induced 133–79. Addition of sCSP_{133–79} increased expression of *comX* and *comYA* in 133–79 above levels in non-induced SK36, whereas *comC* and *comD* were induced in 133–79 at lower levels than in non-induced SK36 (Fig. 5A to D).

Effect of cell density on the expression of competence genes

The relative expression of *comX* and *comYA* was compared in high cell-density cultures of 133–79 and SK36. Cells were grown on blood agar plates for 5 hours, harvested from the plates, and RNA was isolated. The expression of both genes was similar in 133–79 and SK36 (data not shown).

Increased biofilm formation by strain 133–79

We therefore tested biofilm development of both strains with and without addition of sCSP. sCSPs showed no influence on static biofilm formation by either strain on polystyrene surfaces. Strain 133–79, however, formed significantly denser biofilms than SK36 ($p = 0.0001$) (Fig. 5).

Discussion

Competence development is part of a CSP-mediated response mechanism to stress. The uptake and integration of DNA into the chromosome via homologous recombination might select for new, beneficial genetic traits and is the main source of horizontal gene transfer in streptococci (Havarstein, 2010). Competence development mutants show a reduced ability to take up or integrate DNA into the chromosome, form biofilms, and produce bacteriocins (Li *et al.*, 2002, Kreth *et al.*, 2005). These mutants might, therefore, be less competitive in a multispecies community as found in the oral biofilm. *S. sanguinis* is naturally competent, represented by strain SK36. To our surprise, we found an endocarditis isolate 133–79 to be non-transformable under laboratory conditions. We sought to understand why the two strains differed in competence.

The competence genes are functional in 133–79. To establish this conclusion, we sequenced the respective *com* genes in 133–79 and found them to be intact and almost identical on the nucleotide level when compared to SK36, confirming the ability of 133–79 to develop competence. When added exogenously, homologous (133–79) or heterologous (SK36) sCSPs induced expression of competence genes and transformability in strain 133–79.

Since 133–79 could respond to sCSPs, production or export of endogenous CSP was strongly suggested to be impaired in strain 133–79. The *comCDE* genes are encoded in an operon (Fig. 1B). The expression of *comC* is considerably lower in 133–79 than SK36, suggesting a much lower production of CSP. Since *comCDE* expression is regulated in an auto-regulatory circuit, CSP concentrations below a signaling threshold could explain the non-competent phenotype observed for 133–79.

The export of CSP in several streptococci requires the dedicated export system ComAB (Hui & Morrison, 1991, Claverys *et al.*, 2009). For *S. pneumoniae*, CSP export is limiting in competence development (Martin *et al.*, 2000). Interestingly, SK36 does not encode ComAB homologues, suggesting that an alternative CSP-export mechanism is used by *S. sanguinis* (Xu *et al.*, 2007). CSPs of SK36 and 133–79 lack the typical double glycine motif for Com *comCDE* AB transport, which is found in CSPs of other sequenced streptococci. Since the dedicated transport system has not yet been identified in *S. sanguinis*, we are unable to test if the low expression of is caused by impaired transport of CSP.

We therefore concentrated on regulation of the *comCDE* operon. Investigation of the promoter region showed no difference between SK36 and 133–79. No obvious -35 site could be identified. A search for LysRT type regulator binding sites did not show a ComE binding site upstream of the extended -10 region as describe for the *S. pneumoniae* ComE binding site (Ween *et al.*, 1999). Further experimental proof is required to show whether ComE interacts with the *comCDE* promoter and which region is covered by the bound protein. The promoter region was also searched for the presence of a *comX* box using the recently described sequences for *S. mutans* as template (Okinaga *et al.*). No *comX* binding site was identified. To consider whether the transcription start of the *comCDE* operon differs in SK36 and 133–79, RACE PCR was performed with CSP-induced and non-induced cells of both strains. The start site appeared the same in induced cells and uninduced controls. We

conclude therefore that the regulation of *comCDE* operon expression is the same in both strains.

In strain 133–79, the most likely explanation for the lack of autonomous competence development is impaired export of CSP and little or no CSP is available for auto-induction. Deletion of the *comC* gene in SK36 results in the same non-transformable phenotype, which could be rescued to wild type transformability when sCSP was added (T. Kitten, VCU School of Dentistry; personal communication). These data support the conclusion that that *S. sanguinis* has CSP-inducible competence development.

The transcriptional response of *S. gordonii* to CSP has been reported to change the expression of several surface proteins potentially involved in host cell interactions (Vickerman et al., 2007). Homologs of these surface proteins are found in *S. sanguinis*. We, therefore, tested whether CSP auto-induction changes surface proteins on *S. sanguinis* sufficiently to alter the ability to trigger platelet aggregation (reviewed in (Herzberg, 1996)). Autologous sCSP induction of SK36 does not affect the platelet aggregation response (data not shown). We have also tested whether CSP addition affects biofilm formation by *S. sanguinis*. Although strain 133–79 forms denser biofilms than SK36, this function was unaffected by CSP. Hence, we found no evidence that CSP could contribute to virulence by affecting biofilm development or platelet aggregation in *S. sanguinis*.

In summary, the non-transformable *S. sanguinis* 133–79 has the ability to develop competence when exogenous CSP is supplied. Why *S. sanguinis* 133–79 is naturally non-transformable appear to reflect differences in release or processing of CSP when compared with SK36, although there is no obvious advantage of this phenotype based on the similar platelet aggregation and biofilm formation traits. From an ecological standpoint, if strain 133–79 was originally part of a multispecies community including CSP-producing *S. sanguinis*, a classification as “cheater” might be possible. Strain 133–79 would take advantage of the heterologous produced CSP by other strains of *S. sanguinis* in the oral biofilm. It could therefore respond with full competence development, but without bearing the costly production of an excreted peptide. More detailed competition experiments in the future will answer this question.

Acknowledgments

The support from NIH/NIDCR grants R01DE08590 and R01DE11831 to MCH and 4R00DE018400 to JK is gratefully acknowledged. We thank Dr. Justin Merritt for helpful discussions.

References

- Ashby MT, Kreth J, Soundarajan M, Sivulu LS. Influence of a model human defensive peroxidase system on oral streptococcal antagonism. *Microbiology*. 2009; 155:3691–3700. [PubMed: 19684069]
- Becker MR, Paster BJ, Leys EJ, Moeschberger ML, Kenyon SG, Galvin JL, Boches SK, Dewhirst FE, Griffen AL. Molecular analysis of bacterial species associated with childhood caries. *J Clin Microbiol*. 2002; 40:1001–1009. [PubMed: 11880430]
- Caufield PW, Dasanayake AP, Li Y, Pan Y, Hsu J, Hardin JM. Natural history of *Streptococcus sanguinis* in the oral cavity of infants: evidence for a discrete window of infectivity. *Infect Immun*. 2000; 68:4018–4023. [PubMed: 10858217]
- Claverys JP, Martin B, Polard P. The genetic transformation machinery: composition, localization, and mechanism. *FEMS Microbiol Rev*. 2009; 33:643–656. [PubMed: 19228200]
- Claverys JP, Prudhomme M, Martin B. Induction of competence regulons as a general response to stress in gram-positive bacteria. *Annu Rev Microbiol*. 2006; 60:451–475. [PubMed: 16771651]

- Gaustad P, Havarstein LS. Competence-pheromone in *Streptococcus sanguis*. Identification of the competence gene *comC* and the competence pheromone. *Adv Exp Med Biol*. 1997; 418:1019–1021. [PubMed: 9331824]
- Havarstein LS. Increasing competence in the genus *Streptococcus*. *Molecular Microbiology*. 2010
- Havarstein LS, Coomaraswamy G, Morrison DA. An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae*. *Proc Natl Acad Sci U S A*. 1995; 92:11140–11144. [PubMed: 7479953]
- Herzberg MC. Platelet-streptococcal interactions in endocarditis. *Crit Rev Oral Biol Med*. 1996; 7:222–236. [PubMed: 8909879]
- Herzberg MC, Gong K, MacFarlane GD, Erickson PR, Soberay AH, Krebsbach PH, Manjula G, Schilling K, Bowen WH. Phenotypic characterization of *Streptococcus sanguis* virulence factors associated with bacterial endocarditis. *Infect Immun*. 1990; 58:515–522. [PubMed: 2137112]
- Hui FM, Morrison DA. Genetic transformation in *Streptococcus pneumoniae*: nucleotide sequence analysis shows *comA*, a gene required for competence induction, to be a member of the bacterial ATP-dependent transport protein family. *J Bacteriol*. 1991; 173:372–381. [PubMed: 1987129]
- Johnsborg O, Havarstein LS. Regulation of natural genetic transformation and acquisition of transforming DNA in *Streptococcus pneumoniae*. *FEMS Microbiol Rev*. 2009; 33:627–642. [PubMed: 19396959]
- Kreth J, Merritt J, Qi F. Bacterial and host interactions of oral streptococci. *DNA Cell Biol*. 2009; 28:397–403. [PubMed: 19435424]
- Kreth J, Merritt J, Shi W, Qi F. Co-ordinated bacteriocin production and competence development: a possible mechanism for taking up DNA from neighbouring species. *Mol Microbiol*. 2005; 57:392–404. [PubMed: 15978073]
- Lee MS, Morrison DA. Identification of a new regulator in *Streptococcus pneumoniae* linking quorum sensing to competence for genetic transformation. *JBacteriol*. 1999; 181:5004–5016. [PubMed: 10438773]
- Li YH, Tang N, Aspiras MB, Lau PC, Lee JH, Ellen RP, Cvitkovitch DG. A quorum-sensing signaling system essential for genetic competence in *Streptococcus mutans* is involved in biofilm formation. *J Bacteriol*. 2002; 184:2699–2708. [PubMed: 11976299]
- Luo P, Li H, Morrison DA. ComX is a unique link between multiple quorum sensing outputs and competence in *Streptococcus pneumoniae*. *Mol Microbiol*. 2003; 50:623–633. [PubMed: 14617184]
- Luo P, Morrison DA. Transient association of an alternative sigma factor, ComX, with RNA polymerase during the period of competence for genetic transformation in *Streptococcus pneumoniae*. *J Bacteriol*. 2003; 185:349–358. [PubMed: 12486073]
- Martin B, Prudhomme M, Alloing G, Granadel C, Claverys JP. Cross-regulation of competence pheromone production and export in the early control of transformation in *Streptococcus pneumoniae*. *Mol Microbiol*. 2000; 38:867–878. [PubMed: 11115120]
- Moreillon P, Que YA. Infective endocarditis. *Lancet*. 2004; 363:139–149. [PubMed: 14726169]
- Morrison DA. Streptococcal competence for genetic transformation: regulation by peptide pheromones. *Microb Drug Resist*. 1997; 3:27–37. [PubMed: 9109094]
- Morrison DA, Lee MS. Regulation of competence for genetic transformation in *Streptococcus pneumoniae*: a link between quorum sensing and DNA processing genes. *Res Microbiol*. 2000; 151:445–51. [PubMed: 10961457]
- Okinaga T, Xie Z, Niu G, Qi F, Merritt J. Examination of the *hdrRM* regulon yields insight into the competence system of *Streptococcus mutans*. *Mol Oral Microbiol*. 25:165–177. [PubMed: 20536745]
- Paik S, Senty L, Das S, Noe JC, Munro CL, Kitten T. Identification of virulence determinants for endocarditis in *Streptococcus sanguinis* by signature-tagged mutagenesis. *Infect Immun*. 2005; 73:6064–6074. [PubMed: 16113327]
- Pestova EV, Havarstein LS, Morrison DA. Regulation of competence for genetic transformation in *Streptococcus pneumoniae* by an auto-induced peptide pheromone and a two-component regulatory system. *Mol Microbiol*. 1996; 21:853–862. [PubMed: 8878046]

- Turner LS, Das S, Kanamoto T, Munro CL, Kitten T. Development of genetic tools for in vivo virulence analysis of *Streptococcus sanguinis*. *Microbiology*. 2009; 155:2573–2582. [PubMed: 19423626]
- Vickerman MM, Iobst S, Jesionowski AM, Gill SR. Genome-wide transcriptional changes in *Streptococcus gordonii* in response to competence signaling peptide. *J Bacteriol*. 2007; 189:7799–7807. [PubMed: 17720781]
- Ween O, Gaustad P, Havarstein LS. Identification of DNA binding sites for ComE, a key regulator of natural competence in *Streptococcus pneumoniae*. *Mol Microbiol*. 1999; 33:817–827. [PubMed: 10447890]
- Westphal N, Plicht B, Naber C. Infective endocarditis--prophylaxis, diagnostic criteria, and treatment. *Dtsch Arztebl Int*. 2009; 106:481–489. quiz 490. [PubMed: 19730718]
- Xu P, Alves JM, Kitten T, Brown A, Chen Z, Ozaki LS, Manque P, Ge X, Serrano MG, Puiu D, Hendricks S, Wang Y, Chaplin MD, Akan D, Paik S, Peterson DL, Macrina FL, Buck GA. Genome of the opportunistic pathogen *Streptococcus sanguinis*. *J Bacteriol*. 2007; 189:3166–3175. [PubMed: 17277061]

Fig. 1.

A) Alignment of CSP from SK36 and 133–79. Arrows indicate amino acid exchanges. Underlined sequence marks the mature peptide. B) *comCDE* operon organization. C) Promoter sequence of strain SK36 and 133–79 with putative regulatory elements. underlined = ribosome-binding-site; bold and underlined = putative extended -10 region; bold = start codon; bold with arrow = transcriptional start.

Fig. 2.

Transformation efficiency of SK36 and 133-79. Cells were grown to an $A_{600\text{nm}}$ of 0.08 in TH medium plus 2.5% heat-inactivated horse serum. 8 $\mu\text{g/ml}$ chromosomal DNA (KanR) and sCSP (0.5 $\mu\text{g/ml}$) was added, incubation continued for 2h at 37°C, and the cells were plated on selective TH plates. Efficiency was calculated as the number of kan-resistant CFUs relative to the number of CFUs on non-selective TH plates. Data presented are the means and standard deviations of three independent experiments.

Fig. 3.

Influence of various sCSP concentrations and the growth phase on transformability. A) Cells were grown to an $A_{600\text{nm}}$ of 0.08 in TH medium plus 2.5% heat-inactivated horse serum. 8 $\mu\text{g/ml}$ chromosomal DNA (KanR) and strain specific sCSP in different concentration was added, incubation continued for 2h at 37°C, and the cells were cells plated on selective TH plates. B) To test for growth phase dependent transformability, cells were grown to different cell densities in TH medium plus 2.5% heat-inactivated horse serum. 8 $\mu\text{g/ml}$ chromosomal DNA (KanR) was added, incubation continued for 2h at 37°C, and the cells were plated on selective TH plates. Data presented are the means and standard deviations of two independent experiments done in duplicates.

Fig. 4.

Real-Time RT PCR analysis of early and late competence genes in SK36 (A to C) and 133–79 (D to F). Cells were induced with their own or heterologous competence-stimulating peptides. The expression levels are presented relative to the non-induced controls, which were arbitrarily assigned a value of 1. Strains were induced in mid-log phase with 0.5 µg/ml CSP. Data presented are the means and standard deviations of three independent experiments done in duplicate from different days.

Fig. 5. Comparative Real-Time RT PCR analysis of early and late competence genes in SK36 and 133-79. The expression levels of 133-79 are presented relative to that of SK36, which was arbitrarily assigned a value of 1. Data presented are the means and standard deviations of three independent experiments done in duplicate from different days.

Fig. 6. Quantification of biofilms. Biofilm mass of attached cells was quantified with a crystal violet microtiter well assay. Data presented are the means and standard deviations of two independent experiments done in multiple replicates from different days. Asterisks indicate statistical significance ($p = 0.0001$).

Table 1

Real Time PCR (RT) and RACE (RA) primers

Primer	Sequence (5' to 3')
RT 133-79 comC F	TGAAAATCTATTCTTTCAAATTGC
RT 133-79 comC R	CAATCCCATGGATTTGGAAT
RT SK36 comC F	CCATGGATTGGAACACCTC
RT comX R	CCAATCCAACATCATCAAGAGTTC
RT comX F	CCGCCAGCCTTATGAAGAAG
RT 133-79 comD F	GCTATTGAGAGTGCCATTGACAG
RT 133-79 comD R	CTCGGGCTGATAGAGTTCTTCC
RT SK36 comD F	GGAGATTCAGCTTTAAGGAGTGTC
RT SK36 comD R	ACAACCTTGATTGGAAGGCGTTC
RT comYA R	AAGCGGCGTTCATCACCAATTCTC
RT comYA F	TGATTAGGCAGGCTCGGCAAGAAG
RT gyrA F	GCCGTGAGCGAATTGTCGTAA
RT gyrA R	CGAACAGCAGTGATACCGTCAATG
RA IP 133-79	GGATGAACAATTAATCCTATTATTCAA
RA IP SK36	GAATGAATAATTAATCCAATTATTCAA
RA OP 133-79	GAAATGAGATTTCAAATGAAAACACAT
RA OP SK36	GAATTGTGCTAGAAAATGAAAATATGT

Table 2

Comparison of primer efficiencies

	Strain	C _t ratio
C _t <i>gyrA</i> /C _t <i>comC</i>	SK36	0.99 ± 0.04
	133-79	0.97 ± 0.04
C _t <i>gyrA</i> /C _t <i>comD</i>	SK36	1.07 ± 0.01
	133-79	1.07 ± 0.02
C _t <i>gyrA</i> /C _t <i>comYA</i>	SK36	1.05 ± 0.02
	133-79	1.01 ± 0.05
C _t <i>gyrA</i> /C _t <i>comX</i>	SK36	1.13 ± 0.02
	133-79	1.09 ± 0.02