Published in final edited form as:

Neuroreport. 2011 March 9; 22(4): 151–156. doi:10.1097/WNR.0b013e328342ba50.

A basis for the pathological oscillations in basal ganglia: the crucial role of dopamine

Moran Weinberger¹ and Jonathan O Dostrovsky^{1,2}

¹ Department of Physiology, Faculty of Medicine, University of Toronto, Toronto, Canada

² Faculty of Dentistry, University of Toronto, Toronto, Canada

Keywords

basal ganglia; oscillations; Parkinson's disease; subthalamic nucleus; dopamine; beta-activity

Introduction

Parkinson's disease is a progressive neurodegenerative movement disorder characterized by poverty and slowness of movement (akinesia and bradykinesia), muscle rigidity and tremor. The core pathology of the disease is the degeneration of midbrain dopaminergic neurons in the substantia nigra pars compacta that project to the striatum and other basal ganglia nuclei. The striatum is the main input structure of the basal ganglia and receives excitatory projections from many cortical areas and thalamic nuclei. The output neurons of the striatum, named medium spiny neurons, are GABAergic and project to both the globus pallidus internus and externus, and to substantia nigra pars reticulata which are also GABAergic. The other nucleus within the basal ganglia is the subthalamic nucleus which receives direct cortical inputs as well as an input from the globus pallidus externus and its glutamatergic neurons project to the globus pallidus internus and substantia nigra pars reticulata (see Figure 1).

Until recently it was thought that the loss of dopaminergic inputs to the striatum led to hyperactivity of the globus pallidus internus and substantia nigra pars reticulata, and this caused hypokinetic motor symptoms. This model, frequently termed the "rate" model, provided a very plausible explanation for the bradykinesia and akinesia that develops in Parkinson's disease, since the proposed hyperactivity of the GABAergic output nuclei, globus pallidus internus and substantia nigra pars reticulata would lead to hypoactivity in the thalamic relay to motor cortex thus leading to a decreased cortical excitability and slowness and poverty of movements. Initial experimental evidence provided support for this model and was the basis for introduction of pallidotomies and deep brain stimulation in the subthalamic nucleus and globus pallidus internus for treating Parkinson's disease. However, recent findings have cast doubt on the validity of this model and prompted a re-examination of the mechanisms underlying the symptoms of Parkinson's disease [1]. In particular, electrophysiological recordings in cortico-basal ganglia circuits in animal models and Parkinson's disease patients revealed that chronic dopamine depletion is associated with alterations in firing patterns and oscillatory activity and it is now widely believed that these changes play a more important role in mediating the pathophysiology than changes in firing rates.

Many of these advances were facilitated by the opportunity afforded by the therapeutic procedure of implantation of deep brain stimulation electrodes in the subthalamic nucleus and globus pallidus internus which permitted the recordings of neuronal activity and local field potentials in Parkinson's disease patients. Also the use of animal models of Parkinson's disease, in particular monkeys treated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and rats treated with 6-hydroxydopamine, chemicals leading to substantia nigra pars compacta cell death has permitted investigation of the effects of DA depletion on basal ganglia function.

There is growing evidence that increased synchrony between neurons and increased oscillatory firing within the basal ganglia play a key role in the pathophysiology of Parkinson's disease. The origin of these pathological activities, however, is still under extensive investigation and is reviewed below.

Pathological oscillations in Parkinson's disease

The demonstration that loss of dopaminergic input to the striatum leads to excessive synchronized oscillatory firing within the basal ganglia-cortical loops was first reported in the MPTP monkey. Later on, the presence of exaggerated oscillatory activity of single neurons and neuronal populations as reflected in the oscillatory local field potential activity, especially within the subthalamic nucleus and globus pallidus, was seen in recordings from Parkinson's disease patients who were withdrawn from their antiparkinsonian medications [2]. The oscillatory discharges occurred mainly in two frequency bands: the tremor frequency (which is mainly seen in tremulous patients) and the beta frequency (between 11 to 35 Hz). The presence of beta oscillations was also reported in the 6-hydroxydopamine rat model of Parkinson's disease [3].

The link between synchronized oscillations and dopamine depletion is most clearly revealed by the fact that treatment with dopaminergic agents rapidly reduces oscillatory and synchronized firing [2], but the mechanisms by which dopamine exerts these effects are still unclear. Since the basal ganglia-thalamo-cortical loop is to some extent a closed circuit, oscillations are likely to emerge and spread out within this loop. Indeed, there is evidence for increased functional connectivity and coherence between cortex, subthalamic nucleus and pallidum in the dopamine depleted state [3–6]. However, little is known about the cellular and network mechanisms underlying the generation and propagation of these oscillations through the basal ganglia, or why this synchronized oscillatory activity results in bradykinesia.

Basis of neuronal oscillations in the dopamine depleted basal ganglia

Dopamine depletion leads to increased excitability of striatal neurons

Medium spiny neurons normally display a very polarized resting potential (down state) that is interrupted by robust depolarizing events (up states). Down states are determined by inwardly rectifying potassium currents that clamp the membrane potential at a hyperpolarized level, whereas up states are initiated and maintained by concurrent excitatory inputs. Action potential discharges occur only during the up states, when the excitatory inputs are strong enough to overcome the various inhibitory forces generated by the rectifying currents, as well as by feed-forward and feed-back inhibitions from local interneurons and axon collaterals, respectively [7]. Thus, the plateau depolarizations in medium spiny neurons allow "reading" of cortical information by the basal ganglia.

Dopamine plays a key role in modulating the up and down states of medium spiny neurons. Following dopamine depletion, striatal medium spiny neurons exhibit shorter and less

hyperpolarized down states, and fire more frequently in conjunction with cortical inputs [8;9]. After systemic administration of D1 receptor agonists, most medium spiny neurons become silent. In addition to its postsynaptic modulation, dopamine directly inhibits glutamate release from corticostriatal terminals by stimulating D2 receptors located on a subpopulation of cortical afferents [10]. Thus, presynaptic D2 receptors, together with postsynaptic D1 receptors, provide a mechanism for modulating cortical signals. Dopamine loss therefore facilitates the passage and spreading of cortical rhythms throughout the basal ganglia network by disrupting striatal “filtering” of cortical inputs.

Apart from depolarizing medium spiny neurons, dopamine loss has been shown to facilitate electrotonic coupling between neighboring striatal neurons [11]. This effect was hypothesized to result from upregulation of gap junctions, which might further facilitate interactions and synchronization among neurons.

Altered striatal-globus pallidus transmission

Findings in the striatum and globus pallidus in rodents (which is equivalent to the primate globus pallidus externus) support the view that after dopamine loss, an increase in striatal output contributes to the emergence of oscillations in the globus pallidus externus [12]. Moreover, in addition to its effect on the striatum, dopamine acts directly on extra-striatal basal ganglia nuclei. In the globus pallidus, dopamine acts to increase the neuronal activity by attenuating GABAergic inhibition. This is mediated by activation of presynaptic D2 receptors on striatopallidal terminals, which reduces GABA release [13;14], as well as by activation of postsynaptic D4 receptors that inhibit GABA receptor-mediated currents [15]. The loss of GABAergic inhibition in the globus pallidus externus can potentially amplify synchronized oscillations within the subthalamic nucleus-globus pallidus externus network (see below).

In the globus pallidus internus (entopeduncular nucleus in rodents), in contrast, dopamine acts on presynaptic D1-like receptors and enhances GABA release from terminals of the striato-globus pallidus internus projections, therefore reducing neuronal activity in the globus pallidus internus (and the same is true for substantia nigra pars reticulata) [16;17]. Activation of D1-like receptors increases oscillatory and bursting firing in the globus pallidus internus in both normal and MPTP monkeys, whereas blockade of these receptors has the opposite effect [17;18]. This effect is paradoxical since increased bursting and oscillatory activities resemble the pathological state, suggesting that reduced dopamine action in globus pallidus internus does not contribute to the oscillatory firing there following dopamine depletion in animals and Parkinson’s disease patients. Instead, network interactions may be more important in determining these abnormalities which would be consistent with the finding that the transmission of information between the primary motor cortex and the globus pallidus is greatly enhanced after MPTP treatment [6].

Spontaneous activity in subthalamic nucleus neurons: from tonic firing to bursts

The subthalamic nucleus is densely populated by glutamatergic projection neurons which are capable of firing independently of synaptic input due to their pacemaker voltage-gated sodium channels. These neurons are most likely to display tonic firing, and only very few spontaneously fire in bursts [19]. The vast majority of the neurons, however, can switch from a relatively regular firing pattern to an irregular/bursting pattern under various conditions [20–22].

In animal models and Parkinson’s disease, a significantly higher proportion of subthalamic nucleus neurons display burst-like firing patterns, as opposed to regular discharge characteristic of the non-parkinsonian state. D1-like agonists increase the spontaneous and

stimulation-evoked firing rates of tonic subthalamic nucleus neurons, and D2-like agonists influence bursting neurons by reducing their burst duration and turning bursting into a tonic firing pattern [19;23]. Thus, concurrent activation of D1 and D2 receptors promotes tonic firing of subthalamic nucleus neurons. Moreover, by acting on postsynaptic D2-like receptors, dopamine increases the variability of autonomous firing of subthalamic nucleus neurons, and acts to de-correlate subthalamic nucleus activity [24]. Thus in the dopamine-depleted state there is enhancement of the intrinsic membrane properties that enable synchronized burst firing.

In addition to its effect on spontaneous firing patterns, dopamine binding to presynaptic D2 receptors reduces the release probability of glutamate and GABA [25;26] and thus suppresses both glutamatergic excitatory postsynaptic potentials (EPSPs) and GABAergic inhibitory postsynaptic potentials (IPSPs) in subthalamic nucleus [25;27]. Dopamine loss therefore results in enhanced synaptic input to the nucleus, and this contributes even further to the emergence of synchronized oscillations (discussed below).

Subthalamic nucleus-globus pallidus externus network: enhanced feedback inhibition and rebound bursting

The globus pallidus externus is reciprocally connected with the subthalamic nucleus (see Figure 1), creating a network that can potentially generate and maintain low-frequency (~5 Hz) rhythmic activity in the absence of cortical and striatal input [28]. This low-frequency pacemaker activity has the potential to facilitate the faster oscillations that are observed *in vivo* in the parkinsonian state. Extrastriatal dopaminergic modulation is thought to sustain the normal interaction between subthalamic nucleus and globus pallidus externus and its loss may therefore contribute to pathological oscillatory activity.

The autonomous activity in subthalamic nucleus leads to increased numbers of inactivated sodium channels, which can be transiently de-inactivated by the hyperpolarization induced by the pallidal input. Thus, if the inhibitory input is strong enough to completely reset the autonomous, it can promote synchronization because it results in de-inactivation of all of the voltage-gated sodium channels that underlie the autonomous activity [20]. In contrast, weak individual IPSPs promote only partial resetting of subthalamic nucleus neurons, which results in variable delays between spikes and de-synchronization among the neurons. Strong multiple IPSPs do not only reset subthalamic nucleus firing but occasionally promote rebound bursts by bringing the membrane potential to a more hyperpolarized state, which primes low-threshold calcium channels and hyperpolarization-activated cationic currents that are not active during spontaneous activity [20;22].

These various properties suggest that feedback inhibition from the globus pallidus externus has the potential to synchronize the activity of subthalamic nucleus neurons. Because dopamine acts locally to reduce inhibitory synaptic input to the subthalamic nucleus [25;29], its absence may enhance the impact of synchronous GABAergic inputs. In addition, dopamine acts in the globus pallidus both presynaptically via D2/3-like receptors and postsynaptically via D4-like receptors to reduce excitatory subthalamopallidal input [30]. Thus, its absence may facilitate the excitatory feedback from subthalamic nucleus to globus pallidus externus, and thus strengthen the spatiotemporal rhythmic and synchronous oscillatory activity in the subthalamic nucleus-globus pallidus externus network.

After loss of dopamine, pallidal oscillatory firing is influenced more by inhibitory input from the striatum than by excitatory input from the subthalamic nucleus [12]. This is in contrast to the normal situation where the firing of pallidal neurons is entrained by excitatory subthalamic nucleus input [31]. Thus, the level of dopaminergic stimulation is an

important factor in determining the relative influence of the striatum versus the subthalamic nucleus on the globus pallidus externus.

Enhancement of Cortico-subthalamic integration

Since subthalamic nucleus neurons are the only glutamatergic neurons of the basal ganglia, they can function as a direct excitatory relay through which cortical information reaches the output nuclei of the basal ganglia.

As previously mentioned, hyperpolarizing events in subthalamic nucleus neurons can potentially increase their firing by decreasing the number of inactivated voltage-gated sodium channels. Indeed, the coupling between postsynaptic EPSPs and the number of action potentials is dramatically increased when IPSPs precede EPSPs [32]. Thus, enhanced GABAergic inhibition from globus pallidus externus can prime subthalamic nucleus neurons to respond more efficiently to excitatory cortical input by increasing the availability of voltage-gated sodium channels. In Parkinson's disease, the loss of dopamine in the subthalamic nucleus amplifies feedback inhibition from the globus pallidus externus. This increased feedback inhibition, together with the enhanced release probability in subthalamic nucleus synapses, enhances the effectiveness of cortical inputs to generate action potentials by reducing threshold, latency and variability [33]. Furthermore, the synchrony between cortical neurons is enhanced following MPTP treatment [34], thus providing a more powerful oscillatory input to the subthalamic nucleus.

In dopamine-depleted animals the interaction between the subthalamic nucleus and cortical areas is enhanced, and low-frequency oscillations become apparent in the globus pallidus and are correlated with cortical oscillatory activity [3;4]. The oscillatory activity is abolished by ipsilateral cortical ablation, suggesting that abnormal oscillatory activity in the subthalamic nucleus-globus pallidus externus network in the dopamine-depleted state is generated by inappropriate enhanced processing of beta oscillatory inputs originating in the cortex.

Evidence from studies in Parkinson's disease patients

Similar to the observations in animal models, the oscillatory activity seen in Parkinson's disease patients is suppressed by dopamine replacement therapies, in tandem with clinical improvement [2]. Recent evidence has emerged of an interesting correlation between the amount of beta activity in the subthalamic nucleus of Parkinson's disease patients and the improvement in motor symptoms after intake of dopaminergic medication [35–37]. This raises the possibility that the amount of oscillatory activity reflects the sensitivity of the network, or at least the subthalamic nucleus, to dopamine. About a quarter of the neurons in the subthalamic nucleus show oscillatory firing at either beta or tremor frequency [35;38], and some can alter their oscillatory firing from one frequency to another over time [38]. It is therefore tempting to hypothesize that the frequency of oscillations is being determined by an external source of input to the subthalamic nucleus. In fact, the oscillatory firing of subthalamic nucleus neurons is almost always time-locked to the simultaneously recorded local field potentials, which are believed to be generated by the temporal summation of postsynaptic potentials and hence to reflect synchronized inputs [35;39].

The cortex is a likely source of rhythmic oscillatory inputs to the basal ganglia, especially through the subthalamic nucleus. The oscillatory activity (in both the neuronal firing and local field potentials) is maximal in the dorsal part of the subthalamic nucleus [35], which is known to receive input from the primary motor cortex [39]. Further evidence for the cortical origin of oscillations comes from EEG recordings in Parkinson's disease patients who show abnormal beta-band synchronization of cortical networks that is coherent with the oscillatory

activities in subthalamic nucleus and globus pallidus internus [5]. Furthermore transcranial magnetic stimulation of the primary motor and supplementary motor cortical areas significantly suppresses beta activity in the subthalamic nucleus [40], also suggesting that the cerebral cortex drives and/or enhances basal ganglia beta oscillations in Parkinson's disease.

Our recordings in subthalamic nucleus and globus pallidus internus in Parkinson's disease and dystonia patients show that there can be considerable variability in the coupling between local field potentials, which primarily reflect synaptic inputs, and neuronal firing and suggest that at least part of this variability is related to extrastriatal dopamine levels. This situation is also predicted on the basis of the experimental findings regarding dopamine's pre- and post-synaptic actions as discussed above. Thus, in Parkinson's disease patients one generally observes many neurons in subthalamic nucleus and globus pallidus internus with strong oscillatory bursting coherent with the beta frequency oscillatory local field potential activity when off dopaminergic medication but much less when they are on medication. In contrast, in dystonia patients the incidence of globus pallidus internus cells with oscillatory activity coherent with the local field potential is significantly lower than in Parkinson's disease patients (unpublished data). Table 1 summarizes the proposed mechanisms involved in the generation of oscillatory activity in the basal ganglia network following dopamine depletion.

In summary, the data from Parkinson's disease patients add to the growing evidence emerging from animal models and *in vitro* studies that dopaminergic loss in Parkinson's disease may increase the sensitivity of the basal ganglia-thalamo-cortical network to rhythmic oscillatory inputs, perhaps of cortical origin, leading to pathological oscillatory synchronization within and between these structures which interferes with the processing of movement-related signals and results in motor deficits.

Acknowledgments

Supported by: US NIH and Canadian Institutes of Health Research

Reference List

1. Wichmann T, DeLong MR. Deep brain stimulation for neurologic and neuropsychiatric disorders. *Neuron*. 2006; 52:197–204. [PubMed: 17015236]
2. Brown P. Oscillatory nature of human basal ganglia activity: relationship to the pathophysiology of Parkinson's disease. *Movement Disorders*. 2003; 18:357–363. [PubMed: 12671940]
3. Sharott A, Magill PJ, Harnack D, Kupsch A, Meissner W, Brown P. Dopamine depletion increases the power and coherence of beta-oscillations in the cerebral cortex and subthalamic nucleus of the awake rat. *European Journal of Neuroscience*. 2005; 21:1413–1422. [PubMed: 15813951]
4. Magill PJ, Bolam JP, Bevan MD. Dopamine regulates the impact of the cerebral cortex on the subthalamic nucleus-globus pallidus network. *Neuroscience*. 2001; 106:313–330. [PubMed: 11566503]
5. Williams D, Tijssen M, van Bruggen G, Bosch A, Insola A, Di Lazzaro V, et al. Dopamine-dependent changes in the functional connectivity between basal ganglia and cerebral cortex in humans. *Brain*. 2002; 125:1558–1569. [PubMed: 12077005]
6. Rivlin-Etzion M, Marmor O, Saban G, Rosin B, Haber SN, Vaadia E, et al. Low-pass filter properties of basal ganglia cortical muscle loops in the normal and MPTP primate model of parkinsonism. *Journal of Neuroscience*. 2008; 28:633–649. [PubMed: 18199764]
7. Hammond C, Bergman H, Brown P. Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends in Neuroscience*. 2007; 30:357–364.

8. Tseng KY, Kasanetz F, Kargieman L, Riquelme LA, Murer MG. Cortical slow oscillatory activity is reflected in the membrane potential and spike trains of striatal neurons in rats with chronic nigrostriatal lesions. *Journal of Neuroscience*. 2001; 21:6430–6439. [PubMed: 11487667]
9. Murer MG, Tseng KY, Kasanetz F, Belluscio M, Riquelme LA. Brain oscillations, medium spiny neurons, and dopamine. *Cellular and Molecular Neurobiology*. 2002; 22:611–632. [PubMed: 12585682]
10. Bamford NS, Zhang H, Schmitz Y, Wu NP, Cepeda C, Levine MS, et al. Heterosynaptic dopamine neurotransmission selects sets of corticostriatal terminals. *Neuron*. 2004; 42:653–663. [PubMed: 15157425]
11. Onn SP, Grace AA. Alterations in electrophysiological activity and dye coupling of striatal spiny and aspiny neurons in dopamine-denervated rat striatum recorded in vivo. *Synapse*. 1999; 33:1–15. [PubMed: 10380846]
12. Walters JR, Hu D, Itoga CA, Parr-Brownlie LC, Bergstrom DA. Phase relationships support a role for coordinated activity in the indirect pathway in organizing slow oscillations in basal ganglia output after loss of dopamine. *Neuroscience*. 2007; 144:762–776. [PubMed: 17112675]
13. Floran B, Floran L, Sierra A, Aceves J. D2 receptor-mediated inhibition of GABA release by endogenous dopamine in the rat globus pallidus. *Neuroscience Letters*. 1997; 237:1–4. [PubMed: 9406865]
14. Cooper AJ, Stanford IM. Dopamine D2 receptor mediated presynaptic inhibition of striatopallidal GABA(A) IPSCs in vitro. *Neuropharmacology*. 2001; 41:62–71. [PubMed: 11445186]
15. Shin RM, Masuda M, Miura M, Sano H, Shirasawa T, Song WJ, et al. Dopamine D4 receptor-induced postsynaptic inhibition of GABAergic currents in mouse globus pallidus neurons. *Journal of Neuroscience*. 2003; 23:11662–11672. [PubMed: 14684868]
16. Ferre S, O'Connor WT, Svenningsson P, Bjorklund L, Lindberg J, Tinner B, et al. Dopamine D1 receptor-mediated facilitation of GABAergic neurotransmission in the rat strioventrodorsal pathway and its modulation by adenosine A1 receptor-mediated mechanisms. *European Journal of Neuroscience*. 1996; 8:1545–1553. [PubMed: 8758962]
17. Kliem MA, Maidment NT, Ackerson LC, Chen S, Smith Y, Wichmann T. Activation of nigral and pallidal dopamine D1-like receptors modulates basal ganglia outflow in monkeys. *Journal of Neurophysiology*. 2007; 98:1489–1500. [PubMed: 17634344]
18. Kliem MA, Pare JF, Khan ZU, Wichmann T, Smith Y. Ultrastructural localization and function of dopamine D1-like receptors in the substantia nigra pars reticulata and the internal segment of the globus pallidus of parkinsonian monkeys. *European Journal of Neuroscience*. 2010; 31:836–851. [PubMed: 20374284]
19. Baufreton J, Zhu ZT, Garret M, Bioulac B, Johnson SW, Taupignon AI. Dopamine receptors set the pattern of activity generated in subthalamic neurons. *The FASEB Journal*. 2005; 19:1771–1777. [PubMed: 16260646]
20. Bevan MD, Magill PJ, Hallworth NE, Bolam JP, Wilson CJ. Regulation of the timing and pattern of action potential generation in rat subthalamic neurons in vitro by GABA-A IPSPs. *Journal of Neurophysiology*. 2001; 87:1348–1362. [PubMed: 11877509]
21. Beurrier C, Congar P, Bioulac B, Hammond C. Subthalamic nucleus neurons switch from single-spike activity to burst-firing mode. *Journal of Neuroscience*. 1999; 19:599–609. [PubMed: 9880580]
22. Bevan MD, Wilson CJ, Bolam JP, Magill PJ. Equilibrium potential of GABA(A) current and implications for rebound burst firing in rat subthalamic neurons in vitro. *Journal of Neurophysiology*. 2000; 83:3169–3172. [PubMed: 10805713]
23. Baufreton J, Garret M, Rivera A, de la Calle A, Gonon F, Dufy B, et al. D5 (not D1) dopamine receptors potentiate burst-firing in neurons of the subthalamic nucleus by modulating an L-type calcium conductance. *Journal of Neuroscience*. 2003; 23:816–825. [PubMed: 12574410]
24. Ramanathan S, Tkatch T, Atherton JF, Wilson CJ, Bevan MD. D2-like dopamine receptors modulate SKCa channel function in subthalamic nucleus neurons through inhibition of Cav2.2 channels. *Journal of Neurophysiology*. 2008; 99:442–459. [PubMed: 18094105]

25. Shen KZ, Johnson SW. Presynaptic dopamine D2 and muscarine M3 receptors inhibit excitatory and inhibitory transmission to rat subthalamic neurones in vitro. *Journal of Physiology*. 2000; 525:331–341. [PubMed: 10835037]
26. Baufreton J, Bevan MD. D2-like dopamine receptor-mediated modulation of activity-dependent plasticity at GABAergic synapses in the subthalamic nucleus. *Journal of Physiology*. 2008; 586:2121–2142. [PubMed: 18292127]
27. Hassani OK, Feger J. Effects of intrasubthalamic injection of dopamine receptor agonists on subthalamic neurons in normal and 6-hydroxydopamine-lesioned rats: an electrophysiological and c-Fos study. *Neuroscience*. 1999; 92:533–543. [PubMed: 10408602]
28. Pleniz D, Kital ST. A basal ganglia pacemaker formed by the subthalamic nucleus and external globus pallidus. *Nature*. 1999; 400:677–682. [PubMed: 10458164]
29. Zhu Z, Bartol M, Shen K, Johnson SW. Excitatory effects of dopamine on subthalamic nucleus neurons: in vitro study of rats pretreated with 6-hydroxydopamine and levodopa. *Brain Research*. 2002; 945:31–40. [PubMed: 12113949]
30. Hernandez A, Ibanez-Sandoval O, Sierra A, Valdiosera R, Tapia D, Anaya V, et al. Control of the subthalamic innervation of the rat globus pallidus by D2/3 and D4 dopamine receptors. *Journal of Neurophysiology*. 2006; 96:2877–2888. [PubMed: 16899633]
31. Goldberg JA, Kats SS, Jaeger D. Globus pallidus discharge is coincident with striatal activity during global slow wave activity in the rat. *Journal of Neuroscience*. 2003; 23:10058–10063. [PubMed: 14602820]
32. Baufreton J, Atherton JF, Surmeier DJ, Bevan MD. Enhancement of excitatory synaptic integration by GABAergic inhibition in the subthalamic nucleus. *Journal of Neuroscience*. 2005; 25:8505–8517. [PubMed: 16162932]
33. Bevan MD, Hallworth NE, Baufreton J. GABAergic control of the subthalamic nucleus. *Progress in Brain Research*. 2007; 160:173–188. [PubMed: 17499114]
34. Goldberg JA, Boraud T, Maraton S, Haber SN, Vaadia E, Bergman H. Enhanced synchrony among primary motor cortex neurons in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine primate model of Parkinson's disease. *Journal of Neuroscience*. 2002; 22:4639–4653. [PubMed: 12040070]
35. Weinberger M, Mahant N, Hutchison WD, Lozano AM, Moro E, Hodaie M, et al. Beta oscillatory activity in the subthalamic nucleus and its relation to dopaminergic response in Parkinson's disease. *Journal of Neurophysiology*. 2006; 96:3248–3256. [PubMed: 17005611]
36. Ray NJ, Jenkinson N, Wang S, Holland P, Brittain JS, Joint C, et al. Local field potential beta activity in the subthalamic nucleus of patients with Parkinson's disease is associated with improvements in bradykinesia after dopamine and deep brain stimulation. *Experimental Neurology*. 2008; 213:108–113. [PubMed: 18619592]
37. Kuhn AA, Tsui A, Aziz T, Ray N, Brucke C, Kupsch A, et al. Pathological synchronisation in the subthalamic nucleus of patients with Parkinson's disease relates to both bradykinesia and rigidity. *Experimental Neurology*. 2009; 215:380–387. [PubMed: 19070616]
38. Weinberger M, Hutchison WD, Lozano AM, Hodaie M, Dostrovsky JO. Increased gamma oscillatory activity in the subthalamic nucleus during tremor in Parkinson's disease patients. *Journal of Neurophysiology*. 2009; 101:789–802. [PubMed: 19004998]
39. Nambu A, Takada M, Inase M, Tokuno H. Dual somatotopical representations in the primate subthalamic nucleus: evidence for ordered but reversed body-map transformations from the primary motor cortex and the supplementary motor area. *Journal of Neuroscience*. 1996; 16:2671–2683. [PubMed: 8786443]
40. Gaynor LM, Kuhn AA, Dileone M, Litvak V, Eusebio A, Pogosyan A, et al. Suppression of beta oscillations in the subthalamic nucleus following cortical stimulation in humans. *European Journal of Neuroscience*. 2008; 28:1686–1695. [PubMed: 18657185]

Figure 1.

Simplified diagram of basal ganglia connections. Excitatory and inhibitory connections are distinguished by black and gray lines respectively. The overall direct effect of dopamine on synaptic transmission in each nucleus is indicated by the (+) and (-) signs. GPe: globus pallidus externus; GPi: globus pallidus internus; SNc: substantia nigra pars compacta; SNr: substantia nigra pars reticulata; STN: subthalamic nucleus; D1 and D2: dopamine receptor subtypes.

Table 1

Summary of the mechanisms involved in the generation and propagation of oscillatory activity within the basal ganglia following dopamine loss

The pre- and post-synaptic actions of dopamine in the basal ganglia				
Site of Action	Striatum	STN	GPe	GPI
Postsynaptic	Promotes down state of MSNs and filtering of cortical rhythms	Enhances uncorrelated tonic firing	Reduces GABA- mediated and subthalamopallidal currents	
Presynaptic	Inhibits corticostriatal glutamate release	Reduces release of glutamate and GABA	Reduces GABA release at striatal synapses and glutamate release at STN synapses	Enhances GABA release at striatal synapses

Effects of dopamine loss on basal ganglia neurons				
	Striatum	STN	GPe	GPI
	Enhances cortical inputs to the striatum	Enhances intrinsic bursting activity	Enhances responses to striatal inputs	Enables efficient transfer of STN rhythmic inputs
	Disrupts striatal filtering and enhances response to cortical oscillations	Enhances GPe- induced hyperpolarization and rebound STN bursting	Promotes synchronized bursting activity driven by STN	
	Facilitates electrotonic coupling	Enhances direct cortical driving		
Overall effect	Promotes synchronized oscillatory bursting activity in the basal ganglia network			

Abbreviations: MSNs—medium spiny neurons.