Published in final edited form as:

Int J Mass Spectrom. 2011 March 30; 301(1-3): 7–11. doi:10.1016/j.ijms.2010.05.026.

Microwave assisted acid cleavage for denaturation and proteolysis of intact human adenovirus

Catherine Fenselau^{*}, Olli Laine¹, and Stephen Swatkoski²

Department of Chemistry & Biochemistry, University of Maryland, College Park MD 20742 USA

Abstract

Microwave assisted acid cleavage was applied directly to intact adenovirus type 5 to achieve denaturation and proteolysis in a single reaction. The speed of the digestion, coupled with the simplicity of MALDI analysis, allowed peptide products to be profiled in less than 5 min. Identification of peptides from a range of proteins by MALDI-TOF confirms that both denaturation and proteolysis were achieved using low concentrations of acetic acid (12.5%) and short incubations (1.5 to 2 min) at high temperatures (140° C). These conditions favor production of peptides that carry Asp on their C-termini. When this cleavage reaction is carried out in highly enriched H₂ ¹⁸O, a single atom of ¹⁸O is introduced site-specifically into the carboxyl terminal. The labeling reaction is applied to label reporter peptides from human adenovirus type 5 harvested from HeLa cells. Small peptide products of endogenous processing were also observed.

Keywords

adenovirus; Asp selective cleavage; chemical proteolysis; isotope labeling; MALDI-TOF; endogenous processing

1.0 Introduction

1.1

Adenoviruses are commonly used as vectors in genetic engineering and gene therapy [1] and, as such, have generated continuing interest in new methods for their rapid characterization [for example, 2–4]. The adenovirus particle is a large macromolecular assembly ($M_r \sim 2 \times 10^8$ Da) composed of a double-stranded DNA genome and multiple copies of at least 13 distinct proteins [3,5–11]. A viral protease is responsible for processing [12–14]. The small proteome of the adenovirus served earlier as a model system for the development in this laboratory of enzyme-catalyzed ¹⁸O-labeling [15]. In the present report adenovirus type 5 is revisited to demonstrate the capability of Asp-selective rapid microwave-assisted acid cleavage [16–22] concurrently to denature a stable protein complex and to cleave its proteins in a single reaction.

© 2010 Elsevier B.V. All rights reserved.

Phone 1-301-405-8616; Fax 1-301-405-8615.

¹Present address: Work Environment Development, Finnish Institute of Occupational Health, Topeliuksenkatu 41 a A, 00250 Helsinki, Finland

²Present address: Department of Pharmacology and Molecular Sciences, Johns Hopkins School of Medicine, 725 N. Wolfe St. Baltimore MD 21205, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1.2

Much work in proteomics comprises comparative or quantitative analysis, exploiting the capability of mass spectrometry to measure ratios of stable isotope-labeled peptides. Differential stable isotope labels can be introduced into proteins (for example, ICAT [23] and metabolic labeling [24] methods), or into their proteolytic peptide products (C-terminal- ^{18}O [25] and iTraQ [26] methods). Desirable attributes of labeling reactions include speed, complete labeling, site specific labeling and the absence of contaminating chemical by-products [27]. The feasibility is examined here of incorporating one atom of O-^{18} during Asp-selective microwave-supported acid cleavage in highly enriched $\text{H}_2\ ^{18}\text{O}$.

1.3

Figure 1 illustrates the dominant mechanism by which acid cleavage is thought to occur [28] in a protein, and the pathway by which it is expected that a single atom of ^{18}O would be incorporated into proteolytic products. ^{18}O -labeling was first optimized on myoglobin and ovalbumin, and then extended to label peptides from human adenovirus type 5. The effect of acid type and concentration on the extent of incorporation of the isotope label has been evaluated. The site of incorporation is confirmed using laser induced dissociation.

2.0 Material and methods

2.1

Human adenovirus type 5 (ATCC # VR-5) was cultured in HeLa host cells (ATCC # CCL-2) following a published method [15]. HeLa cells were grown in 150 cm^2 culture flasks to about 90% confluence, and exposed to virus stock for 1.5 hr with slow rocking. The virus stock was removed and 2% FBS cell maintenance media was added. As soon as cytopathic lysis was observed, at 2.0 to 2.5 days, the media was removed, host cells were detached and all cells were collected by centrifugation at 8000g for 20 min. The cells were washed twice with 0.1 M PBS pH 7.1, resuspended in 10 mM Tris-HCl buffer, pH 7.6, and the crude virus sample was subjected to analysis.

2.2

Acid cleavage was carried out in 12.5% aqueous acetic acid, unless otherwise specified, in 200 μl glass sample tubes in a Benchmate microwave system (CEM Corp., Matthews NC). The temperature was maintained at $140^\circ \pm 5^\circ \text{C}$ for 1.5 to 5 min by fluctuation in the microwave power. Peptides were labeled by cleavage in $\text{H}_2\ ^{18}\text{O}$, 97.7 atom % ^{18}O normalized (Isotech Inc., Miamisburg OH). Standard proteins were prepared at a concentration of 1 mg/ml in deionized water.

2.3

MALDI spectra were measured on an Axima CFR Plus time-of-flight mass spectrometer (Shimadzu Biotech, Columbia MD) using a 337 nm nitrogen laser. Spectra of peptide samples were acquired in both linear and reflectron modes as averages of 100–200 profile scans, using α -cyanohydroxycinnamic acid (10 mg/ml in 70%/30% (v/v) acetonitrile/deionized water containing 0.1% trifluoroacetic acid) as the matrix. In laser induced dissociation (LID) experiments spectra were acquired in the reflectron mode as averages of 250–500 profiles.

2.4

Nano-electrospray mass spectra were acquired using an LC Packings Ultimate HPLC system (San Francisco CA) interfaced to a PE Sciex API QStar Pulsar I Q-TOF mass spectrometer (Concord ON). Peptide products were desalted using C_{18} Ziptips (Millipore, Bedford MA)

dried and redissolved in 40 μ L 0.1% formic acid. Ten μ L was loaded onto a 15 cm C₁₈ reverse phase column (I.D. 75 μ m) (LC Packings) and eluted with a flow rate of 0.20 μ L/min under a gradient of 3 to 97 % acetonitrile/0.1 % formic acid (solvent B) in 90 min. Solvent A was 0.1% formic acid.

2.5

Bioinformatic searches were carried out using peptide masses or sequence tags. All peptide [M+H]⁺ masses between 900 and 2500, obtained using MALDI-TOF mass spectrometry, were searched using the RMIDB website [29]. A tolerance of 1.0 Da was allowed and searches were performed against a custom database comprising peptides formed *in silico* by Asp-specific cleavage of all adenovirus protein sequences compiled from Swiss-Prot, Genbank, RefSeq and TrEMBL.

Product ion spectra from laser induced dissociation were searched by MASCOT with mass tolerances of $\pm$ 1.0 Da and $\pm$ 1.5 Da for parent and fragment ions, respectively. Spectra were searched against all NCBI nr entries, using the “formic acid” cleavage option and “other viruses” as the taxonomic search parameter. Nano-electrospray MS/MS product ion spectra were searched with mass tolerances of $\pm$ 0.3 and $\pm$ 1.0 Da for Parent and fragment ions, respectively. These spectra were searched against the Swiss-Prot database using MASCOT’s “other viruses” taxonomic classification and a “formic acid” option modified in house [19] to accommodate cleavage on both sides of Asp.

3.0 Results and Discussion

3.1

Figure 2 presents a MALDI-TOF spectrum of acid cleavage products from the adenovirus sample in the mass range 950 to 2450 (top), and a control spectrum obtained from the adenovirus sample not subjected to acid cleavage (bottom). The top spectrum confirms that many peptides are formed by the acid cleavage reaction. In the bottom spectrum small peptides are detected when no chemical cleavage is employed. Virtually all of the ions detected in the product mixture formed by acid cleavage could be assigned as virus peptides based on their masses (Table 1). Among these, the identities of selected peptides were confirmed by partial sequencing by LID and MS/MS (Table I).

3.2

Table I indicates that 37 peptides are identified in the MALDI spectrum (Figure 2) from 8 proteins on the basis of molecular masses and LID sequence tags. Another 9 peptides are listed from ESI MS/MS experiments in order to strengthen the protein identifications and the argument for denaturation. A ninth protein, the minor core protein (PV) was detected only by ESI. Hexon (PII) and penton (PIII) proteins are well represented by peptides. The hexon protein is the second most abundant (720 copies) in each virus, while the penton protein III has only 60 copies per virus. It is of special interest that peptides are identified from the fiber protein (PIV), which participates in attachment to host cells [30]. It is readily lost from viral particles and is often not encountered in protein analysis [5]. It may be retained in the samples studied here because they have not been subjected to final cleanup by gradient centrifugation. The late 100 kDa protein is expressed later in the cycle of infection. In cases where LID and CID experiments are reported, Asp is confirmed to be located at the carboxy terminal of peptide products as previously reported [16–22]. Obtaining high protein sequence coverage was not an objective in this experiment, however it is anticipated that additional peptides would be characterized by repeated injections in HPLC-MS/MS experiments.

3.3

Digestion of the intact virus with immobilized trypsin on the MALDI sample plate produced few peptides (data not shown) because the complex was not denatured. Tryptic digestion of adenovirus proteins is typically preceded by separate denaturation with detergents, reduction and alkylation steps [e.g.,3,5–7] and takes much longer than the 5 min required for microwave assisted acid cleavage and MALDI-TOF profiling.

3.4

The smaller of the two endogenous peptides (m/z 1197.5) detected in Figure 2 (bottom) was identified by LID sequencing as GLRFPSKMFGG. This sequence is part of the major core protein VII [13,14], and the peptide has also been detected by others using HPLC-based analyses [5,6]. Cleavage of this small peptide by the adenovirus protease is reported to be required before incorporation of the major capsid protein VII into the mature virion [14]. The other peptide detected in the control spectrum, at m/z 1350.6, may also be a product of endogenous cleavage. The human adenovirus minor core protein VI is known to undergo processing by adenovirus protease at position 239 [13,14] to produce the peptide GVQSLKRRRCF, which has a calculated average $[MH]^+$ mass of 1350.6. This sequence is also supported by an LID experiment. Adenovirus protease activity is required for the production of infectious virus particles and the ability to detect these peptide products rapidly may be useful to researchers evaluating protease inhibitors to prevent viral infections. In our studies the two peptides were not detectable in samples harvested at 2 and 4 hr post-infection. They were clearly detected in samples harvested at 24 hr.

3.5

Using myoglobin and ovalbumin as model proteins, formic acid (6.0 % and 12.5 %) and acetic acid (6.0 % and 12.5 %) were evaluated for efficient incorporation of the isotope label from $H_2^{18}O$. These concentrations were already known to provide extensive cleavage (16–19). Reaction times of 15 sec, 30 sec, 90 sec, 3 min, 5 min, 15 min and 30 min were evaluated at the constant temperature of 140°C. Non-specific or side-chain incorporation of isotopes was observed when formic acid was used, and at extended reaction times in acetic acid. Based on the extent of proteolysis and high incorporation of a single isotope, 12.5 % acetic acid and 90 sec were chosen as standard labeling conditions for the rest of the study. Partial mass spectra are shown in Figure 3 of a peptide, STRTQINKVVRFD, produced from ovalbumin by cleavage at those conditions in $H_2^{16}O$ (top) and in $H_2^{18}O$ (bottom). A single atom of ^{18}O is found to be incorporated >90% in the latter digestion.

3.6

In order to confirm that the label is incorporated at the carboxy-terminal (see mechanism in Figure 1), tandem mass spectrometry experiments were performed. Figure 4 shows a fragment ion spectrum of a peptide AQQAMTKALELFRND from myoglobin cleaved in heavy water. The label is shown to be localized on the carboxy-terminal Asp moiety, based on the 2 Da mass shift of the y ions observed, and the absence of any mass increase in b ions formed. The original protein carboxyl terminal of myoglobin, observed in the peptide (D)IAAKYKELGRQG, remained unlabeled, and no evidence of random labeling was observed among other myoglobin peptides examined under these conditions.

Peptide products that retain Asp at the C-terminus are preferentially formed at digestion times <2 min [16–22]. Depending on the incubation time, some of these peptides undergo a second hydrolysis to lose C-terminal Asp. Loss of C-terminal Asp in this step was observed to proceed as a second reaction with nonspecific incorporation of either ^{18}O or ^{16}O , which originated within the peptide.

3.7

This rapid cleavage and isotope labeling technique was used to follow the growth of human adenovirus type 5 in HeLa cells. Figure 5 shows isotope envelopes for the peptide PEYWNFRNGD from the adenovirus fiber protein. Adenovirus was harvested from four flasks 24 hour after infection and digested in 12.5 % acetic acid in $H_2^{16}O$ for 90 sec. The virus was harvested from four culture flasks at 48 hr and digested in $H_2^{18}O$. Aliquots of peptide products from the two samples were combined 1:1 by volume to produce the mass spectrum shown. After subtraction of contributions by ^{13}C isotope species, the $^{16}O/^{18}O$ ratio indicates the presence of approximately twice as much virus in the 48 hr incubation compared to the 24 hr incubation, and indicates that the growth rate has not yet begun to level off.

4.0 Conclusions

In some circumstances it may be advantageous to characterize viruses rapidly and with a minimum of processing. In the present paper the capability of microwave assisted acid cleavage is demonstrated to denature human adenovirus and simultaneously to produce peptides without the addition of buffers and chaotropic agents. This proteolytic reaction can also incorporate a single atom of ^{18}O quantitatively and site-specifically into peptides that end in Asp. It is compatible with various automated proteomic workflows [17,19,29].

Acknowledgments

The research was supported in part by a grant from the NIH, GM 021248. Dr. Laine was supported by a fellowship from the Helsingin Sanomat 100th Anniversary Foundation. We thank Prof. Nathan Edwards for advice on the use of RMIDB.

References

1. Walther W, Stein U. Viral vectors for gene transfer: a review of their use in the treatment of human diseases. *Drugs*. 2000; 60:249–271. [PubMed: 10983732]
2. Lehmberg E, Traina JA, Chakel JA, Chang RJ, Parkman M, McCaman MT, Murakami PK, Lahidji V, Nelson JW, Hancock WS, Nestaas E, Pungor EJ. Reversed-phase high-performance liquid chromatographic assay for the adenovirus type 5 proteome. *J.Chomatogr. B*. 1999; 732:411–423.
3. Liu Y, Vellekakmp G, Chen G, Mirza UA, Wylie D, Twarowska B, Tang JT, Porter FW, Wang S, Nagabhusan TL, Pramanik BN. Proteomic study of recombinant adenovirus 5 encoding human p53 by matrix-assisted laser desorption/ionization mass spectrometry in combination with database search. *Int. J.Mass Spectrom.* 2003; 226:55–69.
4. Blyn LB, Hall TA, Libby B, Ranken R, Sampath R, Rudnick K, Moradi E, Desai A, Metzgar D, Russell KL, Freed NE, Balansay M, Broderick MP, Osuna MA, Hofstadler SA, Ecker DJ. Rapid detection and molecular serotyping of adenovirus by use of PCR followed by electrospray ionization mass spectrometry. *J.Clin. Microbiol.* 2008; 46:644–651. [PubMed: 18094138]
5. Blanche F, Monegier B, Faucher D, Duchesne M, Audhuy F, Barbot A, Bouvier S, Daude G, Dubois H, Guillemain T, Maton L. Polypeptide composition of an adenovirus type 5 used in cancer gene therapy. *J. Chromatog. A*. 2001; 921:39–48. [PubMed: 11461012]
6. Roitsch C, Achstetter T, Benchaibi M, Bonfils E, Cauet G, Gloeckler R, L'hote H, Keppi E, Nguyen M, Spehner D, Van Dorsselaer A, Malatme D. Characterization and quality control of recombinant adenovirus vectors for gene therapy. *J.Chromatog. B*. 2001; 752:263–280.
7. Chelius D, Huhmer AR, Shieh CH, Hehmberg E, Traina JA, Slattery TK, Pungor E. Analysis of the adenovirus type 5 proteome by liquid chromatography and tandem mass spectrometry methods. *J.Prot.Res.* 2002; 1:501–513.
8. Chroboczek J, Bieber F, Jacrot B. The sequence of the genome of adenovirus type 5 and its comparison with the genome of adenovirus type 2. *Virol.* 1992; 186:280–285.

9. Shenk, T. Adenoviridae: the viruses and their replication. In: Fields, BN.; Knipe, DM.; Howley, PM.; Chanock, RM.; Melnick, JL.; Monath, TP.; Roizman, B.; Straus, SE., editors. Fundamental virology. 3rd edn.. Philadelphia: Lippincott-Raven; 1996. p. 2111-2148.
10. Stewart PL, Burnett RM. Adenovirus structure by X-ray crystallography and electron microscopy. Curr. Top. Microbiol Immunol. 1995; 199:25–38. [PubMed: 7555058]
11. van Oostrum J, Burnett RM. Molecular Composition of the adenovirus type 2 virion. J. Virol. 1985; 56:439–448. [PubMed: 4057357]
12. Anderson CW. The proteinase polypeptide of adenovirus type 2 virions. Virol. 1990; 177:259–272.
13. Webster A, Leith IR, Hay RT. Activation of adenovirus-coded protease and processing of preterminal protein. J. Virology. 1994; 68:7292–7300. [PubMed: 7933113]
14. Diouri M, Keyvani-Amineh H, Geoghegan KF, Weber JM. Cleavage efficiency by adenovirus protease is site-dependent. J. Biol. Chem. 1996; 271:32511–32514.
15. Yao X, Freas A, Ramirez J, Demirev PA, Fenselau C. Proteolytic ¹⁸O labeling for comparative proteomics: model studies with two serotypes of adenovirus. Anal. Chem. 2001; 73:2836–2842. [PubMed: 11467524]
16. Swatkoski S, Russell S, Edwards N, Fenselau C. Rapid chemical digestion of small acid-soluble spore proteins for Rapid Analysis of *Bacillus* Spores. Anal. Chem. 2006; 78:181–188. [PubMed: 16383326]
17. Swatkoski S, Gutierrez P, Ginter J, Petrov A, Dinman JD, Edwards N, Fenselau C. Integration of residue-specific acid cleavage into proteomic workflows. J. Proteome Res. 2007; 6:4525–4527. [PubMed: 17902642]
18. Li J, Shefcheck K, Callahan J, Fenselau C. Extension of microwave-accelerated residue-specific acid cleavage to proteins with carbohydrate side chains and disulfide linkages. Internat. J. Mass Spectrom. 2008; 278:109–113.
19. Swatkoski S, Gutierrez P, Wynne C, Petrov A, Dinman JD, Edwards N, Fenselau C. Evaluation of microwave-accelerated residue-specific acid cleavage for proteomic applications. J. Proteome Res. 2008; 7:579–586. [PubMed: 18189344]
20. Hauser NJ. Online Microwave D-cleavage LC-ESI-MS/MS of intact proteins: Site - specific cleavages at aspartic acid residues and disulfide bonds. J. Proteome Res. 2008; 7:1012–1026. [PubMed: 18198820]
21. Hauser NJ, Han HL, McLuckey SA, Basile F. Electron transfer dissociation of peptides generated by microwave D-cleavage digestion of proteins. J. Prot. Res. 2008; 7:1867–1872.
22. Hua L, Low TY, Sze SK. Microwave-assisted specific chemical digestion for rapid protein identification. Proteomics. 2006; 6:586–591. [PubMed: 16342144]
23. Tao WA, Aebersold R. Advances in quantitative proteomics via stable isotope tagging and mass spectrometry. Curr. Opin. Biotechnol. 2003; 14:110–118. [PubMed: 12566010]
24. Ong SE, Kratchmarova I, Mann M. Properties of ¹³C-substituted arginine in stable isotope labeling by amino acids in cell culture. J. Proteome Res. 2003; 2:173–181. [PubMed: 12716131]
25. Yao X, Afonso C, Fenselau C. Dissection of Proteolytic ¹⁸O Labeling: Endoprotease-Catalyzed ¹⁶O-to-¹⁸O Exchange of Truncated Peptide Substrates. J. Proteome Res. 2003; 2:147–152. [PubMed: 12716128]
26. Ross PI, Huang YN, Marchese JN, Williamson V, Parker K, Hattan S, Khainovski N, Pillai S, Dey S, Daniels S, Purkayastha S, Juhasz P, Martin S, Bartlett-Jones M, He F, Jacobson A, Pappin DJ. Multiplexed Protein Quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents. Molec. Cell. Proteomics. 2004; 3:1154–1169. [PubMed: 15385600]
27. Fenselau C, Yao X. ¹⁸O₂-Labeling in Quantitative proteomic Strategies: A Status Report. J. Proteome Res. 2009; 8:2140–2143. [PubMed: 19338309]
28. Li AQ, Sowder RC, Henderson LE, Moore SP, Garfinkel DJ, Fisher RJ. Chemical Cleavage at Aspartyl Residues for Protein Identification. Anal. Chem. 2001; 73:5395–5402. [PubMed: 11816565]
29. www.RMIDB.org 01-20-10
30. Philipson L. Structure and assembly of adenoviruses, Current Top. Microbiol. Immun. 1983; 109:1–52.

Figure 1. Mechanistic scheme for acid catalyzed cleavage at aspartic acid in a protein (28). Bolded **O** predicts incorporation of labeled oxygen atoms.

Figure 2.
MALDI-TOF mass spectra of peptides from adenovirus sample harvested at 48 hr. Top: 2 min acid digestion; bottom: no digestion

Figure 3. Partial MALDI-TOF mass spectra of the peptide STRTQINKVVRFD produced by acid digestion of ovalbumin in $H_2^{16}O$ (top) and $H_2^{18}O$ (bottom).

Figure 4. Laser induced dissociation spectrum of the peptide AQGAMTKALELFRND produced by acid digestion of myoglobin. Inset is a list of predicted b and y ions.

Figure 5.

Partial MALDI-TOF mass spectrum of the peptide PEYWNFRNGD in solutions of peptide products combined 1:1 by volume from labeled adenovirus (48 hr harvest) and unlabeled adenovirus (24 hr harvest). Ion intensities are shown in parentheses.

TABLE 1

Characterization of Peptides Detected by MALDI TOF MS

Protein	Observed [M+H] ⁺	Calculated [M+H] ⁺	Peptide	Confirmed by MS/MS	MASCOT Expect Value
Hexon (PII)	1183.3	1183.4	D.SITQKKFLCD.R		
	1298.0	1298.5	V.DSITQKKFLCD.R		
	1350.7	1350.6	D.RNTELSYQLLL.D		
	1423.7*	1423.7	D.PYYTYSGSIPYL.D	ESI CID	0.036
	1462.3	1462.7	D.YMNKRVVAPGLVD.C		
	1494.4	1493.6	G.DNLTPKVVLVLYSED.V		
	1530.7	1530.7	D.GEGYNVAQCNMTKD.W		
	1538.8*	1538.7	D.PYYTYSGSIPYLD.G	ESI CID	1.7e-5
	1602.1	1601.9	D.RSQRLTLRFIPVD.R		
	1674.3	1674.9	D.RTRYFSMWNQAVD.S		
	1760.4	1761.0	D.VNMVLQSSGLGNDLRVD.G		
	1818.2	1818.1	D.RLLTPNEFEIKRSVD.G		
	1886.8	1887.2	D.RSQRLTLRFIPVDRE.D		
	2002.8	2002.3	D.RSQRLTLRFIPVDRED.T		
	2002.8	2003.3	D.GTFYLNHTFKKVAITFD.S	ESI CID	0.053
	2040.7	2040.2	D.RTRYFSMWNQAVDSYD.P	ESI CID	0.011
	2060.8	2061.3	L.DSIGDRTRYFSMWKQAVD.S		
Penton (PIII)	2167.9	2167.4	D.NPNTYDYMNKRVVAPGLVD.C	ESI CID	0.0034
	2425.2	2424.7	D.RMYSFFRNFPMSRQVVD.D		
	994.8	995.1	D.VDAYQASLK.D		
	1011.0	1011.1	V.DAYQASLK.D		
	1011.0	1011.2	D.TRNFRLGFD		
	1126.3	1126.3	D.TRNFRLGFD.P		
	1573.5	1573.8	D.PQTGIRSWTLLCTP.D		
	1689.2	1688.9	D.PQTGIRSWTLLCTPD.V	ESI CID	0.0057
	2044.1*	2044.1	D.PVTGLVMFPGVYTNEAFHPD.I	ESI CID	1.5e-6
	2135.0*	2134.9	D.STFTQYRSWYLAYNYGD.P	ESI CID	0.016
IVa2 (PIVa2)	1016.9	1017.1	D.LILEHNYD.V		

Protein	Observed [M+H] ⁺	Calculated [M+H] ⁺	Peptide	Confirmed by MS/MS	MASCOT Expect Value
Fiber (P1V)	1462.3	1461.6	D.RDAVEQVTELWD.R		
	1658.2	1659.0	D.RLELLGQTLKSMPTA.D		
	2425.2	2425.9	D.LVRENMRVRDMLNEVAPLLR.D		
	2442.4	2442.8	Y.DVSDPRNIFAQAARGPIAIMD.E		
	1183.3	1183.3	D.PEYWNFRNG.D	MALDI LID	0.023
Hexon-associated (PIX)	1298.0	1298.3	D.PEYWNFRNGD.L	ESI CID	9.7e-5
	1412.8	1413.4	L.DPEYWNFRNGD.L		
	1674.3	1673.9	D.SQGNMQLNVAGGLRID.S		
	1767.3	1767.0	D.LRQQVSALKASSPPNA V.-	ESI CID	2.8e-5
	1498.8 [*]	1498.7	D.SLTAPSEFATTASTD.A	ESI CID	1.9e-5
Late 100 kDa.	1802.8	1802.0	D.LWTAFNERSVAAHLAD.I		
	1858.1 [*]	1858.0	D.IASLNEVPKIFEGGLGRD.E	ESI CID	0.00048
	2709.5 [*]	2709.3	D.AANAPTTFPVEAPPLEEEVHIEQD.F	ESI CID	2.3e-5
	1350.7	1350.7	GVQSLKRRRCF (Endogenous)	MALDI LID	
	1655.0 [*]	1655.9	D.LANQAVQNKINSKLD.P	ESI CID	0.0037
Minor Capsid (PVI)	1197.5	1197.5	GLRFP SKMFGG (Endogenous)	MALDI LID	
Major Core (PVII)	2442.4	2442.7	D.AIDAVVEEARNYTPPPPVSTVD.A		
Minor Core (PV)	1919.1 [*]	1919.0	D.AAVQAVAAAASKTSTEVQTD.P	ESI CID	2.7e-5

^{*} indicates that peptide was only found in ESI mass spectra. D is shown at the carboxy terminus.