Published in final edited form as:

Tetrahedron Lett. 2011 April 27; 52(17): 2072–2075. doi:10.1016/j.tetlet.2010.10.111.

Stereoselective Synthesis of the Decahydrofluorene Core of the Hirsutellones[#]

Geoff T. Halvorsen^a and William R. Roush^{a,*}

^a Department of Chemistry, Scripps Florida, 130 Scripps Way, Jupiter, FL 33458, USA

Abstract

A stereoselective synthesis of the decahydrofluorene core of the hirsutellones was accomplished in 8 steps and in 43% overall yield. The key step of the synthesis is the highly stereoselective intramolecular Diels-Alder cyclization of the siloxacyclopentene constrained tetraene **1**.

Keywords

hirsutellone B; Intramolecular Diels-Alder reaction; Siloxacyclopentene-constrained dienophile; stereoselective synthesis; Oxonium ion activated dienophile

The hirsutellones are a family of polyketides isolated from the insect pathogenic fungus *Hirsutella nivea* BCC 2594 that display potent antitubercular activity, with minimum inhibitory concentrations against *Mycobacterium tuberculosis* H₃₇R ranging from 0.78 to 3.125 µg/mL.^{1,2} The interesting polycyclic structures consisting of a decahydrofluorene core with a 12- or 13-membered *para*-cyclophane, as well as the antitubercular activity make the hirsutellones intriguing synthetic targets (Figure 1).³

We reasoned that the tricyclic core common to the hirsutellones could be assembled by an intramolecular Diels-Alder (IMDA) cycloaddition⁴ of tetraene **1**, utilizing a siloxacyclopentene-constrained dienophile to ensure proper stereochemical control of the *trans*-fused C(14) ring juncture relative to the C(13)-hydroxyl group in **2** (Figure 2).^{5,6} Control of the hydroxyl stereochemistry at this position is often difficult to achieve in IMDA reactions.^{4,7} Thus, implementation of this strategy would enable us to control all of the stereocenters present in the decahydrofluorene core while simultaneously introducing functionality at C(13) and C(15) that could be used to append the macrocycle en route to the natural products.

The synthesis of the common hirsutellone core **2** commenced with the synthesis of triene aldehyde **9** as presented in Scheme 1. The enantioselective Diels-Alder reaction of commercially available (*E,E*)-hexa-2,4-dien-1-ol (**3**) with methyl acrylate was performed in the presence of the reagent generated from (*R*)-1,1'-bi-2-naphthol ((*R*)-BINOL), Me₂Zn and MeMgBr as described by Ward.⁸ This reaction provided lactone **4** in 95% yield and with

[#]This manuscript is dedicated to Professor Harry H. Wasserman in celebration of his 90th birthday, and in recognition of his fundamental contributions to organic chemistry and his extraordinary service and leadership on behalf of the community of organic chemists.

*Corresponding author. Tel.: 561-228-2450; fax: 561-228-3052; roush@scripps.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

96% e.e. Hydrogenation of **4** over 10% Pd/C gave the saturated lactone **5** (99% yield), which was then reduced to lactol **6** using diisobutylaluminum hydride (DIBAL) in CH₂Cl₂ at -78 °C. Wittig olefination of lactol **6** using the ylide generated by treatment of trimethylphosphonium salt **7**⁹ with potassium hexamethyldisilazide in THF at 0 °C then gave trienol **8a** with no detectable (*Z*)-olefin at the newly formed double bond, but with approximately 15% of the *E,Z* isomer **8b** which could not be separated from **8a**.^{9,10} Trienol **8a** was then oxidized under Parikh-Doering conditions,¹¹ followed by treatment with potassium carbonate in methanol, which led to smooth epimerization of the aldehyde and provided exclusively **9** with the desired equatorial carboxaldehyde stereochemistry.

Aldehyde **9** was further elaborated to triene **1** as summarized in Scheme 2. Addition of the lithium acetylide generated by treatment of propiolaldehyde acetal **10**¹² with *n*-BuLi to a solution of **9** in 1:1 THF and *tert*-butyl methyl ether (TBME) at 0 °C, with warming to 60 °C, provided propargyl alcohol **11** in 60% yield with excellent Felkin-Anh diastereoselectivity (>95 : 5 d.r.). Hydrosilylation of propargyl alcohol **11** was accomplished using the procedure developed by Lee.¹³ Thus, treatment of **11** with tetramethyldisilazane to give the silyl ether followed by addition of potassium *tert*-butoxide in THF gave siloxacyclopentene **1** in 84% yield. Because we had anticipated at the outset that the diastereoselectivity of the key IMDA reaction might be poor if the siloxacyclopentene strategy was not utilized,^{4,7} we also synthesized the non-constrained triene **13** for use in comparative IMDA reactions. Therefore, hydroalumination of propargyl alcohol **11** with sodium bis-(2-methoxyethoxy)-aluminumhydride (RED-Al®) followed by standard workup gave allylic alcohol **12** (84%) which was then protected as the triethylsilyl ether triene **13** (Scheme 2).

Cycloaddition reactions of **1** and **13** were performed by treatment of a solution of the triene in CH₂Cl₂ at -78 °C with trimethylsilyltriflate followed by warming to -20 °C.¹⁴ These reactions undoubtedly proceed via oxonium ion activated dienophiles, generated by TMS-OTf promoted ring opening of the cyclic acetals.¹⁴ After the cycloadditions were complete, the crude reaction mixtures were immediately subjected to desilylation conditions (TBAF in THF) to facilitate product isolation and purification.^{15,16}

The cycloaddition of **1** and subsequent proteodesilylation of the crude cycloadduct gave decahydrofluorene **2** in excellent yield and as a single diastereomer as determined by ¹H NMR analysis of the crude reaction mixture. The stereochemistry of **2** was assigned by using a combination of coupling constant analysis and NOE studies, as summarized in Scheme 3.

In contrast, the Lewis acid promoted cyclization of the unconstrained tetraene **13** was less selective, giving an inseparable 85:15 mixture of cycloadduct diastereomers, among which **2** predominated, following deprotection of the silyl ether (Scheme 3). Analysis of the key coupling constants identified for **14** enabled us to identify the minor diastereomer in this reaction as decahydrofluorene **14**, which is epimeric to **2** at the vinyl substituent (in particular *J*_{f,g} = 11.8 Hz in **14**, vs. 6 Hz in **2**; see Scheme 3). This compound presumably arises from the IMDA cyclization of the minor (*E,Z*)-tetraene present in triene **13** (see Scheme 4). This raises the question, why was the analogous cycloadduct not realized in the IMDA cyclization of (*Z*)-**1** (triene **1** used in Scheme 3 was also an 85 : 15 mixture of *E* and *Z* olefin isomers, deriving from **7**).⁹

Examination of transition states available for the IMDA reactions of **1** and **13** are presented in Schemes 4 and 5. Examination of **TS13A** and **TS13B** in Scheme 5 for the IMDA cyclization of the (*E*)- and (*Z*)-isomers of **13** indicate that the (*Z*)- isomer can readily cyclize via transition state **TS 13B**. In contrast, **TS1B** in Scheme 5 for the cyclization of the (*Z*)-isomer present in **1** suffers from serious non-bonded interactions between the (*Z*)-vinyl

group and the Me₂Si— unit of the siloxacyclopentene constrained dienophile. Therefore, the IMDA cyclization of **1** gives a single cycloadduct, **2**, whereas the IMDA cyclization of **13** gives a mixture of **2** and **14** corresponding to the mixture of (E)- and (Z)-triene isomers present in the cyclization substrate. However, it remains unclear why the IMDA reaction of **13** does not provide a third cycloadduct, **15**, deriving from the alternative transition state **TS13C**. Transition states analogous to **TS 13C** are sometimes competitive in IMDA reactions of trienes with alkoxy substituents at the position allylic to the dienophile⁷ (but is physically inaccessible in the IMDA cyclization of **1**, owing to the siloxacyclopentene unit in the dienophile).

In summary, we have completed an 8-step asymmetric synthesis of the decahydrofluorene core unit common to the hirsutellone family of natural products that proceeds in 43% overall yield from the commercially available diene **3**. The key step of this synthesis is the highly diastereoselective cyclization of siloxacyclopentene-constrained triene **1**. The diastereoselectivity of this cyclization proved superior to that of the unconstrained triene **13**, which gave an 85 : 15 mixture of diastereomeric cycloadducts. Further efforts towards the completion of syntheses of hirsutellone family members will be reported in due course.

Acknowledgments

Financial support provided by the National Institutes of Health (GM026782) is gratefully acknowledged.

References and notes

1. Isaka M, Rugseree N, Maithip P, Kongsaree P, Prabpia S, Thebtaranonth Y. *Tetrahedron*. 2005; 61:5577.
2. Isaka M, Prathumpai W, Wongsa P, Tanticharoen M. *Org Lett*. 2006; 8:2815. [PubMed: 16774264]
3. (a) Nicolaou KC, Sarlah D, Wu TR, Zhan W. *Angew Chem Int Ed*. 2009; 48:6870. (b) Tilley SD, Reber KP, Sorensen EJ. *Org Lett*. 2009; 11:701. [PubMed: 19119873] (c) Huang M, Huang C, Liu B. *Tetrahedron Lett*. 2009; 50:2797.
4. Reviews of intramolecular Diels-Alder reactions: (a) Ciganek B. *Org React*. 1983; 32:1. (b) Craig D. *Chem Soc Rev*. 1986; 16:187. (c) Roush, WR. *Comprehensive Organic Synthesis*. Trost, BM., editor. Vol. 5. Pergamon Press; Oxford: 1991. p. 513-550. (d) Takao K, Munakata R, Tadano K. *Chem Rev*. 2005; 105:4779. [PubMed: 16351062] (e) Juhl M, Tanner D. *Chem Soc Rev*. 2009; 38:2983. [PubMed: 19847335]
5. Halvorsen GT, Roush WR. *Org Lett*. 2007; 9:2243. [PubMed: 17480094]
6. Halvorsen GT, Roush WR. *Org Lett*. 2008; 10:5513. [PubMed: 19053726]
7. (a) Roush WR, Peseckis SM. *J Am Chem Soc*. 1981; 103:6696. (b) Funk RL, Zeller WE. *J Org Chem*. 1982; 47:180. (c) Suzuki T, Tanaka N, Matsumura T, Hosoya Y, Nakada M. *Tetrahedron Lett*. 2006; 47:1593.
8. Ward DE, Souweha MS. *Org Lett*. 2005; 7:3533. [PubMed: 16048335]
9. The phosphonium salt **7** was generated from 2,4-pentadienyl iodide and Me₃P, and contains c.a. 15% of the (Z)-isomer which we were unable to separate. Therefore, the 85 : 15 mixture of isomeric phosphonium salts was used in the Wittig reaction. The corresponding mixture of triene isomers produced at the stage of **8** could not be separated at any stage of this synthesis.
10. Vedejs E, Marth CF. *J Am Chem Soc*. 1988; 110:3948.
11. (a) Parikh JR, Doering WvE. *J Am Chem Soc*. 1967; 89:5505. (b) Tidwell TT. *Org React*. 1990; 39:297.
12. Chemler SR, Roush WR. *J Org Chem*. 2003; 68:1319. [PubMed: 12585871]
13. Maifeld SV, Lee D. *Org Lett*. 2005; 7:4995. [PubMed: 16235941]
14. Gassman PG, Singleton DA, Wilwerding JJ, Chavan SP. *J Am Chem Soc*. 1987; 109:2182.
15. The siloxacyclopentane unit of the cycloadduct derived from **1** was unstable and could not be isolated.

16. (a) Stork G, Mah R. *Tetrahedron Lett.* 1989; 30:3609. (b) Heitzman CL, Lambert WT, Mertz E, Shotwell JB, Tinsley JM, Va P, Roush WR. *Org Lett.* 2004; 7:2405. [PubMed: 15932209]

Figure 1.
Structures of hirsutellones A-C

Figure 2.
IMDA cycloaddition strategy for synthesis of the hirsutellone core.

Scheme 1.
Synthesis of triene aldehyde **9**

Scheme 2.
Synthesis of tetraenes **1** and **13**

Scheme 3.
Intramolecular cycloadditions of **1** and **13**

Scheme 4.
Transition state analysis for IMDA cyclization of **1**

Scheme 5.
Transition state analysis for IMDA cyclization of **13**