Published in final edited form as:

Clin Immunol. 2011 May ; 139(2): 185–192. doi:10.1016/j.clim.2011.02.003.

Antigen-specific IgA B memory cell responses to *Shigella* antigens elicited in volunteers immunized with live attenuated *Shigella flexneri* 2a oral vaccine candidates

J. K. Simon^{a,b,c}, M. Maciel Jr.^{a,d}, E.D. Weld^{a,b}, R. Wahid^a, M.F. Pasetti^a, W.L. Picking^e, K. L. Kotloff^{a,b}, M. M. Levine^{a,b}, and M. B. Sztein^{a,b,*}

^aDivision of Infectious Diseases and Tropical Pediatrics, Department of Pediatrics, University of Maryland School of Medicine, Baltimore, Maryland, USA

^bDivision of Geographic Medicine, Department of Medicine, Center for Vaccine Development, University of Maryland School of Medicine, Baltimore, Maryland, USA

^eDepartment of Microbiology and Molecular Genetics, Oklahoma State University, Stillwater, Oklahoma, USA

Abstract

We studied the induction of antigen-specific IgA memory B cells (B_M) in volunteers who received live attenuated *Shigella flexneri* 2a vaccines. Subjects ingested a single oral dose of 10⁷, 10⁸ or 10⁹ CFU of *S. flexneri* 2a with deletions in *guaBA* (CVD 1204) or in *guaBA*, *set* and *sen* (CVD 1208). Antigen-specific serum and stool antibody responses to LPS and Ipa B were measured on days 0, 7, 14, 28 and 42. IgA B_M cells specific to LPS, Ipa B and total IgA were assessed on days 0 and 28. We show the induction of significant LPS-specific IgA B_M cells in anti-LPS IgA seroresponders. Positive correlations were found between anti-LPS IgA B_M cells and anti-LPS IgA in serum and stool; IgA B_M cell responses to IpaB were also observed. These B_M cell responses are likely play an important role in modulating the magnitude and longevity of the humoral response.

Keywords

Vaccine; *Shigella*; B cell memory; Immunoglobulin A; IgA; mucosal immunity

1. Introduction

Shigella infections continue to be a major cause of morbidity and mortality among children under 5 years old living in the developing world. Every year, there are 165 million cases of shigellosis worldwide and 14,000 cases reported in the United States; it is estimated that because of underreporting, the number of actual cases may be twenty times higher [1,2]. The

© 2010 Elsevier Inc. All rights reserved.

*Corresponding author: Marcelo B. Sztein, M.D., Tel.: +1 410 706 5328; fax: +1 410 706 6205. msztein@medicine.umaryland.edu, 685 W. Baltimore Street, Suite 480, Baltimore MD 21201, United States.

^cPresent address: NanoBio Corporation, Ann Arbor, Michigan, USA

^dPresent address: ETEC Vaccine Program, Naval Medical Research Center, Silver Spring, Maryland, USA

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

increasing prevalence of resistance to multiple antimicrobials is of concern [3] and *Shigella* is considered a Category B bioterror agent by the CDC [4]. *Shigella flexneri* is endemic throughout the developing world, and causes more mortality than any other species of *Shigella* [5]. There is a high demand for a safe and effective oral vaccine, and the WHO has prioritized the development of a well-tolerated vaccine that induces durable immunity against shigellosis [1,6].

By engineering rational deletions in the wild-type *Shigella flexneri* 2a strain 2457T, two vaccine candidates, designated CVD 1204 and CVD 1208, were constructed at the Center for Vaccine Development (CVD). CVD 1204 contains deletions in *guaA* (encoding a guanosine monophosphate synthase) and *guaB* (encoding an inositol monophosphate dehydrogenase), which impair the biosynthesis of guanine nucleotides; CVD 1208 has additional deletions of *set* and *sen* genes that encode *Shigella* enterotoxins 1 and 2, respectively. In a Phase 1 trial CVD 1204 was shown to be clearly attenuated compared to its wild type parent (based on comparison with data from multiple previous challenge studies), while CVD 1208 appeared fully attenuated yet immunogenic [7]. Clinical adverse reactions (diarrhea, dysentery and/or fever) occurred in 8 of 23 recipients of CVD 1204 but in only 1 of 21 recipients of CVD 1208 [7].

Putative correlates of protection against shigellosis reported in the literature include serum IgG antibodies against lipopolysaccharide (LPS) O antigen and serotype specific O antigen peripheral blood IgA antibody secreting cells (ASC) [2,8,9]. Other antibody and cell-mediated immune responses (CMI) against conserved antigens such as invasion plasmid antigens (Ipa) may also play a role in protective immunity [2,10–13]. An optimal vaccine should not only induce enduring systemic and mucosal antibody responses but also allow the host to mount an anamnestic immune response upon subsequent re-exposure to antigen. This response is faster, stronger, and qualitatively better than primary responses and depends on the presence of B_M cells [14]. Following natural *Shigella* infection, as well as after ingestion of some live attenuated *Shigella* vaccines, relatively long-term humoral and secondary secretory IgA immune responses to LPS in stool have been described [15]. We have previously demonstrated the induction of IgG B_M responses by live attenuated *Shigella* vaccines in human volunteers [16]. However, the presence of IgA B_M responses has not been reported. In this study we examined the hypothesis that volunteers who display mucosal and serum antibody responses to CVD 1204 and CVD 1208 live-attenuated oral *Shigella* vaccines also exhibit IgA B_M cell responses specific to LPS, IpaB and other *Shigella* antigens.

2. Materials and methods

2.1. Specimens

46 healthy adult volunteers 18–45 years of age from the Baltimore–Washington area received a single oral dose of *S. flexneri* 2a Δ *guaBA* (CVD 1204) or *S. flexneri* 2a Δ *guaBA* Δ *sen* Δ *set* (CVD 1208) as previously described [7]. Volunteers received 10⁷, 10⁸, or 10⁹ CFU of each vaccine strain or placebo, and sera and stools were collected on days 0, 7, 14, 28, and 42. In addition, peripheral blood mononuclear cells (PBMC) were obtained on days 0 and 28 after oral vaccination. PBMC specimens were cryopreserved and stored in liquid nitrogen until use as previously described [17]. Seroresponse, measured by ELISA [7], was defined as $\geq$ 4-fold rise of antigen-specific IgA antibody in serum (seroresponders) and a $\geq$ 4-fold rise of antigen-specific IgA/total IgA in stool (mucosal responders) after oral vaccination as compared to pre-vaccination. Adequate specimens were available to assay 13 seroresponders and 11 non-seroresponders; these included subjects immunized with placebo or 10⁷, 10⁸, or 10⁹ CFU of the *Shigella* strains. For B_M assays, subjects from all three dosage level cohorts were analyzed. Prior to enrollment, the purpose of the study was

explained to the subjects and they passed a written test containing questions regarding the rationale for the study, risks and procedures. Informed consent was obtained from all participants and the study was approved by the UMD Institutional Review Board.

2.2. Antigen preparation

LPS was purified by the hot aqueous phenol method of Westphal [18]. IpaB, IpaC, IpaD, MxiH, VirG and *Yersinia pestis* LcrV antigens were purified as recombinant proteins from *E. coli*. A PCR fragment of virG encompassing amino acids 68–774 was copied from the vir plasmid of *S. flexneri* 2457T by standard PCR. The fragment was ligated into pET22b (Novagen, Madison, WI) and the ligation product used to transform *E. coli* NovaBlue. The resulting plasmid was sequenced and used in the protein expression system. IpaD, VirG and LcrV were purified by standard His tag chromatography and dialyzed into PBS as described previously [19,20]. IpaC and MxiH were solubilized from inclusion bodies with 6M urea, purified by standard His tag chromatography, and refolded by step dialysis into PBS as previously described [21,22]. IpaB was co-expressed with its cognant chaperone, IpgC, as described previously [23]. The complex was purified by standard His tag chromatography via the His tag fused to the IpgC. IpaB was released from IpgC with 1% *n*-octyl-polyoxyethylene.

2.3 Mucosal and systemic antibodies

Serum antibodies specific for *S. flexneri* LPS and Ipa B were measured by ELISA as we previously described [7]. Briefly, plates were coated with LPS (5 µg/mL) or IpaB (0.1 µg/mL) and blocked with 10% dried milk in PBS. Samples were evaluated in serial 2-fold dilutions. HRP-labeled goat anti-human Fc α chain (ICN) was used as conjugate and TMB microwell Peroxidase (Kirkegaard & Perry Laboratories, KPL) as substrate. Titers were calculated from linear regression curves as the reciprocal serum dilution that produced an OD of 0.2 above the blank (EU/ml). Total and LPS-specific fecal IgA were also measured by ELISA as previously described [7]. Plates were coated with either α-chain specific anti-human IgA (1 µg/mL; Jackson ImmunoResearch Laboratories) or LPS (10 µg/mL). Stool supernatants were tested in serial 2-fold dilutions. HRP-labeled goat anti-human IgA (Jackson) was used as conjugate and TMB (KPL) as substrate. IgA concentrations were calculated by interpolation into a standard curve of human IgA (Calbiochem). Data are reported as the ratio of LPS-specific/ total IgA levels.

2.4 Antibody secreting cells

IgA and IgG ASCs specific for *S. flexneri* LPS and IpaB were detected by ELISpot as we previously described [7,24]. A positive ASC response was defined as a post-vaccination count at least 3 SD above the mean prevaccination count and at least 8 cells/10⁶ PBMC.

2.5. PBMC expansion

Expansion of PBMC to measure B_M cells was performed as described by Crotty *et al.* [25]. Briefly, PBMC specimens were thawed, washed with complete cRPMI 1640 containing 100 IU/mL penicillin, 100 µg/mL streptomycin (CellGro, Manassas, VA), 2 mM L-glutamine (BioWhittaker, Walkersville, MD), and 10% heat-inactivated fetal bovine serum (HyClone, Logan, UT), and expanded for 5 days in 6-well plates (1 × 10⁶ cells/well) in the presence of 1/100,000 pokeweed mitogen (PWM, kindly provided by Dr. S. Crotty), 6 µg/mL CpG-2006 (Qiagen/Operon, Huntsville, AL), 50 uM β-Mercapto-ethanol, and 1/10,000 *Staphylococcus aureus* Cowan (SAC, Sigma-Aldrich, St. Louis, MO) in cRPMI (expansion media) in a total volume of 2 mL/well. After 2 days of incubation, cells were fed by replacing 2 mL of cRPMI in each well.

2.6. B Memory cell assays

96-Well ELISPOT MAHA (Millipore, Billerica, MA) plates were coated with 5 µg/mL LPS, 0.5 µg/mL of IpaB, IpaC, IpaD, VirG, MxiH or *Yersinia pestis* LcrV (as a negative control) or 5 µg/mL of mouse anti-human IgA in PBS and incubated overnight at 4°C. The plates were then blocked with 1% bovine serum albumin (Sigma) in RPMI for 2 h at 37° C and 5% CO₂, and 10⁵ expanded PBMC added per well coated with LPS, IpaB, IpaC, IpaD, VirG, MxiH or LcrV. For total IgA measurements, serial 2-fold dilutions were performed starting at 7,500 expanded cells/well down to 234 cells/well. Cells were incubated for 5 h at 37° C and 5% CO₂, washed with PBST followed by PBS, and incubated with mouse anti-human pan IgA Biotin Conjugated Antibody (Hybridoma Reagent Laboratory, Baltimore, MD) overnight at 4° C. Subsequently plates were washed with PBST and PBS and labeled with horseradish peroxidase-conjugated Avidin D (Vector Laboratories, Burlingame, CA) for 1 h at room temperature. The substrate 3-Amino-9 ethylcarbazole C (Calbiochem, La Jolla, CA, USA) was added at 100 µL per well for 20 minutes at room temperature and the reaction was stopped with ddH₂O. Final enumeration of specific and total SFC was performed using the Immunospot Series 3B Analyzer ELISPOT reader (Cellular Technologies Ltd, Shaker Heights, OH) with aid of the Immunospot software version 4.0 (Cellular Technologies Ltd).

Adequate expansion of B_M cells, assessed by the frequency of total IgA detected by ELISPOT, is critical to the sensitivity and consistency of this method. Thus, specimens which did not reach a minimum cut-off level following expansion (arbitrarily defined as the 10th percentile of the levels reached by all volunteers at any time point: i.e., > 8,300/10⁶ total IgA SFC/10⁶ expanded cells) were excluded from further analysis. Statistical analysis was performed on the mean number of SFC in antigen-coated wells minus the mean number of SFC in the negative control wells. The limit of detection of antigen-specific/total expanded cells in our assays was 1 in 100,000 (0.001%). The limit of detection of antigen-specific/total IgA-secreting cells in each volunteer was determined by taking into account the maximum number of total IgA SFC for that subject in anti-IgA coated ELISPOT wells. The latter had an acceptable minimum of 0.005% (i.e., 1 specific SFC in the 8,300 total IgA SFC/10⁶ expanded cells cut-off).

2.7. Flow cytometry

Eight volunteers had sufficient PBMC available after expansion pre and post vaccination to enable flow cytometric measurements. Of these subjects, 3 volunteers were responders by LPS IgA in serum and stool by ELISA and 5 were non responders. Of the 3 LPS responders, 2 were also IpaB responders. Expanded PBMC were washed with 1% FBS in PBS and labeled with fluorochrome-labeled mAbs against the following antigens: (1) CD19-ECD (clone J3.119, Beckman-Coulter, Fullerton, CA), (2) CD20-APC-Cy7 (clone L27, BD Biosciences, San Jose, CA), (3) CD27-APC-A700 (clone 1A4CD27, Beckman-Coulter), (4) IgA-Biotin (clone G20-359, BD Biosciences - subsequently labeled with streptavidin-Pacific Orange (InVitrogen, Carlsbad, CA), (5) IgG-PE-Cy5 (clone G18-145, BD Biosciences), (6) integrin α_4/β_7 -Alexa 647 (the anti-integrin α_4/β_7 ACT-1 mAb was kindly provided by Dr. W. Newman, PaxVax Inc., San Diego, CA and conjugated to Alexa 647 using an Alexa 647-labeling kit (Molecular probes, Eugene, OR), and (7) CD14-PacBlue (clone TÜK4 InVitrogen), CD3-PacBlue (clone UCHT1 BD Biosciences) and Vivid (InVitrogen), which were used to exclude cells staining positively with these mAbs using a “dump” channel gating strategy. Incubation with the mAbs was performed in volumes of 50 µl per tube for 20–30 minutes at 4°C, washed with 1% FBS in PBS, and fixed in 300 µl of 1% formaldehyde until run. Events were acquired on a MoFlow flow cytometer/cell sorter system (Beckman-Coulter) and analyzed using WinList 6.0 (Verity Software House, Topsham, ME) software.

2.8. Statistical analysis

Microsoft® Office Excel 2007, GraphPad Prism 5.0, and STATA 9.0 were used for statistical analysis. Our hypotheses were evaluated using non-parametric two-sided tests. Pre- and post-vaccination results were paired. Antigen-specific SFC/10⁶ expanded cells were divided by total IgA SFC. The Wilcoxon signed rank test was used to assess continuous pre- to post-vaccination antigen-specific B_M responses. Wilcoxon Rank Sum was used to compare responders to non-responders. Correlations between seroresponse and B_M responses were assessed using Spearman rho test for continuous variables and Fisher's exact test for dichotomous variables. Two sided p values <0.05 were considered significant.

3. Results

3.1. Evaluation of total and specific SFC after expansion of B_M cells

Twenty four volunteers who received vaccine had sufficient PBMC pre- and post-vaccination to be included in these studies. Thirteen of the 24 subjects were anti-LPS IgA seroresponders. All 13 seroresponders were also mucosal responders by sIgA anti-LPS measured in stool. Three individuals were mucosal responders only with ≥ 4 fold increases in anti-LPS/total stool sIgA without a serum response. Cells from one anti-LPS seroresponder had to be excluded from analysis due to inadequate expansion. LPS-specific B_M cells increased from a median of 7 SFC/10⁶ expanded cells pre-vaccination to a median of 53 SFC/10⁶ expanded cells 28 days post-vaccination ($p=0.005$) (Figure 1). Increases were observed in LPS-specific IgA B_M in 10 out of 12 IgA anti-LPS (83%) seroresponders with adequate expansion. Of 24 vaccinated volunteers who had sufficient PBMC to perform these studies, 7 also displayed an IgA anti-IpaB seroresponse. Mean IgA anti-IpaB B_M cells increased from 4 to 20 SFC/10⁶ expanded cells pre- to post-vaccination ($p=0.062$); 4 out of 7 (57%) IgA IpaB seroresponders manifested an increase in IgA anti-IpaB B_M cells. The median percentages of antigen-specific SFC as a proportion of total IgA SFC showed increases from 0.03% pre-vaccination to 0.20% post-vaccination for LPS (p value = 0.005) and 0.03% pre-vaccination to 0.10% post-vaccination for IpaB ($p=0.2$) among seroresponders (Figure 1). Individuals who were not seroresponders did not exhibit a statistically significant increase in antigen-specific B_M responses pre- to post-vaccination.

Because of the importance of other proteins in *Shigella* pathogenesis, e.g., VirG (IcsA), MxiH, IpaB, IpaC and IpaD [26,27], we studied whether immunization with a live oral *Shigella* vaccine also elicited specific B_M to these antigens. Only sporadic B_M responses were observed to IpaC, IpaD, MxiH, and VirG. No responses were detected against *Y. pestis* LcrV (negative control).

The frequency of B_M can be quantified as the number of anti-LPS or as anti-IpaB specific IgA per 10⁶ expanded PBMC or the % of anti-LPS or anti-IpaB specific IgA divided by the total number of IgA producing cells in expanded PBMC. As can be observed in Figure 2, highly significant correlations were observed between the two methods of quantification.

3.2. Correlation of B_M responses with antibody responses in serum and stool

Mucosal responses to vaccination (i.e., the production of secretory IgA in stool/total ≥ 4 fold pre to post oral vaccination) was observed among 16 out of 24 subjects in this study. Among these 16 subjects, strong correlations were observed when comparing their LPS-specific B_M (SFC/10⁶ expanded cells) frequencies measured on day 28 with their respective post-vaccination peak seroresponse (Figure 3A), peak mucosal (stool) IgA response (Figure 3B), and peak peripheral blood IgA ASC responses (Figure 3C). Strong correlations were also observed among anti-LPS IgA B_M and both mucosal and peripheral anti-LPS IgA responses among the 12 individuals who were seroresponders (data not shown). Of importance, the

correlations remain significant when results were analyzed as percentages of specific anti-LPS IgA B_M divided by total IgA secreting cells in each volunteer. No significant correlations were observed among antibody levels in serum and stool and peripheral B_M cells on day 0 (pre-vaccination, data not shown). Calculations taking into account individual spot sizes (a relative measurement of the amount of antibody produced by each cell) did not change the results. Interestingly, no correlations were observed between the levels of IpaB-specific IgA B_M cells and the serum levels of anti-IpaB IgA antibodies or the levels of anti-IpaB ASC in circulation (stool anti-IpaB antibody levels were not measured)(data not shown). Among the vaccine recipients who exhibited a mucosal response, the frequencies of IgA B_M cells were better correlated with serum LPS IgA titers (Figure 3) than with serum LPS IgG (Spearman Rho 0.5 p=0.05).

3.3 Expression of gut homing molecules in plasmablasts and B_M following immunization

To evaluate whether immunization with CVD 1204 or CVD 1208 resulted in changes in the proportion of B cell subsets of defined phenotypes, PBMC obtained before and after oral vaccination were expanded and stained to examine B cell subsets and expression of the gut homing integrin α_4/β_7 receptor. Cells were sequentially gated based on forward versus side scatter ("lymph region"), followed by the electronic elimination of doublets, dead cells, CD3⁺ T cells and CD14⁺ macrophages. The CD19⁺ cells co-expressing integrin α_4/β_7 (i.e., with the potential to home to the gut) or not were then gated and analyzed for their expression of IgG or IgA; each of these cell subsets was further gated based on their expression of CD27 and CD20 to define the CD19⁺ integrin α_4/β_7 ⁺ CD27⁺ CD20⁺ [largely B_M] and CD19⁺ integrin α_4/β_7 ⁺ CD27⁺ CD20^{-/low} [largely plasmablasts/plasmocytes] subsets [28,29].

The results from these studies demonstrated an increase in IgA secreting cells post-oral vaccination compared to pre-vaccination among the LPS seroresponders. This increase was statistically significant when comparing responders to non-responders by Wilcoxon Rank Sum in the CD19⁺ integrin α_4/β_7 ⁺ IgA⁺ CD27⁺ CD20⁺ as well as CD19⁺ integrin α_4/β_7 ⁺ IgA⁺ CD27⁺ CD20^{-/low} subsets (Figure 4). No significant differences among responders and non-responders were observed in the percentages of any of the other subsets evaluated (i.e., IgG⁺ subsets, CD19⁺ integrin α_4/β_7 ⁻ subsets, CD19⁺ integrin α_4/β_7 ⁺ CD27⁻ CD20⁺ and CD19⁺ integrin α_4/β_7 ⁺ CD27⁻ CD20⁻ subsets).

4. Discussion

Immune responses induced locally provide the first line of defense against the many pathogens that invade the human host via mucosal surfaces. Humans make more IgA than IgG, IgM and IgE combined and allows for active transport of sIgA across mucosal epithelia to facilitate antigen exclusion and neutralization [30]. The normal human colon has a higher proportion of IgA than IgG-producing plasma cells as evidenced by immunohistochemistry [31]. The induction of B_M responses is widely accepted to be a major factor in the ability of vaccines to elicit long lasting, effective immunity. However, the precise role of IgA ASC and B_M cells in primary and secondary [anamnestic] immune responses to infection remains ill defined.

A B_M response can be demonstrated by documenting: (1) an anamnestic secondary antibody response, both stronger and faster than the initial immune response, (2) avidity maturation, and (3) the presence of B_M cells [32]. B_M cells have been described in humans against vaccines known to induce a T-cell dependent response, including live viral vaccines administered parenterally (e.g. smallpox vaccine [33]) and orally (e.g., rotavirus vaccines [34]) and parenteral conjugate vaccines consisting of bacterial polysaccharides (such as pneumococcal [35] and meningococcal [36] capsular polysaccharides) covalently linked to

carrier proteins. B_M cells have also been reported following natural infection with bacterial enteropathogens and after the administration of live oral bacterial enteric vaccines [16,37]. We here report the observation that oral immunization with attenuated *S. flexneri* 2a vaccines elicits the generation specific IgA B_M cells.

Although the immunological correlates of protection following *Shigella* infection have not yet been fully elucidated, high numbers of antigen-specific IgA ASC post oral vaccination have been found to be associated with protection from shigellosis after experimental challenge [8]. Interestingly, the volunteers who have high numbers of antigen-specific ASC after vaccination and are asymptomatic after challenge have low levels of antigen-specific IgA ASC in peripheral blood after secondary antigen exposure [38]. In the present study we observed that a single oral immunization with live attenuated *Shigella* vaccine CVD 1204 or CVD 1208 elicited significant increases in antigen-specific IgA anti-LPS B_M responses among subjects who mounted $\geq$ fourfold specific sIgA or IgA antibody responses as measured by ELISA, respectively, in stool and serum. IpaB seroresponders also exhibited antigen-specific IgA B_M responses to IpaB. Strong correlations were observed between the magnitude of IgA anti-LPS B_M cells and increases in anti-LPS IgA responses, both among individuals who displayed a strong mucosal sIgA stool response and among IgA seroresponders. B_M cells also correlated highly with anti-LPS IgA ASC, both among seroresponders and mucosal responders. It will be important to determine in future experimental challenge studies with wild-type *Shigella* whether antigen-specific IgA and IgG B_M cells are elevated in subjects who had been (or not) previously immunized with oral live attenuated *Shigella* strains and whether B_M cells are associated with protection.

We were surprised that no differences were detected in the magnitude of B_M cell responses among recipients of the slightly reactogenic CVD 1204 or the very well tolerated CVD 1208, or related to the various dosage levels evaluated (10^7 , 10^8 , or 10^9 CFU). An optimistic interpretation of this observation is that the loss of reactogenicity resulting in a well tolerated further attenuated *Shigella* vaccine strains such as strain CVD 1208 is not accompanied by a diminution of the ability to elicit strong immune responses, including the generation of antigen-specific IgA B_M cells.

In previous studies we determined by flow cytometry the proportions of phenotypically defined total B, B_M and IgG⁺ B_M cell populations before and after expansion [16]. This information helped validate the B_M cell ELISPOT assay as well as provide the rationale for limiting the analyses to specimens that showed evidence of appropriate in vitro expansion in the presence of mitogens. By eliminating specimens that exhibited suboptimal expansion of functional cells (defined as $\leq 8,300$ total IgA SFC/ 10^6 expanded cells) in the current dataset, we observed increased levels of circulating antigen-specific IgA B_M responses elicited by oral vaccination of up to 1.1% antigen-specific B_M SFC/total IgA SFC post-vaccination for T-independent antigen LPS and up to 1.0% for the T-dependent antigen IpaB. These proportions of antigen-specific B_M cells are similar to those previously reported in subjects who received efficacious parenteral vaccines. For example, responses to diphtheria and tetanus toxoids elicited specific B_M cells over total IgG secreting cells in the range of 0.01–1% [39]. Recombinant hepatitis B vaccine showed the induction of 0.07% hepatitis B surface antigen-specific Ig secreting cells over total Ig secreting cells; 88% of vaccinees had detectable levels of IgG B_M cells and 76% had detectable levels of IgA and IgM B_M cells [40]. In natural cholera infection, anti-LPS IgA B_M cells have been reported to be 0.6% of total IgA B_M cells and proposed to play an important role in the anamnestic mucosal immune response [37]. To our knowledge, the data included in the present manuscript is the first description of the induction of specific IgA B_M cell response to LPS in recipients of an oral live-attenuated bacterial vaccine.

Since *Shigella* enters the host via the gut, it is important to define the local immune responses elicited by immunization. Due to regulatory constraints and other factors, intestinal mucosal biopsies are very difficult to obtain following vaccine administration or challenge with wild-type organisms. Alternative ways to investigate whether antigen-specific IgA and IgG ASC and B_M found to be elevated in the periphery after oral vaccination have the potential to migrate to the gut mucosa involve indirect measurements, such as sIgA in stool and the measurement of gut homing molecule expression, e.g., integrin α_4/β_7 , on IgA⁺ and IgG⁺ ASC and B_M cells. Although in our experience flow cytometry is not as sensitive as ELISPOT in assessing antigen-specific B_M responses, it nevertheless allowed us to characterize the phenotype of the B cell subsets that increased following oral vaccination.

In the present study we found significant increases in the proportion of vaccine-induced CD19⁺ CD27⁺ CD20⁺ integrin α_4/β_7 ⁺ IgA⁺ B_M cells with a gut homing pattern in seroresponders as compared to non-seroresponders. Smaller, yet significant, increases have also been observed in CD19⁺ CD27⁺ CD20⁻ integrin α_4/β_7 ⁺ IgA⁺ cells (a phenotype typically associated with plasmablasts) [29]. While B_M only express surface Ig, expression of Ig in plasmablasts is largely cytoplasmic with lower levels of surface Ig expression [41]. Because this manuscript is focused on B_M and sufficient cells were not available for independent staining panels (i.e., extracellular and intracellular IgA and/or IgG staining), we stained extracellularly for Ig. Thus, it is likely that the smaller differences observed in integrin α_4/β_7 ⁺ IgA⁺ plasmablasts in responders as compared to non-responders are the result of a lower efficiency in the detection of IgG⁺ and IgA⁺ plasmablasts. Future studies using intracellular Ig staining will allow this issue to be addressed directly. In contrast to the increases in IgA⁺ cells, this phenomenon was not observed in CD19⁺ α_4/β_7 ⁺ IgG B_M cells (irrespective of expression of CD27 and/or CD20). In sum, we found that IgA⁺, but not IgG⁺, CD19⁺ B cells expressing the gut homing receptor integrin α_4/β_7 increased 28 days after oral vaccination, and that gating on subsets based on the expression of CD27 and CD20 enhanced the ability to identify this population. This observation is consistent with Crotty's original description of the B_M cell ELISPOT where sorting experiments were used to identify that B_M have a CD27⁺ CD20⁺ phenotype [25]. The present results suggest an increase in circulating *Shigella* specific IgA B_M and plasmablasts with gut homing potential in individuals in whom immunization elicited anti-*Shigella* humoral responses.

A limitation of this study is the relatively small number of volunteers who could be evaluated, which is largely due to the fact that it employed "convenience" specimens based on availability. In spite of this limitation, we observed strong, statistically significant, specific anti-LPS IgA B_M responses and associations with anti-LPS IgA antibody and ASC levels. However, it is very likely that the small sample size provided insufficient power to adequately estimate the presence and association of anti-IpaB IgA B_M cells and seroresponses. Future studies will address this issue by evaluating larger numbers of subjects and vaccine candidates.

Virulent *Shigella* target the M cells that overlie gut-associated lymphoid tissue and rapidly attain an intracellular niche within epithelial cells. SIgA anti-*Shigella* antibodies can prevent mucosal invasion. Moreover, wild type organisms coated with such antibody elicit less profound inflammatory responses than *Shigella* not coated with antibody [42]. It is not known how long sIgA intestinal and serum IgG anti-*Shigella* antibodies stimulated by clinical infection or oral immunization with *Shigella* vaccines remain elevated at levels adequate to prevent invasion and clinical illness. If antibody levels decline to non-protective levels, it falls upon B_M cells to mount a sufficiently rapid anamnestic response to limit the clinical consequence of exposure to virulent *Shigella*. Experimental challenge studies with immunologically-naïve healthy adult volunteers and epidemiologic investigations of point

source outbreaks suggest that for most *Shigella* serotypes the usual incubation is ~ 1–3 days from the point of ingestion of virulent organisms until the onset of clinical illness. Thus, an effective anamnestic response mediated by B_M cells must occur within a mere 1–2 days. Few reports have directly addressed the duration of immunity that follows an initial clinical *Shigella* infection in the absence of repetitive boosting (as would occur in an endemic situation). Similarly, to our knowledge the only controlled studies that describe the duration of protection conferred by immunization with live oral *Shigella* vaccines were those of Mel et al who immunized children living in an endemic area where *Shigella* was highly seasonal [43,44]. Oral immunization with four doses over two weeks of streptomycin-dependent live oral *Shigella* vaccines led to significant serotype-homologous protection that endured for one year [44]. However, the administration of just a single oral booster dose was able to prolong the protection for an additional year [44].

In summary, oral vaccination with a single oral dose of a live-attenuated *Shigella* vaccine elicited antigen-specific IgA B_M responses among IgA seroresponders and sIgA mucosal responders with a high degree of correlation between antigen-specific B_M cells and both serum and mucosal titers as well as antigen-specific ASC in peripheral circulation. Mucosal sIgA responses are more difficult to measure than serum antibody, as is the measurement of small number of IgA B_M cells. Nevertheless, we observed positive correlations between antigen-specific anti-LPS B_M cells and antigen-specific anti-LPS sIgA/total IgA in stool. As *Shigella* invades via the gastrointestinal mucosa, fecal sIgA responses likely play an important role as the first line of defense in protection. Antigen-specific IgA B_M cells should be further studied in natural infection and challenge studies to determine their role in the anamnestic responses and protection against this important and devastating pathogen. For expedited identification of future candidate vaccines, it will be necessary to identify reliable surrogates of protection against symptomatic infection. A strong correlation between anti-LPS IgA B_M cells and peak anti-LPS IgA antibody and ASC responses further reinforces our contention that B_M cells may be an important indicator for long-term humoral immunity and a possible surrogate of protection against shigellosis.

5. Conclusions

Oral vaccination with live-attenuated *S. flexneri* 2a elicited detectable IgA B_M cells to LPS and, to a lesser extent, to IpaB. Positive correlations were observed between anti-LPS IgA B_M and both anti-LPS secretory IgA /total IgA in stool and anti-LPS IgA in serum as well as circulating ASC. In addition, flow cytometric analyses revealed significant differences in IgA⁺ B_M subsets expressing integrin α_4/β_7 pre to post vaccination when comparing seroresponders to non-responders. Our results support the contention that B_M cells may be an important indicator for long-term humoral immunity and a possible surrogate of protection against shigellosis

Abbreviations

ASC	antibody secreting cell
B _M	memory B cells
CD	cluster designation
CDC	Center for Disease Control and Prevention
CVD	Center for Vaccine Development
CFU	colony forming units
CMI	cell-mediated immunity

ELISA	enzyme-linked immunosorbent assay
ELISPOT	enzyme-linked immunosorbent spot
Ig	immunoglobulin
Ipa	Invasion plasmid antigen
IU	international units
LPS	lipopolysaccharide
mAb	monoclonal antibodies
mL	milliliter
mM	millimolar
NIH	National Institutes of Health
PBMC	peripheral blood mononuclear cells
PBS	phosphate buffered saline
PBST	phosphate buffered saline + 0.05% Tween 20
PCR	polymerase chain reaction
PWM	pokeweed mitogen
cRPMI	complete Roswell Park Memorial Institute medium
SAC	<i>Staphylococcus aureus</i> Cowan
SFC	spot forming cells
µg	microgram
uM	micromolar
VTEU	Vaccine and Treatment Evaluation Unit
WHO	World Health Organization

Acknowledgments

We thank the volunteers for participating in the clinical trial, the clinical and regulatory staff at the CVD, Drs. William Blackwelder and Yhukun Wu for helpful statistical discussions, and Dr. S. Crotty for providing PWM. Support for this research was provided by NIH R01-AI057927 (to M.B.S.), U19-AI-082655 (CCHI; to M.B.S.), K23-AI065759 (to J.S.) and N01-AI25461 (VTEU, to M.M.L). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Allergy and Infectious Diseases or the NIH.

References

1. Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, Swerdlow DL, Sansonetti PJ, Adak GK, Levine MM. Global burden of Shigella infections: implications for vaccine development and implementation of control strategies. Bull. World Health Organ. 1999; 77:651–666. [PubMed: 10516787]
2. Levine MM, Kotloff KL, Barry EM, Pasetti MF, Sztein MB. Clinical trials of Shigella vaccines: two steps forward and one step back on a long, hard road. Nat. Rev. Microbiol. 2007; 5:540–553. [PubMed: 17558427]
3. Pickering LK. Antimicrobial resistance among enteric pathogens. Semin. Pediatr. Infect. Dis. 2004; 15:71–77. [PubMed: 15185189]

4. Bioterrorism Agents/Diseases. Center for Disease Control and Prevention Website. 2010. Available from: URL: <http://www.bt.cdc.gov/agent/agentlist.asp>
5. Bennish ML, Wojtyniak BJ. Mortality due to shigellosis: community and hospital data. *Rev. Infect. Dis.* 1991; 13 Suppl 4:S245–S251. [PubMed: 2047645]
6. Future needs and directions for Shigella vaccines. *Weekly Epidemiological Record.* 2006; 81:51–58.
7. Kotloff KL, Pasetti MF, Barry EM, Nataro JP, Wasserman SS, Sztein MB, Picking WD, Levine MM. Deletion in the Shigella Enterotoxin Genes Further Attenuates Shigella flexneri 2a Bearing Guanine Auxotrophy in a Phase 1 Trial of CVD 1204 and CVD 1208. *J. Infect. Dis.* 2004; 190:1745–1754. [PubMed: 15499528]
8. Kotloff KL, Losonsky GA, Nataro JP, Wasserman SS, Hale TL, Taylor DN, Newland JW, Sadoff JC, Formal SB, Levine MM. Evaluation of the safety, immunogenicity, and efficacy in healthy adults of four doses of live oral hybrid Escherichia coli- Shigella flexneri 2a vaccine strain EcSf2a-2. *Vaccine.* 1995; 13:495–502. [PubMed: 7639017]
9. Cohen D, Block C, Green MS, Lowell G, Ofek I. Immunoglobulin M, A, and G antibody response to lipopolysaccharide O antigen in symptomatic and asymptomatic Shigella infections. *J Clin. Microbiol.* 1989; 27:162–167. [PubMed: 2463995]
10. Oaks EV, Hale TL, Formal SB. Serum immune response to Shigella protein antigens in rhesus monkeys and humans infected with Shigella spp. *Infect. Immun.* 1986; 53:57–63. [PubMed: 3721580]
11. Oberhelman RA, Kopecko DJ, Salazar-Lindo E, Gotuzzo E, Buysse JM, Venkatesan MM, Yi A, Fernandez-Prada C, Guzman M, Leon-Barua R. Prospective study of systemic and mucosal immune responses in dysenteric patients to specific Shigella invasion plasmid antigens and lipopolysaccharides. *Infect. Immun.* 1991; 59:2341–2350. [PubMed: 2050402]
12. Barzu S, Nato F, Rouyre S, Mazie JC, Sansonetti P, Phalipon A. Characterization of B-cell epitopes on IpaB, an invasion- associated antigen of Shigella flexneri: identification of an immunodominant domain recognized during natural infection. *Infect. Immun.* 1993; 61:3825–3831. [PubMed: 7689541]
13. Samandari T, Kotloff KL, Losonsky GA, Picking WD, Sansonetti PJ, Levine MM, Sztein MB. Production of IFN-gamma and IL-10 to Shigella invasins by mononuclear cells from volunteers orally inoculated with a Shiga toxin-deleted Shigella dysenteriae type 1 strain. *J. Immunol.* 2000; 164:2221–2232. [PubMed: 10657678]
14. Traggiai E, Puzone R, Lanzavecchia A. Antigen dependent and independent mechanisms that sustain serum antibody levels. *Vaccine.* 2003; 21 Suppl 2:S35–S37. [PubMed: 12763680]
15. Li A, Karnell A, Huan PT, Cam PD, Minh NB, Tram LN, Quy NP, Trach DD, Karlsson K, Lindberg G. Safety and immunogenicity of the live oral auxotrophic Shigella flexneri SFL124 in adult Vietnamese volunteers. *Vaccine.* 1993; 11:180–189. [PubMed: 8438616]
16. Simon JK, Wahid R, Maciel M Jr, Picking WL, Kotloff KL, Levine MM, Sztein MB. Antigen-specific B memory cell responses to lipopolysaccharide (LPS) and invasion plasmid antigen (Ipa) B elicited in volunteers vaccinated with live-attenuated Shigella flexneri 2a vaccine candidates. *Vaccine.* 2009; 27:565–572. [PubMed: 19022324]
17. Sztein MB, Wasserman SS, Tacket CO, Edelman R, Hone D, Lindberg AA, Levine MM. Cytokine production patterns and lymphoproliferative responses in volunteers orally immunized with attenuated vaccine strains of Salmonella typhi. *J. Infect. Dis.* 1994; 170:1508–1517. [PubMed: 7995991]
18. Westphal O, Jann K, Himmelsbach K. Chemistry and immunochemistry of bacterial lipopolysaccharides as cell wall antigens and endotoxins. *Prog. Allergy.* 1983; 33:9–39. [PubMed: 6187018]
19. Espina M, Ausar SF, Middaugh CR, Baxter MA, Picking WD, Picking WL. Conformational stability and differential structural analysis of LcrV, PcrV, BipD, and SipD from type III secretion systems. *Protein Sci.* 2007; 16:704–714. [PubMed: 17327391]
20. Espina M, Olive AJ, Kenjale R, Moore DS, Ausar SF, Kaminski RW, Oaks EV, Middaugh CR, Picking WD, Picking WL. IpaD localizes to the tip of the type III secretion system needle of Shigella flexneri. *Infect. Immun.* 2006; 74:4391–4400. [PubMed: 16861624]

21. Harrington A, Darboe N, Kenjale R, Picking WL, Middaugh CR, Birket S, Picking WD. Characterization of the interaction of single tryptophan containing mutants of IpaC from *Shigella flexneri* with phospholipid membranes. *Biochemistry*. 2006; 45:626–636. [PubMed: 16401091]
22. Darboe N, Kenjale R, Picking WL, Picking WD, Middaugh CR. Physical characterization of MxiH and PrgI, the needle component of the type III secretion apparatus from *Shigella* and *Salmonella*. *Protein Sci*. 2006; 15:543–552. [PubMed: 16501225]
23. Birket SE, Harrington AT, Espina M, Smith ND, Terry CM, Darboe N, Markham AP, Middaugh CR, Picking WL, Picking WD. Preparation and characterization of translocator/chaperone complexes and their component proteins from *Shigella flexneri*. *Biochemistry*. 2007; 46:8128–8137. [PubMed: 17571858]
24. Van De Verg L, Herrington DA, Murphy JR, Wasserman SS, Formal SB, Levine MM. Specific immunoglobulin A-secreting cells in peripheral blood of humans following oral immunization with a bivalent *Salmonella typhi*-*Shigella sonnei* vaccine or infection by pathogenic *S. sonnei*. *Infect. Immun*. 1990; 58:2002–2004.
25. Crotty S, Aubert RD, Glidewell J, Ahmed R. Tracking human antigen-specific memory B cells: a sensitive and generalized ELISPOT system. *J. Immunol. Methods*. 2004; 286:111–122. [PubMed: 15087226]
26. Parsot C. *Shigella* spp. and enteroinvasive *Escherichia coli* pathogenicity factors. *FEMS Microbiol. Lett*. 2005; 252:11–18. [PubMed: 16182469]
27. Zhang L, Wang Y, Olive AJ, Smith ND, Picking WD, De Guzman RN, Picking WL. Identification of the MxiH needle protein residues responsible for anchoring invasion plasmid antigen D to the type III secretion needle tip. *J. Biol. Chem*. 2007; 282:32144–32151. [PubMed: 17827155]
28. Wrammert J, Smith K, Miller J, Langley WA, Kokko K, Larsen C, Zheng NY, Mays I, Garman L, Helms C, James J, Air GM, Capra JD, Ahmed R, Wilson PC. Rapid cloning of high-affinity human monoclonal antibodies against influenza virus. *Nature*. 2008; 453:667–671. [PubMed: 18449194]
29. Avery DT, Ellyard JJ, Mackay F, Corcoran LM, Hodgkin PD, Tangye SG. Increased expression of CD27 on activated human memory B cells correlates with their commitment to the plasma cell lineage. *J Immunol*. 2005; 174:4034–4042. [PubMed: 15778361]
30. Brandtzaeg P. Induction of secretory immunity and memory at mucosal surfaces. *Vaccine*. 2007; 25:5467–5484. [PubMed: 17227687]
31. Brandtzaeg P. The mucosal immune system and its integration with the mammary glands. *J Pediatr*. 2010; 156:S8–S15. [PubMed: 20105666]
32. Tarlinton D. B-cell memory: are subsets necessary? *Nat. Rev. Immunol*. 2006; 6:785–790. [PubMed: 16998511]
33. Crotty S, Felgner P, Davies H, Glidewell J, Villarreal L, Ahmed R. Cutting edge: long-term B cell memory in humans after smallpox vaccination. *J. Immunol*. 2003; 171:4969–4973. [PubMed: 14607890]
34. Rojas OL, Caicedo L, Guzman C, Rodriguez LS, Castaneda J, Uribe L, Andrade Y, Pinzon R, Narvaez CF, Lozano JM, De VB, Franco MA, Angel J. Evaluation of circulating intestinally committed memory B cells in children vaccinated with attenuated human rotavirus vaccine. *Viral Immunol*. 2007; 20:300–311. [PubMed: 17603846]
35. Clutterbuck EA, Salt P, Oh S, Marchant A, Beverley P, Pollard AJ. The kinetics and phenotype of the human B-cell response following immunization with a heptavalent pneumococcal-CRM conjugate vaccine. *Immunology*. 2006; 119:328–337. [PubMed: 17067312]
36. Kelly DF, Snape MD, Clutterbuck EA, Green S, Snowden C, Diggle L, Yu LM, Borkowski A, Moxon ER, Pollard AJ. CRM197-conjugated serogroup C meningococcal capsular polysaccharide, but not the native polysaccharide, induces persistent antigen-specific memory B cells. *Blood*. 2006; 108:2642–2647. [PubMed: 16675705]
37. Harris AM, Bhuiyan MS, Chowdhury F, Khan AI, Hossain A, Kendall EA, Rahman A, LaRocque RC, Wrammert J, Ryan ET, Qadri F, Calderwood SB, Harris JB. Antigen-specific memory B-cell responses to *Vibrio cholerae* O1 infection in Bangladesh. *Infect Immun*. 2009; 77:3850–3856. [PubMed: 19528207]

38. Kotloff KL, Nataro JP, Losonsky GA, Wasserman SS, Hale TL, Taylor DN, Sadoff JC, Levine MM. A modified *Shigella* volunteer challenge model in which the inoculum is administered with bicarbonate buffer: clinical experience and implications for *Shigella* infectivity. *Vaccine*. 1995; 13:1488–1494. [PubMed: 8578831]
39. Nanan R, Heinrich D, Frosch M, Kreth HW. Acute and long-term effects of booster immunisation on frequencies of antigen-specific memory B-lymphocytes. *Vaccine*. 2001; 20:498–504. [PubMed: 11672915]
40. Tuaillon E, Tabaa YA, Petitjean G, Huguet MF, Pajaux G, Fondere JM, Ponselle B, Ducos J, Blanc P, Vendrell JP. Detection of memory B lymphocytes specific to hepatitis B virus (HBV) surface antigen (HBsAg) from HBsAg-vaccinated or HBV-immunized subjects by ELISPOT assay. *J Immunol. Methods*. 2006; 315:144–152. [PubMed: 16959261]
41. Jaimes MC, Rojas OL, Kunkel EJ, Lazarus NH, Soler D, Butcher EC, Bass D, Angel J, Franco MA, Greenberg HB. Maturation and trafficking markers on rotavirus-specific B cells during acute infection and convalescence in children. *J. Virol*. 2004; 78:10967–10976. [PubMed: 15452217]
42. Boullier S, Tanguy M, Kadaoui KA, Caubet C, Sansonetti P, Corthesy B, Phalipon A. Secretory IgA-mediated neutralization of *Shigella flexneri* prevents intestinal tissue destruction by down-regulating inflammatory circuits. *J. Immunol*. 2009; 183:5879–5885. [PubMed: 19828639]
43. Mel D, Gangarosa EJ, Radovanovic ML, Arsic BL, Litvinjenko S. Studies on vaccination against bacillary dysentery. 6. Protection of children by oral immunization with streptomycin-dependent *Shigella* strains. *Bull. World Health Organ*. 1971; 45:457–464. [PubMed: 4948417]
44. Mel DM, Arsic BL, Radovanovic ML, Litvinjenko SA. Live oral *Shigella* vaccine: vaccination schedule and the effect of booster dose. *Acta Microbiol. Acad. Sci. Hung*. 1974; 21:109–114. [PubMed: 4613128]

Figure 1. Antigen-specific IgA B_M cell responses

Shown are recipients of 10^7 (triangles), 10^8 (circles) and 10^9 (squares) CFU of CVD 1204 (solid lines) and CVD 1208 (broken lines) who mounted a ≥ 4 -fold rise of anti-LPS ($n=12$) and anti-IpaB ($n=7$) IgA pre to post vaccination and evidenced appropriate B_M expansion in vitro; LPS (a) and IpaB (b) ELISPOT performed on days 0 and 28; comparisons made by Wilcoxon Signed Rank Test for 1204 and 1208 combined. Results are expressed as the % of specific B_M SFC per total IgA⁺ expanded cell populations; horizontal lines represent the median of the corresponding groups.

Figure 2. Correlation of antigen-specific B_M cells per 10⁶ expanded PBMC with antigen-specific B_M cells/total IgA B_M cells

B_M cells specific for LPS (a) and IpaB (b) /10⁶ expanded cells 28 days after immunization were plotted against the % of specific SFC divided by total IgA-secreting expanded cell populations for all volunteers studied; dotted lines represent the 95% confidence interval.

Figure 3. Correlation of anti-LPS B_M cell responses with serum and stool antibodies
 Recipients of CVD 1204 and CVD 1208 who mounted a ≥ 4 -fold rise of anti-LPS/total stool sIgA pre to post vaccination (mucosal responders) and evidenced appropriate B_M expansion in vitro ($n=16$) were included in the analysis. Peak serum IgA titers to LPS (a), peak stool IgA specific for LPS divided by total IgA (b), and peak LPS specific IgA ASC (c) were plotted against the number of LPS-specific B_M cells / 10^6 expanded cells 28 days after immunization; dotted lines represent the 95% confidence interval.

Figure 4. Pre to post vaccination assessment of B cell subsets expressing the gut homing receptor integrin α_4/β_7

Shown are the % of CD19⁺ integrin α_4/β_7 ⁺ cells co-expressing CD27 and CD20 (left panels) and CD27 but not CD20 (right panels) pre and day 28 post vaccination for IgA⁺ (upper panels) and IgG⁺ cells (lower panels) among LPS seroresponders and non-responders. *p < 0.05 by Wilcoxon Rank Sum comparing net change in % of the indicated subsets post minus pre immunization among LPS seroresponders versus non-responders; LPS seroresponse is defined as ≥ 4 -fold rise of antigen-specific antibody in serum (all responders were both IgA and IgG seroresponders).