Published in final edited form as:

Biochemistry. 2011 April 12; 50(14): 2820–2828. doi:10.1021/bi102002v.

Alternative initial proton acceptors for the D pathway of *Rhodobacter sphaeroides* cytochrome *c* oxidase

Lakshman Varanasi and Jonathan Hosler*

Department of Biochemistry, University of Mississippi Medical Center, Jackson, MS 39216

Abstract

In order to characterize protein structures that control proton uptake, forms of cytochrome *c* oxidase (CcO) containing a carboxyl or a thiol group in line with the initial, internal waters of the D pathway for proton transfer have been assayed in the presence and absence of subunit III. Subunit III provides approximately half of the protein surrounding the entry region of the D pathway. The mutant N139D-D132N contains a carboxyl group 6Å within the D pathway and lacks the normal, surface-exposed proton acceptor, Asp-132. With subunit III, the steady-state activity of this mutant is slow but once subunit III is removed its activity is the same as wild-type CcO lacking subunit III ($\sim 1800 \text{ H}^+ \text{ s}^{-1}$). Thus, a carboxyl group $\sim 25\%$ within the pathway enhances proton uptake even though the carboxyl has no direct contact with bulk solvent. Protons from solvent apparently move to internal Asp-139 through a short file of waters, normally blocked by subunit III. Cysteine-139 also supports rapid steady-state proton uptake, demonstrating that an anion other than a carboxyl can attract and transfer protons into the D pathway. When both Asp-132 and Asp/Cys-139 are present, the removal of subunit III increases CcO activity to rates greater than that of normal CcO due to simultaneous proton uptake by two initial acceptors. The results show how the environment of the initial proton acceptor for the D pathway in these CcO forms dictates the pH range of CcO activity, with implications for the function of Asp-132, the normal proton acceptor.

The *aa*₃-type cytochrome *c* oxidase complex (CcO) is the final member of the electron transfer chain of oxidative phosphorylation in the inner membrane of mitochondria and in the cytoplasmic membrane of many bacteria. In order to generate a voltage gradient across these membranes, CcO transfers electrons from its outer (positive) surface and protons from its inner (negative) surface to a buried heme-Cu active site where the charges combine as O₂ is reduced to water (1-3). As a proton pump, CcO also uses the energy of O₂ reduction to transfer protons completely through the protein, across the membrane (3-6). The four-subunit *aa*₃-type CcO of the bacterium *Rhodobacter sphaeroides* is one of the popular, mutable models of the catalytic core of mitochondrial CcO (7,8).

Cytochrome *c* oxidase (CcO) takes up the protons necessary for catalysis via two long pathways, the D and K paths, leading from the inner surface of the complex to the buried active site (Fig. 1) (2,4,9,10). Protons are introduced into the D pathway by Asp-132 of subunit I (*R. sphaeroides* numbering) and transferred 26Å to Glu-286, near the active site, through a series of hydrogen-bonded water molecules (11-14). From Glu-286, protons branch to O₂ reduction intermediates in the active site or to the site of proton pumping (15-17). The K pathway begins far from the entry region of the D pathway, at Glu-101 of subunit II (18). The K pathway transfers protons primarily via protein side chains to the cross-linked Tyr-288/His-284 group of the active site (6,11,12).

*Phone: 601-984-1861; FAX: 601-984-1501; jhosler@umc.edu.

Aspartate-132 is defined as the initial proton acceptor of the D pathway based upon its absolute requirement for D pathway activity and its partial exposure on the surface of the protein. Alteration of Asp-132 to alanine, histidine or asparagine essentially eliminates proton uptake (19,20), indicating that Asp-132 provides the sole entry point into the D pathway of normal CcO. Only one of the carboxyl oxygens of Asp-132 is exposed at the bottom of narrow cavity some 6-10 Å deep (21). The protein partner of subunit I in the membrane, subunit III, provides about half of the protein residues that form the cavity surrounding Asp-132 (21-24).

Subunit III can be removed from *R. sphaeroides* CcO using detergent, leaving an active CcO form composed of subunits I and II (25). The removal of subunit III results in a sizeable acid shift (greater than one pH unit) of the pH profile of proton uptake into the D pathway (20,26). The shift apparently reflects a decrease in the affinity of Asp-132 for protons (20,26) consistent with a greater exposure of Asp-132 to solvent water and increased stability of the carboxyl anion.

The removal of subunit III also restores significant rates of proton uptake ($\sim 400 \text{ H}^+ \text{ s}^{-1}$) to D pathways inactivated by the alteration of Asp-132 to alanine, histidine or asparagine (20,25,27). The absence of subunit III allows protons from solvent to access components of the D pathway beyond residue 132, such as the crystal water seen between Asp-132 and Asn-139 in wild-type CcO structures (14,17,20,28). Ergo, subunit III normally provides a structural barrier that prevents protons from solvent from bypassing Asp-132.

Even following the removal of subunit III, steady-state proton uptake by the D pathways of D132A III (-) and D132H III (-) is considerably slower than that of WT III (-) ($\sim 400 \text{ s}^{-1}$ vs $\sim 1800 \text{ s}^{-1}$). The addition of arachidonic acid to D132A III (-) stimulates the rate of proton uptake, restoring up to 60% of normal TN_{max} , as the carboxyl of the fatty acid takes over as the initial proton acceptor (20,25). This demonstrates the utility of an anionic proton acceptor at the entrance of the pathway even though full activity is not restored and the precise location of the fatty acid carboxyl group remains unknown.

Work in *Paracoccus denitrificans* (29) and *R. sphaeroides* (30) has shown that mutant CcOs with a substitution of aspartic acid for an asparagine that lies in the D pathway 6 Å above Asp-132 (Asn-131 in *Paracoccus*, Asn-139 in *Rhodobacter*) retain high rates of O_2 reduction activity. Consistent with this, N139D of *R. sphaeroides* exhibits rapid proton uptake in single turnover experiments (30,31). In a study of proton pumping, Asp-132 of N139D was altered to asparagine, creating N139D-D132N (32). Of particular interest for this study of proton uptake is the finding that N139D-D132N III (+) shows greater activity than D132A/H/N III (+) ($\sim 250 \text{ s}^{-1}$ vs $\sim 40 \text{ s}^{-1}$) (20,25,32,33). This suggests that Asp-139, even though buried, facilitates the entry of solvent protons into the D pathway.

Based on previous results (20,25), we first hypothesized that the access of solvent protons to Asp-139 of N139D-D132N is restricted by subunit III. In fact, we find that without subunit III, buried Asp-139 supports as much CcO activity as surface-exposed Asp-132 of WT III (-). We propose that Asp-139 becomes the initial proton acceptor for the D pathway in the absence of Asp-132 and subunit III. A short file of waters provides the likely pathway for proton transfer between Asp-139 and solvent. The general utility of an anionic proton acceptor is demonstrated by examining CcO forms with cysteine at position 139. Further experiments show that the environment of the initial acceptor can control the pH range of CcO activity, that multiple acceptors can increase proton uptake to rates greater than normal CcO, and that proton uptake by the D pathway is inherently more limiting to CcO activity than proton uptake by the K pathway.

Materials and Methods

Materials

Ni²⁺-NTA agarose was obtained from Qiagen. Arachidonic acid, methyl arachidonate (mAA), horse heart cytochrome *c*, CCCP, valinomycin and soybean phospholipids (asolectin) were from Sigma. Stock solutions (50 mM) of arachidonic acid and mAA were prepared in ethanol and stored at -20° C. Soybean phospholipid (asolectin, Sigma type II-S) was recrystallized using the procedure of Sone et al (34). N-dodecyl β -D-maltoside was from Anatrace. All other chemicals were of ACS grade.

Construction of N139C-D132N and N139D-D132A

A pUC18 construct with a ~2 kbp *HindIII* - *EcoRI* fragment containing *R. sphaeroides* *coxI*, the gene for subunit I of CcO, with the N139C mutation, was obtained from Prof. R.B. Gennis, University of Illinois. Quik-Change mutagenesis (Stratagene) was used to insert the D132N mutation and the presence of the N139C and the D132N mutation, and the absence of other alterations, was verified by DNA sequencing. The altered *EcoRI-HindIII* fragment was then excised from the pUC plasmid and moved into the broad host range vector pRK415 (35) to create pJG273. This plasmid was conjugated into *R. sphaeroides* JS100, a strain lacking the wild-type gene for subunit I (36), for the expression of N139C-D132N. The same procedure was used to create N139D-D132A, starting from the *HindIII* - *EcoRI* fragment of *coxI* containing the N139D mutation. The *R. sphaeroides* CcO mutants N139D (30), N139C and N139D-D132N (32) were obtained from the Gennis laboratory, University of Illinois.

CcO isolation, subunit III removal, COV preparation and activity measurements

Bacterial growth and oxidase purification by Ni-affinity chromatography were performed as in Zhen et al (37). The removal of subunit III was performed as in Mills et al. (25). Measurement of the rates of O₂ reduction (consumption) by purified, detergent-solubilized CcO were performed using an oxygen electrode as described previously (25) in assay mixtures containing 25 mM MES-KOH, pH 6.5, adjusted to 50 mM *I* with KCl, 0.3 mM TMPD, 3.0 mM ascorbate, 0.1% dodecyl maltoside (DDM), 1000 units of catalase, 1 mg/ml soybean phospholipids, 5-25 pmol CcO and horse cytochrome *c* and other additions as noted. For determinations of CcO activity at different pH values, the buffer was 25 mM MES-KOH for pH 6.2 to 6.8, 25 mM Hepes-KOH for pH 6.8 to 8.2 and 25 mM CHES-KOH for pH 8.2 to 9.2. Each buffer was adjusted to a final ionic strength of 50 mM using KCl. The actual pH was measured in the O₂ electrode chamber before and after each activity measurement. The rates of non-enzymatic reduction of O₂ by ascorbate/TMPD at each pH were subtracted. Apparent p*K*_a values for steady-state O₂ reduction were determined by fitting the pH dependence data to a sigmoidal function that assumes a single p*K*_a using the equation $TN = TN_{min} + (TN_{max} - TN_{min}) / (1 + 10^{(pH - pK_a)})$ in GraphPad Prism 5.0.

Cytochrome *c* oxidase was reconstituted into phospholipid vesicles (COVs) as previously described (30). Controlled steady-state activities of the COVs were measured using the O₂ electrode in a reaction mixture containing 25 mM Hepes, pH 7.4, 24 mM KCl, plus ascorbate, TMPD, horse cytochrome *c* and catalase as above. Uncontrolled (uncoupled) activities were measured in the same reaction mixture plus 5 μ M CCCP and 3 μ M valinomycin.

Results

The effect of removing subunit III on the steady-state proton uptake activity of D132N and the Asn-139 mutants

Beginning with normal CcO, the alteration of Asp-132, the initial proton acceptor of the D pathway, to alanine, histidine or asparagine (D132A/H/N) nearly eliminates proton uptake by this pathway (19,20). Slow O₂ reduction activity continues ($\sim 40 \text{ e}^- \text{ s}^{-1}$), as substrate protons normally supplied by the D pathway are slowly taken up by another route, apparently from the opposite side of the complex (25,33,38). The removal of subunit III from D132A and D132H restores steady-state CcO activity to rates 20-25% that of normal CcO by increasing the rate of D pathway proton uptake to $\sim 400 \text{ H}^+ \text{ s}^{-1}$ (20,25). (All proton uptake rates presented here are those for substrate protons.) D132N exhibits the same behavior as D132A/H: removing subunit III increases its low activity to a $\text{TN}_{\text{max}} \sim 400 \text{ s}^{-1}$ ((27); Table 1).

Starting with D132N, the further alteration of Asn-139 to aspartic acid creates N139D-D132N (32), thereby placing a carboxyl group within the D pathway, 6 Å above the normal position of Asp-132 (Figs. 1 & 2). Computer modeling of the Asn-139→Asp-139 substitution into the structure of the subunit I-II form of wild-type *R. sphaeroides* CcO (2GSM; (39)), using the Swiss PDB Viewer DeepView program (40), shows that the carboxyl of Asp-139 occupies nearly the same position as the amide group of Asn-139 in normal CcO. The group has no exposure to bulk solvent. One carboxyl O of Asp-139 is 2.8 Å from the water above (O to O, center to center) while the other lies 2.9 Å from the water below, which is the water normally seen between Asp-132 and Asn-139 (Fig. 2). While a structure of the double mutant N139D-D132N is not yet available, a high resolution structure of N139D of *R. sphaeroides* (Jiapeng Zhu and R. Gennis, Univ. Illinois, unpublished) agrees with the modeled position of Asp-139. In addition, the same position for the equivalent of Asp-139 is seen in a structure of N131D of *Paracoccus denitrificans*, although the water normally present above Asn-139 is not present in the structure of N131D (41). The carboxyl of Asp-139 appears well placed to transfer protons, consistent with published results showing high activity for N131D of *P. denitrificans* (29,41) and N139D of *R. sphaeroides* (30,42), plus the observation of rapid proton uptake in single turnover experiments (30,31). Prior to the removal of subunit III, the double mutant N139D-D132N exhibits low activity with respect to normal CcO (Table 1; (32)) due to absence of Asp-132. However, with the removal of subunit III the O₂ reduction rate of N139D-D132N increases to $\sim 1800 \text{ s}^{-1}$, the same as WT III (-) (Table 1). This shows that Asp-139 of N139D-D132N and Asp-132 of wild-type CcO are equally efficient at supporting steady-state proton uptake, as long as subunit III is removed to allow protons from solvent to access Asp-139.

With subunit III in place, the activity of N139D-D132N is low but significantly greater than D132N III (+) ($\sim 300 \text{ s}^{-1}$ vs $\sim 40 \text{ s}^{-1}$; Table 1; (32)). This suggests that Asp-139 obtains protons from solvent even in the presence of subunit III, albeit at a much slower rate.

Similar to N139D, the mutant N139C is highly active ((41,42); Table 1), indicating that a cysteine at position 139 also permits rapid proton flow through the D pathway. We created the double mutant N139C-D132N to determine if Cys-139, like Asp-139, is capable of functioning as the sole initial proton gate for the D pathway, i.e. without Asp-132. As for Asp-139, computer modeling indicates that the thiol of a cysteine at position 139 is well placed to transfer protons from water to water given the longer hydrogen bonds of a thiol (Fig. 2). The thiol of Cys-139 is 3.5 Å from the “upstream” water and 3.4 Å from the “downstream” water. This is confirmed by a nearly identical position of the thiol group in a high resolution structure of *R. sphaeroides* N139C (Zhu and Gennis, unpublished). Similar to N139D-D132N, the activity of N139C-D132N is more rapid than D132N even with

subunit III present ($\sim 250 \text{ s}^{-1}$ vs $\sim 40 \text{ s}^{-1}$; Table 1), indicating that Cys-139 stimulates D pathway proton uptake. Upon the removal of subunit III, the activity of the double mutant increases to $\sim 1600 \text{ s}^{-1}$, indicating enhanced access of solvent protons to Cys-139. The TN_{max} of N139C-D132N III (-) is only slightly lower than that of WT III (-) (Table 1), showing that Cys-139 functions as an efficient initial proton gate in the absence of Asp-132.

The structures of CcO indicate that the most probable pathway for proton transfer from solvent to Asp-139 or Cys-139 is a short series of hydrogen bonded waters within the normal D pathway channel (Fig. 2). Subunit III appears to inhibit the formation of this water chain. If this is so, then creating more space for waters to enter this channel should enhance proton uptake even in the presence of subunit III. To test this, Asn-132 of N139D-D132N and N139C-D132N was altered to alanine, which replaces the amide group at the mouth of the channel with the smaller methyl group. Indeed, N139D-D132A has significantly greater activity *with subunit III present* than does N139D-D132N (1017 s^{-1} vs. 328 s^{-1} ; Table 1). Thus, creating more space at the mouth of the D channel increases the rate of proton transfer to Asp-139, most probably by making it more possible for waters to enter and form a hydrogen-bonded path for proton transfer from solvent. A similar result was obtained using N139C-D132A III (+) (data not shown).

In the mutant N139D, two carboxylic groups are present in the initial region of the D pathway, Asp-132 and Asp-139. The situation is similar in N139C, except that the smaller thiol group is present at position 139. With subunit III present, the steady-state activities of both mutants are roughly similar to our preparations of wild-type CcO; N139D is slightly, but consistently, more active than normal CcO while N139C is slightly less active (Table 1). The removal of subunit III stimulates the steady-state O_2 reduction activity of both mutant CcOs to rates *greater* than that of normal CcO, at least up to pH 8 (Table 1; Fig. 4). The result requires the conclusion that the altered D pathways of N139D III (-) and N139C III (-) take up substrate protons more rapidly than wild-type CcO that contains subunit III.

Arachidonic acid enhances the D pathway activity of the Asn-139 mutants

The addition of long-chain, unsaturated fatty acids, notably arachidonic acid (Aa), restores activity to CcO forms lacking Asp-132 (33), especially in the absence of subunit III (20,25,27). Arachidonic acid also enhances the activity of WT III (-), which contains Asp-132, but if subunit III is present Aa has no effect (25). Single turnover experiments confirm that Aa increases the activity of these CcO forms by increasing the rate of proton uptake into their D pathways (20). Methyl arachidonate has no effect (33,43) (Fig. 3), indicating that the carboxyl group is required for the stimulation of proton uptake. Apparently Aa binds on the surface of CcO in such a way that its carboxyl group transfers protons into the D pathway in the absence of Asp-132, or, in the presence of Asp-132, it provides an additional entry point for protons. The ability of the fatty acid carboxyl to transfer protons into the D pathway is enhanced by removing subunit III. Arachidonic acid can also stimulate proton uptake by the K pathway of *R. sphaeroides* CcO, but only when the surface exposed proton acceptor of the K pathway, Glu-101 of subunit II, is replaced by a residue that cannot be protonated (43). Therefore, the K pathways of the CcO forms presented here will not be affected by Aa. Figure 3 shows that Aa enhances the activity of D132N III (-), as it does for D132A III (-) (25). The addition of Aa also stimulates the activities of the double mutants containing Asp-139 or Cys-139 as initial proton gates, similar to the effect of Aa on WT III (-) (Fig. 3). Methyl arachidonate has no effect. Based on the background work described above, we conclude that Aa increases the rate of proton uptake into the D pathways of these CcOs by restoring an initial proton gate in D132N III (-) and by providing an additional proton entry point in N139D/C-D132N III (-). The data also indicate that proton uptake into the D pathway is a rate-limiting process for steady-state O_2 reduction by these CcO forms, as suggested by previous studies (20,25,44). Interestingly, Aa

also stimulates the activities of N139C III (-) and N139D III (-) (Fig. 3), even though these CcO forms already take up substrate protons at a greater rate than normal CcO.

Estimating the functional pK_a values of the initial proton acceptors of the Asn-139 mutants

In comparison to normal CcO, subunit III-depleted CcO exhibits significantly slower proton uptake into the D pathway at pH values greater than ~ 6.5 (21,26) (Fig. 4). Analyses of single turnover experiments, during which only D pathway protons are taken up, support the argument that the shift in the pH dependence of proton uptake occurs with a decrease in the apparent pK_a of Asp-132 to ~ 7 (20,26). This is consistent with the structure of CcO, since the removal of subunit III will increase the exposure of Asp-132 to solvent water and thereby confer greater stability to the anionic carboxyl. When proton uptake into the D pathway has been made rate-limiting, the pH dependence of steady-state O_2 reduction can be used to estimate the pK_a of the initial proton acceptor. (20). For example, the pH dependence of steady-state O_2 reduction by WT III (-) yields a pK_a of 7.0-7.3 (Fig. 4; Table 2; (21,25)), similar to that obtained from the pH dependence of proton uptake during a single turnover (20,26). The pK_a values obtained for Asp-132 and other initial proton gates from steady-state activity data are not 'true' pK_a values since they are not derived from direct, equilibrium measurements of the protonation state of these residues. Rather, these are 'apparent' or 'functional' pK_a values that characterize the group(s) whose protonation and deprotonation controls proton uptake into the D pathway during continuous catalytic turnover.

For N139D-D132N III (-), a pK_a of 7.7 is obtained from the pH dependence of steady-state activity, ~ 0.5 pH units higher than that of WT III (-) (Fig. 4; Table 2). Similar to WT III (-), this value should reflect the functional pK_a of Asp-139 (sans subunit III) if the rate of CcO activity is limited by the rate at which protons enter the D pathway via Asp-139. This assumption is supported by the close similarity of N139D-D132N III (-) to WT III (-). For both CcO forms, an altered D pathway leads to O_2 reduction activity that is slower than WT III (+) over a wide pH range (Fig. 4). Furthermore, the activity of both N139D-D132N III (-) and WT III (-) is further enhanced by arachidonic acid (Fig. 3), indicating that the transfer of solvent protons into the D pathway limits the rate of O_2 reduction, as discussed above. The same arguments hold true for N139C-D132N III (-), where a functional pK_a of 8.0 is assigned to Cys-139 (Fig. 4; Table 2).

For N139D III (-) and N139C III (-), the pH dependencies show their activities to be greater than normal CcO containing subunit III, up to approximately pH 8 (Fig. 4). The further stimulation of their O_2 reduction activities by Aa (Fig. 3) suggests that proton uptake into the D pathway of these CcO forms may still be a limiting factor for active site chemistry when Aa is not present, regardless of the fact that their O_2 reduction activity is greater than normal. The pH dependencies of steady-state O_2 reduction activity yield functional pK_a values of 8.2 for N139C III (-) and 8.3 for N139D III (-). With at least two potential proton acceptors in each of these mutants, Asp-132 and Asp-139 in N139D III (-) and Asp-132 and Cys-139 in N139C III (-), the assignment of functional pK_a values to a specific residue is not possible.

Activity of the subunit III-depleted Asn-139 mutants in lipid vesicles

Once purified CcO is incorporated into phospholipid vesicles (COVs), proton consumption from the interior of the COVs raises the pH near the entry region of the D pathway. At the same time, charge separation generates a membrane potential, positive outside. Both conditions slow proton uptake from the COV interior via the D and K pathways and lower the O_2 reduction activity of CcO. Under controlled conditions, the rate of controlled O_2 reduction by WT III (-) in COVs is slow, $\sim 15 \text{ s}^{-1}$ (Table 3). In contrast, the controlled

activities of the subunit III-depleted Asp-139 mutants are two to four-fold higher than WT III (-) (Table 3). The high RCR values (Table 3) indicate that the increased activity cannot be ascribed to proton leakage through the vesicle membrane. The controlled activities of the subunit III-depleted Asp-139 mutants in COVs increase in the same order as the functional pK_a values obtained for these CcO forms in detergent solution (Table 2). This indicates that controlled CcO activity by these forms is determined by the activity of D pathway which, in turn, is controlled by the proton affinities of the initial proton acceptors of the different D pathways. The controlled activity of D132N III (-), which lacks an initial proton acceptor, is even lower than WT III (-).

The activities of the uncontrolled COVs of the Asp-139 mutants at pH 7.4 are similar to the detergent-solubilized forms (Table 3), indicating that the lipid environment does not alter the characteristics of the mutant CcOs.

Discussion

In the absence of subunit III and Asp-132, Asp-139 within the D pathway becomes the initial anionic proton acceptor

The normal initial proton acceptor of the D pathway is the carboxyl group of Asp-132 of subunit I (2). When Asp-132 is altered to asparagine, alanine or histidine, proton uptake by the D pathway is essentially eliminated (19,20). However, ~20% of the normal rate of steady-state proton uptake is restored by removing subunit III (20,21,25), apparently because this modification creates space for a short chain of waters to transfer protons from solvent to the first crystal water of the D pathway, immediately below Asn-139 (Fig. 1). Because the amide side chain of Asn-139 of normal CcO is argued to adopt a conformation that allows the formation of a continuous chain of waters between Asp-132 and Glu-286 during proton transfer (13), the D pathways for proton transfer in D132N/A/H III (-) likely consist of a chain of waters all the way from solvent to Glu-286. The relatively low efficiency of proton uptake measured for D132A/H III (-) ($\sim 400\text{s}^{-1}$; (20)) may reflect the lack of a protein group to attract and transfer solvent protons into the D pathways.

The mutant N139D-D132N also lacks Asp-132 near the surface but it contains a carboxyl group at position 139, $\sim 6\text{\AA}$ further into the D pathway, in the interior of subunit I (Fig. 1). Similar to D132N/A/H, the removal of subunit III increases the activity of N139D-D132N, but to a much greater extent (Table 1; Fig. 4). In fact, the TN_{max} value of subunit III-depleted N139D-D132N is similar to that of WT III (-), which retains Asp-132 as the initial proton acceptor of the D pathway.

It seems most likely that Asp-139 plays a direct role in proton transfer, rather than moving aside, as proposed for Asn-139 of normal CcO (13), or by causing a structural disruption that allows the removal of subunit III to create a continuous pathway of water from solvent to Glu-286. The reasons for this conclusion, which also apply to Cys-139, are these. 1) The modeled and actual structures show that Asp-139 does not distort CcO structure and that its carboxyl is positioned to participate in proton transfer (Fig. 2; see Results). (The actual structures are, in fact, obtained in the absence of subunit III.) 2) The structures show no obvious pathways bypassing Asp-139. 3) If the functional D pathway of N139D-D132N III (-) did not include Asp-139, i.e. if it consisted solely of water from solvent to Glu-286, its characteristics should be similar to those described above for D132N/A/H III (-). However, the D pathway of N139D-D132N III (-) is at least four-fold more active than those of D132A/H/N III (-) (steady-state proton uptake of 1800s^{-1} vs $\sim 400\text{s}^{-1}$; Table 1). 4) The functional pK_a value of proton uptake facilitated by Asp-139 (7.7; Table 2) is consistent with an internal carboxyl group and considerably different from the high pK_a measured for

proton uptake by D132A/H III (-) (>10; (20)). We conclude that Asp-139 participates directly in proton transfer.

A separate consideration is whether or not Asp-139 of N139D-D132N III (-) protonates and deprotonates during proton transfer. Given its position, the carboxyl of Asp-139 could remain neutral by participating in proton transfer as a Grotthus-type element (10,45), that is by acquiring a proton from a hydronium 'below' its side chain while simultaneously releasing a proton to the water 'above'. However, the results are more consistent with Asp-139 cycling between its protonated and deprotonated forms. The higher rate of proton uptake supported by Asp-139 is consistent with the presence of an anion at position 139 that attracts and 'traps' protons in the same manner as Asp-132. In contrast, uncharged Asp-139 would not be expected to enhance the rate of proton uptake in comparison to D132A/H/N III (-). In addition, we observe a further increase in the functional pK_a of proton uptake for N139D III (-) due to electrostatic interaction between Asp-139 and Asp-132 (see discussion below). Obviously, this requires that Asp-139 can ionize. Finally, FTIR analysis suggests that the equivalent of Asp-139 in a mutant of the *aa*₃-type CcO of *P. denitrificans* is deprotonated at pH 7 (41). From the sum of these results we conclude that Asp-139 can function as an efficient initial proton acceptor in lieu of Asp-132, once subunit III is removed, and that Asp-139 ionizes during proton transfer.

Cysteine as an initial proton acceptor for the D pathway

Cysteine also provides a side chain capable of protonation and deprotonation in the physiologic pH range. Like Asp-139, the CcO structures show that Cys-139 is well positioned to transfer protons (see Results). The high activity of N139C CcO ((41,42); Table 1) shows that Cys-139 does not interrupt the flow of protons to the site of O₂ reduction. N139C-D132N was created to place a thiol group at position 139 in the absence of Asp-132 (Fig. 2). Similar to N139D-D132N, the removal of subunit III from N139C-D132N increases its TN_{max} to a value similar to that of WT III (-) (Table 1). Cys-139 of N139C-D132N III (-) participates directly in proton transfer and it appears to ionize, based on evidence similar to that discussed above for Asp-139. We conclude that Cys-139 functions as the initial proton acceptor for the D pathway of N139C-D132N III (-). To our knowledge, such a role for cysteine is not found in naturally occurring proton transfer pathways. The success of cysteine-139 verifies the general utility of anionic groups as initial proton acceptors.

Proton transfer from solvent to Asp/Cys-139

Neither Asp-139 nor Cys-139 is exposed to bulk solvent, even in the absence of subunit III. The most likely pathway to transfer protons from solvent to Asp/Cys-139 is a short series of waters within the initial region of the normal D pathway channel. Since protons move rapidly along short chains of hydrogen-bonded waters (10), a water file would allow facile proton transfer from solvent to Asp-139 or Cys-139. Moreover, simulations show that carboxylate side chains organize short water files that facilitate proton transfer to the carboxyl group (46). With subunit III removed, a slight outward movement (~1 Å) of the lower region of helix 3 of subunit I would create space sufficient for the formation of a water file leading from bulk solvent to the water molecule normally observed between the carboxyl group of Asp-132 and the amide side chain of Asn-139 (14,17,28) (Fig. 2). The lower region of helix 3 of subunit I participates directly in the subunit I-III interface, so the removal of subunit III should allow increased flexibility. Indeed, a molecular dynamics simulation of an *R. sphaeroides* CcO form lacking subunit III shows the spontaneous movement of three solvent water molecules into the initial region of the D pathway (13). Experimentally, the replacement of the amide group at position 132 with the smaller methyl group increases the rate of steady-state proton transfer to Asp-139 three-fold even when

subunit III is present (N139D-D132A III (+) vs N139D-D132N III (+); Table 1) This result strongly supports the proposal of a water file to position 139 within the existing D pathway channel (Fig 2).

The finding that Asp-139 and Cys-139 facilitate rapid steady-state proton uptake demonstrates that protein groups that speed proton uptake need not be in direct contact with bulk solvent. This has relevance for the situation of Asp-132 of normal CcO. With subunit III present, Asp-132 has limited accessibility to solvent water because it is located at the bottom of a narrow cavity some 6 to 10 Å deep (22,24,28). As such, it seems likely that the protonation of anionic Asp-132 involves proton transfer from solvent via a short series of hydrogen-bonded waters.

Multiple proton acceptors enhance steady-state proton uptake

The activities of the subunit III-depleted CcO forms containing Asp-132 *plus* Asp-139 or Cys139 are actually *greater* than normal CcO containing subunit III, and this higher activity is sustained at lower pH (Fig 4). Electrostatic interactions cannot account for the greater activity of these CcO forms at low pH since the functional pK_a values show that Asp-132, Asp-139 and Cys-139 will lose charge. It could be argued that the Asp-132/Asp-139 pair or the Asp-132/Cys-139 pair provide tandem side chains for the storage of two protons at the entry of the D pathway, which could speed proton uptake and hence activity. However, this arrangement is also present in N139D/C III (+) and therefore it does not explain the increased steady-state activity of these forms upon the removal of subunit III.

A reasonable explanation for the higher than normal activities of N139D/C III (-) arises from our findings that Asp-132, Asp-139 and Cys-139 can all function, independently, as efficient proton entry points. Therefore, upon the removal of subunit III from N139D/C, two pathways deliver protons to Asp/Cys-139 for transfer into the rest of the D pathway. In one path, protons flow from solvent to Asp-132 and onto Asp/Cys-139 via one intervening water (14). In a second path, protons from solvent bypass Asp-132 and flow to Asp/Cys-139 by the water file shown in Fig. 2. The delivery of solvent protons to Asp/Cys-139 by two pathways speeds proton uptake and CcO activity. The creation of two proton entry points in N139D III (-) appears similar to the stimulation of the activity of WT III (-) by arachidonic acid (20,25). In both cases the addition of a second carboxyl group increases the rate of proton uptake.

The situation is further complicated by the observation that the addition of arachidonic acid (but not its methylated form) to N139D III (-) and N139C III (-) further increases their rates of steady-state proton uptake. The result raises the possibility that three proton acceptors, Asp-132, Asp/Cys-139 and the carboxyl of Aa can simultaneously deliver protons into the D pathway in the absence of subunit III.

The normal K pathway exceeds the proton uptake requirements of wild-type CcO

During each catalytic cycle the D pathway takes up two (possibly three) of the four substrate protons required for O₂ reduction and the K pathway takes up the remaining one or two (2). While the D and K pathways converge near the active site, their proton entry points on the inner surface of CcO are far apart (Fig. 1). There is no experimental or structural evidence that the alteration of residues in the initial regions of either pathway affects the other. The greater than normal O₂ reduction rates of N139D/C III (-) requires greater than normal rates of proton uptake by both their D and K pathways. Since the K pathways in these mutant CcOs are not modified, it follows that the K pathway of normal CcO imposes no limitation on the rate of O₂ reduction. The same argument holds true for CcO in lipid vesicles, where proton uptake into both the K and D pathways is inhibited by a transmembrane voltage

gradient and high pH in the interior of the vesicles. Thus, across the range of possible pH values and voltage gradient conditions encountered by wild-type CcO it appears that the proton uptake capability of the K pathway exceeds the requirements for normal activity. In contrast, proton uptake into the D pathway may be a rate-limiting process.

The initial proton acceptor of the D pathway controls CcO activity

The ability of CcO to function at physiological pH is considerably compromised by the removal of subunit III. For example, the activity of WT III (-) is only 20% that of normal CcO at pH 8, while at pH 6.5 the activities of the two forms are similar (Fig. 4). The primary source of this difference appears to be a decline in the functional pK_a of Asp-132 due to greater exposure of Asp-132 to solvent in the absence of subunit III (20).

With this study, we demonstrate that even in the absence of subunit III the pH range of CcO activity can be extended to match that of normal CcO by manipulating the environment and the identity of the initial proton acceptor of the D pathway. In N139D-D132N III (-), the initial aspartate is moved to a position inside the D pathway that eliminates its exposure to bulk solvent. This change increases the proton affinity of the pathway approximately three-fold in comparison to that of WT III (-) (Table 2; Fig. 4) where the carboxyl of Asp-132 is partially exposed to solvent. Hence, shielding the initial proton acceptor of the D pathway from bulk solvent can extend the pH range of the enzyme without lowering its overall activity. This strategy appears to be used by normal CcO, where a ring of residues from subunits I and III shield the carboxyl of Asp-132 from bulk solvent.

The pH dependence of CcO activity is extended another 0.3 pH units with the introduction of Cys-139 as the initial proton acceptor, presumably due to the greater inherent proton affinity of the thiol group for proton. In fact, a greater pH shift might be expected with a thiol as the proton acceptor since continuum electrostatics calculations predict a higher pK_a for Cys-139 in the resting enzyme (41). However, the functional pK_a value measured here reflects the situation during continuous turnover where the apparent proton affinity of the initial acceptor is also influenced by the rate at which protons are withdrawn to the active site. This may prevent large increases in the functional pK_a value.

When Asp-132 is present along with either Asp-139 or Cys-139, the pH profiles become similar to that of normal CcO containing subunit III (Table 2; Fig. 4). Electrostatic interaction between the deprotonated forms of Asp-132 and Asp-139 or Cys-139 should raise the proton affinities of these groups, providing the likely explanation for the retention of activity at higher pH.

Conclusions

The examination of CcO mutants with an aspartate or a cysteine inside the D pathway for proton transfer reveals new information about the process of proton uptake.

1. In the absence of subunit III, protons from solvent water gain access to Asp-139 in the interior of the D pathway such that Asp-139 becomes as efficient an initial proton acceptor for steady-state activity as surface-exposed Asp-132. Solvent protons are likely transferred to Asp-139 through a short file of waters. In normal CcO, subunit III provides a barrier for the influx of solvent water and the formation of an alternative proton pathway that can bypass Asp-132.
2. A protein group that speeds the uptake of protons from solvent need not be in direct contact with bulk solvent. In these experiments, both a carboxylate group and a thiolate group speed proton uptake from an internal position. It is likely that their anionic nature is key to their function in that the negative charge will both attract

protons and facilitate the formation of the hydrogen-bonded water file from solvent (46).

3. Proton uptake can be driven to higher rates by placing multiple anionic proton acceptors in the initial region of the D pathway.
4. Proton uptake into the K pathway is not rate limiting for CcO activity at physiologic pH.
5. The pH range of CcO activity can be manipulated by altering the proton affinity of the initial proton acceptor for the D pathway. The experiments here show that this can be achieved by restricting the access of the acceptor to bulk solvent, by changing the chemical identity of the acceptor and by the electrostatic interaction of the acceptor with another group. Normal CcO appears to use the first of these strategies to maintain a high functional pK_a for Asp-132.

Acknowledgments

The authors thank Professor R.B. Gennis of the University of Illinois for sharing strains and plasmids encoding mutant CcO forms N139D, D132N, N139C and N139D-D132N. We also thank the Gennis group, particularly Dr. Jiapeng Zhu, for sharing high resolution structures of *R. sphaeroides* CcO forms N139D and N139C.

Supported by National Institutes of Health Grant GM56824

References

1. Richter OM, Ludwig B. Cytochrome *c* oxidase--structure, function, and physiology of a redox-driven molecular machine. *Rev Physiol Biochem Pharmacol*. 2003; 147:47–74. [PubMed: 12783267]
2. Hosler JP, Ferguson-Miller S, Mills DA. Energy transduction: Proton transfer through the respiratory complexes. *Annu Rev Biochem*. 2006; 75:165–187. [PubMed: 16756489]
3. Brzezinski P, Gennis RB. Cytochrome *c* oxidase: exciting progress and remaining mysteries. *J Bioenerg Biomembr*. 2008; 40:521–531. [PubMed: 18975062]
4. Brzezinski P, Ädelroth P. Design principles of proton-pumping haem-copper oxidases. *Curr Opin Struct Biol*. 2006; 16:465–472. [PubMed: 16842995]
5. Wikström M, Verkhovsky MI. Mechanism and energetics of proton translocation by the respiratory heme-copper oxidases. *Biochim Biophys Acta*. 2007; 1767:1200–1214. [PubMed: 17689487]
6. Sharpe MA, Ferguson-Miller S. A chemically explicit model for the mechanism of proton pumping in heme-copper oxidases. *J Bioenerg Biomembr*. 2008; 40:541–549. [PubMed: 18830692]
7. Hosler JP, Ferguson-Miller S, Calhoun MW, Thomas JW, Hill J, Lemieux L, Ma J, Georgiou C, Fetter J, Shapleigh J, Tecklenburg MJ, Babcock GT, Gennis RB. Insight into the active-site structure and function of cytochrome oxidase by analysis of site-directed mutants of bacterial cytochrome *aa₃* and cytochrome *bo*. *J Bioenerg Biomembr*. 1993; 25:121–136. [PubMed: 8389745]
8. Richter OM, Ludwig B. Electron transfer and energy transduction in the terminal part of the respiratory chain - lessons from bacterial model systems. *Biochim Biophys Acta*. 2009; 1787:626–634. [PubMed: 19268423]
9. Sharpe, M.; Qin, L.; Ferguson-Miller, S. Proton Entry, Exit and Pathways in Cytochrome Oxidase: Insight from “Conserved” Water. In: Wikström, M., editor. *Biophysical and Structural Aspects of Bioenergetics*. RSC Publishing; Cambridge UK: 2005. p. 26-54.
10. Wraight CA. Chance and design--proton transfer in water, channels and bioenergetic proteins. *Biochim Biophys Acta*. 2006; 1757:886–912. [PubMed: 16934216]
11. Wikström M, Jasaitis A, Backgren C, Puustinen A, Verkhovsky MI. The role of the D- and K-pathways of proton transfer in the function of the haem-copper oxidases. *Biochim Biophys Acta*. 2000; 1459:514–520. [PubMed: 11004470]
12. Brzezinski P, Ädelroth P. Pathways of proton transfer in cytochrome *c* oxidase. *Journal of Bioenergetics & Biomembranes*. 1998; 30:99–107. [PubMed: 9623811]

13. Henry RM, Yu CH, Rodinger T, Pomes R. Functional hydration and conformational gating of proton uptake in cytochrome *c* oxidase. *J Mol Biol.* 2009; 387:1165–1185. [PubMed: 19248790]
14. Koepke J, Olkhova E, Angerer H, Müller H, Peng G, Michel H. High resolution crystal structure of *Paracoccus denitrificans* cytochrome *c* oxidase: new insights into the active site and the proton transfer pathways. *Biochim Biophys Acta.* 2009; 1787:635–645. [PubMed: 19374884]
15. Kaila VR, Sharma V, Wikström M. The identity of the transient proton loading site of the proton-pumping mechanism of cytochrome *c* oxidase. *Biochim Biophys Acta.* 1807:80–84. [PubMed: 20831859]
16. Wikström M, Verkhovsky MI, Hummer G. Water-gated mechanism of proton translocation by cytochrome *c* oxidase. *Biochim Biophys Acta.* 2003; 1604:61–65. [PubMed: 12765763]
17. Qin L, Liu J, Mills DA, Proshlyakov DA, Hiser C, Ferguson-Miller S. Redox-dependent conformational changes in cytochrome *c* oxidase suggest a gating mechanism for proton uptake. *Biochemistry.* 2009; 48:5121–5130. [PubMed: 19397279]
18. Tomson FL, Morgan JE, Gu G, Barquera B, Vygodina TV, Gennis RB. Substitutions for glutamate 101 in subunit II of cytochrome *c* oxidase from *Rhodobacter sphaeroides* result in blocking the proton-conducting K-channel. *Biochemistry.* 2003; 42:1711–1717. [PubMed: 12578386]
19. Smirnova IA, Ädelroth P, Gennis RB, Brzezinski P. Aspartate-132 in Cytochrome *c* Oxidase from *Rhodobacter sphaeroides* is Involved in a Two-Step Proton Transfer during Oxo-Ferryl Formation. *Biochemistry.* 1999; 38:6826–6833. [PubMed: 10346904]
20. Ädelroth P, Hosler J. Surface proton donors for the D-pathway of cytochrome *c* oxidase in the absence of subunit III. *Biochemistry.* 2006; 45:8308–8318. [PubMed: 16819830]
21. Hosler JP. The influence of subunit III of cytochrome *c* oxidase on the D pathway, the proton exit pathway and mechanism-based inactivation in subunit I. *Biochim Biophys Acta.* 2004; 1655:332–339. [PubMed: 15100048]
22. Svensson-Ek M, Abramson J, Larsson G, Törnroth S, Brzezinski P, Iwata S. The X-ray crystal structures of wild-type and EQ(I-286) mutant cytochrome *c* oxidases from *Rhodobacter sphaeroides*. *J Mol Biol.* 2002; 321:329–339. [PubMed: 12144789]
23. Yoshikawa S, Shinzawa-Itoh K, Tsukihara T. Crystal structure of bovine heart cytochrome *c* oxidase at 2.8 Å resolution. *J Bioenerg Biomembr.* 1998; 30:7–14. [PubMed: 9623800]
24. Harrenga A, Michel H. The cytochrome *c* oxidase from *Paracoccus denitrificans* does not change the metal center ligation upon reduction. *J Biol Chem.* 1999; 274:33296–33299. [PubMed: 10559205]
25. Mills DA, Tan Z, Ferguson-Miller S, Hosler J. A role for subunit III in proton uptake into the D pathway and a possible proton exit pathway in *Rhodobacter sphaeroides* cytochrome *c* oxidase. *Biochemistry.* 2003; 42:7410–7417. [PubMed: 12809496]
26. Gilderson G, Salomonsson L, Aagaard A, Gray J, Brzezinski P, Hosler J. Subunit III of cytochrome *c* oxidase of *Rhodobacter sphaeroides* is required to maintain rapid proton uptake through the D pathway at physiologic pH. *Biochemistry.* 2003; 42:7400–7409. [PubMed: 12809495]
27. Han D, Morgan JE, Gennis RB. G204D, a mutation that blocks the proton-conducting D-channel of the *aa₃*-type cytochrome *c* oxidase from *Rhodobacter sphaeroides*. *Biochemistry.* 2005; 44:12767–12774. [PubMed: 16171391]
28. Muramoto K, Hirata K, Shinzawa-Itoh K, Yoko-o S, Yamashita E, Aoyama H, Tsukihara T, Yoshikawa S. A histidine residue acting as a controlling site for dioxygen reduction and proton pumping by cytochrome *c* oxidase. *Proc Natl Acad Sci USA.* 2007; 104:7881–7886. [PubMed: 17470809]
29. Pfützner U, Hoffmeier K, Harrenga A, Kannt A, Michel H, Bamberg E, Richter OM, Ludwig B. Tracing the D-pathway in reconstituted site-directed mutants of cytochrome *c* oxidase from *Paracoccus denitrificans*. *Biochemistry.* 2000; 39:6756–6762. [PubMed: 10841754]
30. Pawate AS, Morgan J, Namslauer A, Mills D, Brzezinski P, Ferguson-Miller S, Gennis RB. A mutation in subunit I of cytochrome oxidase from *Rhodobacter sphaeroides* results in an increase in steady-state activity but completely eliminates proton pumping. *Biochemistry.* 2002; 41:13417–13423. [PubMed: 12416987]

31. Namslauer A, Pawate AS, Gennis RB, Brzezinski P. Redox-coupled proton translocation in biological systems: proton shuttling in cytochrome *c* oxidase. *Proc Natl Acad Sci USA*. 2003; 100:15543–15547. [PubMed: 14676323]
32. Brändén G, Pawate AS, Gennis RB, Brzezinski P. Controlled uncoupling and recoupling of proton pumping in cytochrome *c* oxidase. *Proc Natl Acad Sci USA*. 2006; 103:317–322. [PubMed: 16407159]
33. Fetter J, Sharpe M, Qian J, Mills D, Ferguson-Miller S, Nicholls P. Fatty acids stimulate activity and restore respiratory control in a proton channel mutant of cytochrome *c* oxidase. *FEBS Lett*. 1996; 393:155–160. [PubMed: 8814281]
34. Sone N, Yoshida M, Hirata H, Kagawa Y. Asolectin purification procedure. *J Biochem (Tokyo)*. 1977; 81:519–528. [PubMed: 14954]
35. Keen NT, Tamaki S, Kobayashi D, Trollinger D. Improved Broad-Host-Range Plasmids for DNA Cloning in Gram-Negative Bacteria. *Gene*. 1988; 70:191–197. [PubMed: 2853689]
36. Shapleigh JP, Gennis RB. Cloning, sequencing, and deletion from the chromosome of the gene encoding subunit I of the *aa3*-type cytochrome *c* oxidase of *Rhodobacter sphaeroides*. *Mol Microbiol*. 1992; 6:635–642. [PubMed: 1313140]
37. Zhen Y, Qian J, Follmann K, Hayward T, Nilsson T, Dahn M, Hilmi Y, Hamer AG, Hosler JP, Ferguson-Miller S. Overexpression and purification of cytochrome *c* oxidase from *Rhodobacter sphaeroides*. *Prot Express Purif*. 1998; 13:326–336.
38. Fetter JR, Qian J, Shapleigh J, Thomas JW, Garcia-Horsman A, Schmidt E, Hosler J, Babcock GT, Gennis RB, Ferguson-Miller S. Possible proton relay pathways in cytochrome *c* oxidase. *Proc Natl Acad Sci USA*. 1995; 92:1604–1608. [PubMed: 7878026]
39. Qin L, Hiser C, Mulichak A, Garavito RM, Ferguson-Miller S. Identification of conserved lipid/detergent-binding sites in a high-resolution structure of the membrane protein cytochrome *c* oxidase. *Proc Natl Acad Sci USA*. 2006; 103:16117–16122. [PubMed: 17050688]
40. Guex N, Peitsch MC. SWISS-MODEL and the Swiss-PdbViewer: an environment for comparative protein modeling. *Electrophoresis*. 1997; 18:2714–2723. [PubMed: 9504803]
41. Dürr KL, Koepke J, Hellwig P, Müller H, Angerer H, Peng G, Olkhova E, Richter OM, Ludwig B, Michel H. A D-pathway mutation decouples the *Paracoccus denitrificans* cytochrome *c* oxidase by altering the side-chain orientation of a distant conserved glutamate. *J Mol Biol*. 2008; 384:865–877. [PubMed: 18930738]
42. Zhu J, Han H, Pawate A, Gennis RB. Decoupling mutations in the D-channel of the *aa3*-type cytochrome *c* oxidase from *Rhodobacter sphaeroides* suggest that a continuous hydrogen-bonded chain of waters is essential for proton pumping. *Biochemistry*. 2010; 49:4476–4482. [PubMed: 20441187]
43. Qin L, Mills DA, Hiser C, Murphree A, Garavito RM, Ferguson-Miller S, Hosler J. Crystallographic location and mutational analysis of Zn and Cd inhibitory sites and role of lipidic carboxylates in rescuing proton path mutants in cytochrome *c* oxidase. *Biochemistry*. 2007; 46:6239–6248. [PubMed: 17477548]
44. Mills DA, Hosler JP. Slow proton transfer through the pathways for pumped protons in cytochrome *c* oxidase induces suicide inactivation of the enzyme. *Biochemistry*. 2005; 44:4656–4566. [PubMed: 15779892]
45. Nagle JF, Tristram-Nagle S. Hydrogen bonded chain mechanisms for proton conduction and proton pumping. *J Membr Biol*. 1983; 74:1–14. [PubMed: 6306243]
46. Gu W, Helms V. Tightly connected water wires facilitate fast proton uptake at the proton entrance of proton pumping proteins. *J Am Chem Soc*. 2009; 131:2080–2081. [PubMed: 19159297]

Abbreviations

Aa	arachidonic acid
CcO	cytochrome <i>c</i> oxidase
COV	cytochrome <i>c</i> oxidase vesicle

CCCP	carbonyl cyanide 3-chloro-phenylhydrazone
Ni-NTA	Ni ²⁺ -nitriloacetic acid agarose
RCR	respiratory control ratio
TMPD	N,N,N',N'-tetramethyl- <i>p</i> -phenylenediamine
TN	turnover number as e ⁻ or H ⁺ s ⁻¹
WT III (+)	wild-type CcO
WT III (-)	wild-type CcO lacking subunit III

Figure 1.

The D and K pathways of the *aa*₃-type cytochrome *c* oxidase of *R. sphaeroides*. Subunits I, II and III are shadowed in cyan, green and tan respectively. The waters and selected residues of the D pathway are shown along the blue arrow, and selected residues of the K pathway along the red arrow.

Figure 2.

The normal D pathway and variants containing aspartic acid or cysteine at position 139. The normal D pathway (A) is that of the structure of the Subunit I-II CcO of *R. sphaeroides* at 2.0 Å resolution (2GSM; (39)). The structures shown for N139D (B) and N139C (C) were modeled from the 2GSM structure using the DeepView/Swiss-PdbViewer software package (40). It is assumed that the alteration of Asp-132 to Asn in N139D/C-D132N does not significantly alter the position of Asp-139 or Cys-139. Therefore, models of these CcO forms are not shown. In panel D, three waters (gold; 2.6 Å apart) have been added to the 2GSM structure using DeepView to show one possibility of a water file sufficient to bypass Asp-132 or Asn-132 and transfer protons from solvent to the first crystal water (blue) shown between Asp-132 and Asn-139 in panel A. In the model of panel D, the lower two waters are both exposed to solvent. Both make steric clashes with Pro-136 and Met-133 of subunit I, which would be relieved if the lower region of helix 3 of subunit I moved less than 1 Å. Such flexibility seems likely in the absence of subunit III.

Figure 3.

The effect of 250 μ M arachidonic acid or 250 μ M methyl arachidonate on the steady-state activities of subunit III-depleted wild-type CcO, D132N and four Asn-139 mutants. Activities were measured at pH 7.5 as described in Methods. Error is standard deviation.

Figure 4.

The pH dependence of steady-state activity by wild-type CcO and subunit III-depleted CcO forms. Green open squares = WT III (-); red open circles = N139D-D132N III (-); brown open squares = N139C-D132N III (-); blue open upward triangles = WT III (+); black filled downward triangles = N139C III (-); red filled circles = N139D III (-). Activity measurements and fits to the Henderson-Hasselbalch equation are as described in Methods.

Table 1

Comparison of the $TN_{\max}$ values at pH 6.5 of wild-type CcO and Asn-139 mutants in the presence and absence of subunit III.

CcO form	III (+)	III (-)
WT	$2062^a \pm 24^b$	1835 ± 48
D132N	43 ± 2	383 ± 11
N139D-D132N	328 ± 7	1799 ± 75
N139C-D132N	251 ± 4	1587 ± 23
N139D	$2187^c \pm 81$	2505 ± 24
N139C	$1781^c \pm 36$	2256 ± 36
N139D-D132A	1017 ± 37	ND

^a e^- or H^+ consumed s^{-1}

^b standard error of the fit to the Michaelis-Menton function

^c The activities of N139D/C III (+) are in good agreement with the relative activities published for the corresponding mutants of *P. denitrificans* CcO (41) and with the absolute activities published previously for these *R. sphaeroides* CcO forms (42)

Table 2
Functional pK_a values of steady-state activity by wild-type CcO and the Asn-139 mutants

CcO form	pK_a of steady-state TN^a	Initial proton acceptor
WT III (+)	8.4 (8.3-8.5) ^b	Asp-132
WT III (-)	7.3 (7.2-7.4)	Asp-132
N139D-D132N III (-)	7.7 (7.5-7.9)	Asp-139
N139C-D132N III (-)	8.0 (7.9-8.1)	Cys-139
N139D III (-)	8.3 (8.2-8.4)	Asp-132 & Asp-139
N139C III (-)	8.2 (8.2-8.3)	Asp-132 & Cys-139

^a TN = turnover number as e^- or H^+ consumed s^{-1}

^b95% confidence interval

Table 3
Controlled and uncoupled activity of COVs containing wild-type CcO and the subunit III-depleted D pathway mutants

COVs	Controlled TN ^a	Uncoupled TN ^b	RCR ^c
WT III (-)	15.5 ± 3.5	647 ± 107	42
D132N III (-)	10.2 ± 0.9	72 ± 8	7.2
N139D-D132N III (-)	31 ± 5	730 ± 31	23.4
N139C-D132N III (-)	37 ± 5	405 ± 12	10.9
N139D III (-)	54 ± 1	1315 ± 56	25
N139C III (-)	55 ± 3	1092 ± 14	20

^aError is standard deviation; TN = turnover number as e⁻ or H⁺ consumed s⁻¹

^bIn the presence of 3 μM valinomycin and 5 μM CCCP

^cControlled TN/Uncoupled TN