Published in final edited form as:

Yeast. 2009 July ; 26(7): 399–406. doi:10.1002/yea.1674.

Additional cassettes for epitope and fluorescent fusion proteins in *Candida albicans*

Maryam Gerami-Nejad¹, Keely Dulmage¹, and Judith Berman^{2,*}

¹ Department of Genetics, Cell Biology and Development, University of Minnesota, MN, USA

² Department of Microbiology, University of Minnesota, MN, USA

Abstract

Epitope tags that confer specific properties, including affinity for resins or antibodies or detection by fluorescence microscopy, are highly useful for biochemical and cell biological investigations. In *Candida albicans* and several other related yeasts, the CUG codon specifies serine instead of leucine, requiring that molecular tools be customized for use in this important human fungal pathogen. Here we report the construction of a set of plasmids containing 13-Myc, 3HA, GST, V5 or His9 epitope cassettes that facilitate PCR-mediated construction of epitope-tagged proteins. Common primer sets amplify the different tags with two different selectable markers. In addition, we report construction of a codon-optimized *Discosoma* red fluorescent protein (DsRFP) gene. Like mCherryRFP, this DsRFP signal is detectable in transformants at the colony level and is useful in double-labelling experiments with green fluorescent protein (GFP). Finally, we describe a construct that directs PCR-mediated two-step insertion of *GFP* internal to a coding sequence, which facilitates tagging of secreted proteins, including GPI-anchor cell wall proteins that require endogenous N- and C-termini for function. These reagents expand the repertoire of molecular tools available for working with *C. albicans* and other members of the CUG clade of pathogenic yeasts.

Keywords

Candida albicans; fluorescent protein; epitope; tagging

Introduction

Candida albicans is a major cause of serious mucosal and systemic fungal infections, especially in immunocompromised patients (Pfaller and Diekema, 2007). The use of PCR-mediated transformation has enabled rapid genetic modification of this obligate diploid yeast (Wilson *et al.*, 1999). Epitope tags are important tools for performing biochemical analyses, such as affinity purification and quantification of protein levels. Especially useful are a series of cassettes that employ a defined set of common primers, so that different strains carrying epitopes inserted at the same gene can be constructed with a single pair of oligonucleotides (Bahler *et al.*, 1998; Krawchuk and Wahls, 1999; Lavoie *et al.*, 2008; Longtine *et al.*, 1998b; Sung *et al.*, 2008). The use of cassettes for tagging proteins with green, cyan and yellow fluorescent proteins (GFP, CFP and YFP) at C-termini has also

facilitated the analysis of protein localization and expression within living *C. albicans* cells (Gerami-Nejad *et al.*, 2001; Schaub *et al.*, 2006). Tagging at the N-terminus has usually involved replacement of the native promoter with a constitutive or regulatable promoter cassette (Gerami-Nejad *et al.*, 2004). Double FP labelling experiments are generally performed using YFP together with CFP. However, the development of *Discosoma* red fluorescent protein (Bevis and Glick, 2002) has facilitated the performance of double labelling experiments using GFP and RFP.

In *C. albicans*, the alternative translation of the CUG codon for serine instead of leucine obviates the use of most heterologous vectors with epitope tags. A single *C. albicans* tagging vector that expresses a histidine 6-mer (His6) C-terminal to the protein of interest has been available for a few years (Schaub *et al.*, 2006); a new set of three epitope tagging vectors (expressing HA, TAP or MYC) was recently described (Lavoie *et al.*, 2008). Here we report a series of plasmids that use a shared set of primers for the amplification of five different epitope tags. In addition, we report the construction of a synthetic, codon-optimized monomeric DsRed fluorescent protein (DsRFP) epitope that produces colonies that can be detected by their fluorescent red colour and compare its properties to those of a purple fluorescent protein recently described for use with *Saccharomyces cerevisiae* colonies (Keppler-Ross *et al.*, 2008). Finally, we describe a plasmid that permits insertion of GFP, in frame, within an open reading frame. Together, this set of epitope-tagging plasmids expands the repertoire of biochemical and cell biological experiments that can be performed in *C. albicans*.

Materials and methods

Strains, growth conditions and DNA methods

All strains used in this study (Table 2) were derived from BWP17 (Wilson *et al.*, 1999) and were grown at 30 °C in rich (YPAD) medium, synthetic complete medium (SC), or synthetic complete medium lacking specific nutrients (SM; Sherman, 1991). *Escherichia coli* strain XLI-blue (Stratagene, La Jolla, CA, USA), growth conditions, DNA manipulations and primer design and synthesis were essentially as described previously (Gerami-Nejad *et al.*, 2004). Transformants were selected on SM medium lacking uridine or histidine and were screened by PCR using oligonucleotide primers listed in Table S1.

Plasmid constructions

To generate pMG1874 (*HA-URA3*) the *Bam*H1/*Hind*III fragment of pGFP-*URA3* (Gerami-Nejad *et al.*, 2001) containing *URA3* was ligated to the *Bg*II-*Hind*III 3HA fragment of pFA6a-3HA-KanMX6 (Longtine *et al.*, 1998a). To construct pMG1921 (*HA-HIS1*), the *HIS1* marker was isolated as a *Pvu*I fragment from pGFP-*HIS1* (Gerami-Nejad *et al.*, 2001) and ligated to *Pvu*I-digested pMG1874, resulting in *HIS1* replacing the *URA3* gene. Analogous approaches were used to insert the *13-MYC* as a *Bg*II and *Hind*III fragment from pFA6a-13myc-KanMX6 (Longtine *et al.*, 1998a) into the *Bam*H1-*Hind*III fragment of pGFP-*URA3* to yield pMG1905 (*13-MYC-URA3*). To construct pMG2093 (*13MYC-HIS1*), the *HIS1* marker was isolated as a *Pvu*I fragment from pGFP-*HIS1* and was ligated into *Pvu*I-digested pMG1905.

To construct pMG2083 (*GST-URA3*) the *Bg*II-*Hind*III *GST* fragment from pFA6a-GST-TRP was ligated to *Bam*H1-*Hind*III-digested pGFP-*URA3*. To construct pMG2090 (*V5-URA3*), the V5 epitope was amplified using forward primer 1490 (designed to include a *Hind*III site; Table S2, italicized), reverse primer 1491 (designed to include a *Bam*H1 site; Table 1) and ptracer-EF/V5-HIS (Invitrogen, Carlsbad, CA, USA). The resulting PCR product was digested with *Bam* H1-*Hind*III and ligated into *Bam*H1- and *Hind*III-digested

pGFP-URA3. pMG2051 (*HIS9-URA3*) was constructed by annealing primer 1286, which includes nine CAC codons and a *HindIII* site, with primer 1287, which includes nine GTGs and a *BglII* site (Table S2), digesting the product with *BglII-HindIII* and then ligating it to *BamHI*- and *HindIII*-digested pGFP-URA3.

A synthetic *CaRFP* gene was produced at Celtek Genes (Nashville, TN, USA) based on the coding sequence of monomeric DsRed (Bevis and Glick, 2002), which was codon-optimized for *C. albicans* by changing all CUG codons to TTG, which is the most frequently used *Candida* codon for leucine (<http://www.kazusa.or.jp/codon>). The complete sequence is provided at <http://cbs.umn.edu/labs/berman/epitope.htm>. The 675 nt codon-optimized DsRed sequence was inserted into the pUC57 that had been digested with *BamHI* and *PstI* site, to generate pMG2162. Plasmid pRFP-URA (pMG2169) was constructed by digesting pMG-2162 with *BamHI* and *SalI*, and ligating the RFP coding sequence adjacent to the *URA3* fragment in pGFP-URA3 that had been digested with *BamHI* and *XhoI*.

The GFP-URA3-GFP (GUG) cassette in pMG2082 was constructed by amplifying 456 bp of the C-terminal portion of *GFP* coding sequence using primers 1402 and 1403 (Table S2) and pGFP-URA3 as template. The PCR product was digested with *XbaI* and cloned into the *XbaI* site in pYPB1-ADHp1 (Bertram *et al.*, 1996) to yield plasmid pMG2079, which contains *URA3* fused in its 3' UTR to a truncated *GFP* lacking the N-terminal 87 codons. To construct GFP-URA3-GFP, the N-terminal 459 bp of *GFP* coding sequence was amplified from pGFP-URA3 using primers 1400 and 1401 (Table S2). The PCR product was digested with *BglII* and *XhoI* and cloned into pMG2079 digested with *XhoI* and *BglII*.

To generate pMG2254 (mCherryRFP), plasmid mRFP (yEmRFP; Keppler-Ross *et al.*, 2008), kindly provided by Dr Neta Dean, was amplified using primers 3345 and 3346 (Table S2). The PCR product was cut with both *BamHI* and *EcoRV* and subsequently ligated to *BamHI* and *EcoRV*-digested pGFP-URA3.

Requests for plasmids should be sent to jber-man@umn.edu. Electronic sequence files of the tagging cassettes are available at <http://www.cbs.umn.edu/labs/berman/fluorescent.shtml> under the 'epitope tags' link.

Transformation, identification of integration events and protein detection methods

PCR was performed using one of the plasmids described above as template, and sequences homologous to the appropriate target-gene integration site sequence. Primer pairs were designed as indicated in Figure 1 and Table 1. To tag *CLB2* and *CLN1* with the V5 epitope, *CLB2* or *CLN1*-specific homologous sequence was added to the universal primer sequences described in Table 1. PCR reactions were performed as described previously (Gerami-Nejad *et al.*, 2001). The products from 10 such PCR reactions were pooled, precipitated with ethanol, resuspended in 30 μ l water and used to transform *C. albicans* strain BWP17 (Wilson *et al.*, 1999). Transformants were selected by plating on the appropriate selective media. To identify transformants in which the cassette had correctly integrated into the target gene sequence, genomic DNA was prepared and used as the template in PCR reactions, using one primer that annealed within the transformation module and a second primer that annealed to the target gene locus outside the altered regions. *CDC3* was tagged at the C-terminus with *HA*, *MYC*, *GST* or *HIS9*. *CDC3*-specific homologous sequence (70 nt) was added to the universal primer sequences described in Table 1.

Immunoblot detection of fusion proteins was performed essentially as described previously (Gerami-Nejad *et al.*, 2004), except that starter cultures were grown for ~12 h in SDC medium, diluted to OD₆₀₀ = 0.8 and grown until cultures reached OD₆₀₀ = 1.0. The cultures were collected by centrifugation and stored at -80°C until analysis. Cells were thawed and

resuspended in 50 µl Thorner buffer and heated at 100 °C for 3 min, and then vortexed with an equal volume of glass beads for 4 min at 4 °C. Prior to analysis on 10% SDS–polyacrylamide gels, samples were denatured at 100 °C for 4 min, vortexed for 2 min and lysis was confirmed by microscopy.

Proteins were transferred to a PVDF membrane, blocked for 30 min with 2% skimmed milk in TBST (20 mM Tris, pH 7.6, 137 mM NaCl, 0.1% Tween-20). Membranes were then incubated for 1 h with HRP-conjugated mouse anti-HA antibody (12CAS, 1 : 1000 in 0.2% skimmed milk in TBST), HRP-conjugated rabbit anti-GST Ab (Z-5: sc-459, 1 : 1000 in 0.2% skimmed milk in TBST), mouse Anti-c-Myc Ab (1 : 5000 in 0.2% skimmed milk in TBST) or mouse anti-V5 Ab (1 : 500 in 0.2% skimmed milk in TBST). Blots probed with non-HRP-conjugated antibodies were then washed in TBST and incubated for 1 h with horseradish peroxidase-conjugated goat anti-Mouse IgG (Santa Cruz Biotechnology, Santa Cruz, CA, USA), diluted 1 : 10 000 in 0.2% skimmed milk in TBST. The GST blot was washed in water to remove background (Wu *et al.*, 2002), all other Westerns were washed in 0.67× TBST, and proteins were then detected using Supersignal Femto Chemiluminescent reagent (Pierce, Rockford, IL, USA).

The DsRFP and mCherry variants (Shaner *et al.*, 2008), were inserted at the *ENO1* locus using primers (Table S3) that included 70 bp *ENO1*-homologous sequence plus the universal primer sequences described in Table 1.

Detection of DsRFP and of mCherryRFP fluorescence was performed on 2 day-old colonies using a Nikon SMZ1500 microscope equipped with a Chroma 59 022 filter. Six fluorescent and four non-fluorescent colonies were analysed by PCR using primers 716 and 1363 (Table S3).

Results and discussion

To produce cassettes for epitope tagging the C-terminus of *C. albicans* genes, we modified GFP-tagging vectors pGFP–URA3 and pGFP–HIS1 by replacing the *GFP* gene with sequences encoding different epitope tags. The plasmids retained the transcription termination sequence from *ADHI* and the relevant selectable marker (either *URA3* or *HIS1*). In general, to add a C-terminal epitope tag to a specific *C. albicans* gene, the cassette of interest was amplified using gene-specific forward and reverse primer pairs that included ~70 bp of homology with the gene of interest, followed by the appropriate forward (F1–F6) and reverse (R1–R3) primer sequences (Figure 1, Table 1) with homology to the tagging cassette. A useful feature of the cassettes is that the *HA*, *MYC* and *GST* epitope tags are amplified using the same forward primer (F1) and the reverse primer corresponding to the selectable marker used (R1 or R2). Similarly, the *V5*, *His9* and *DsRFP* epitopes are all amplified using the same reverse primer (R1).

To test the usefulness of the cassettes, we integrated the *HA*, *MYC*, *GST* and *His9* epitopes at the C-terminus of *CDC3* and the *V5* epitope at the C-terminus of *CLB2* and *CLN1* in laboratory strain BWP17. *CDC3* transformants were analysed by PCR for the presence of fragments, using primer 335 from within the cassette together with *CDC3*-specific primer 630 (Table S1). PCR products of the expected size were sequenced to confirm that the insertion was in-frame. For each epitope tagging transformation, PCR products of the expected sizes (Table S4) were detected in at least one and as many as three of the eight screened transformants (12.5–38% *bona fide* transformants).

To investigate whether the epitope tags were expressed at detectable levels, we separated proteins from *C. albicans* crude lysates by electrophoresis, transferred them to a PVDF membrane and probed them with commercially available antibodies that recognize the

individual epitopes (see Materials and methods). Strains expressing fusion proteins all produced a unique prominent band (Figure 2). The HA, V5 and GST fusion proteins ran at the expected electrophoretic mobility, while the Myc fusion protein ran larger than the predicted protein fusion size. This is consistent with the mobility of other Myc-tagged proteins (data not shown). Anti-His6 antibody was less specific (data not shown). Because each epitope is detected with a different antibody, it is difficult to perform a direct comparison of the signals from each epitope.

A recent report of a codon-optimized mCher-ryRFP found that the fluorescent protein was visually detectable as a purple pigment in colonies of *S. cerevisiae*, but not in *C. albicans* (Keppler-Ross *et al.*, 2008). We compared the appearance of codon-optimized DsRFP to that of mCherryRFP in *C. albicans* colonies. While colony colour was not evident by visual inspection, the *ENO1*-DsRFP fusion protein was readily detectable in 2 day-old colonies viewed with a Nikon SMZ1500 microscope equipped with a Chroma 59 022 (Texas red) filter (Figure 3A). Similar levels of fluorescence were detectable with codon-optimized mCher-ryRFP after a single day. The brighter appearance of mCherryRFP relative to DsRFP is consistent with observations in other organisms (Shaner *et al.*, 2005). However, the lower excitation and emission wavelengths of DsRFP relative to mCherryRFP are more compatible with some flow cytometry filter sets. Importantly, transformant colonies can be directly screened for both DsRFP and mCher-ryRFP expression using fluorescence microscopy (Figure 3B). PCR analysis of six colonies that exhibited DsRFP fluorescence revealed that all of them contained the expected junction fragments spanning *ENO1* and the *DsRFP* sequence. As a control, four colonies that did not exhibit DsRFP fluorescence in colonies from the same transformation experiment did not contain junction fragments diagnostic of *bona fide* insertion of the *DsRFP* into *ENO1*. Thus, detection of DsRFP expression at the colony level facilitates the rapid and accurate identification of potential transformants, with low levels of false-positive and false-negative results.

Finally, we generated vector pGUG, in which amplification of a *GFP-URA3-GFP* cassette yields proteins that are tagged in-frame, within the ORF, via a two-step selection scheme (Figure 4A). A similar approach has been used to epitope tag *S. cerevisiae* proteins (Schneider *et al.*, 1995). First, a cassette carrying incomplete copies of *GFP* flanking the counterselectable *URA3* marker is inserted at the locus to be tagged using homology to the insertion site (Table S3) and primers F5 and R3 (Table 1). Insertion of the cassette is facilitated by selection for *URA3*. Subsequently, growth on 5-FOA selects for cells in which recombination between the *GFP* flanking DNA results in deletion of the interrupting *URA3* sequence, while it restores a functional *GFP* gene that is in-frame with the N-terminal and C-terminal coding sequences of the genes being tagged (Figure 4A).

HWPI is a cell wall protein in which the 20 N-terminal amino acids are necessary for entry into the secretory pathway (Mao *et al.*, 2003) and the C-terminus is modified by a GPI-anchor that directs attachment of proteins to the outer surface of the plasma membrane and/or to components of the cell wall (Mao *et al.*, 2008). Expression of C-terminally tagged *HWPI-GFP* results in cytoplasmic localization of the GFP because the protein cannot be appropriately localized (Mao *et al.*, 2003). Guided by the work of Mao *et al.* (2003 (2008)), we inserted pGUG between Gln54 and Thr588, such that the remaining N- and C-terminal signals should be sufficient to localize GFP to the plasma membrane and/or cell wall. As expected, replacement of the middle 534 codons of *HWPI* with pGUG, followed by selection for loss of the *URA3* sequence on 5-FOA, resulted in localization of Hwp1-GFPp, like native Hwp1p, to the cell periphery (Figure 4B). Thus, the pGUG construct described here greatly facilitates the tagging of proteins that require both N- and C-terminal sequences for proper localization and obviates the multiple steps required to clone *GFP* within the ORF of interest prior to transformation (Mao *et al.*, 2003). Furthermore, unlike cloning

approaches used previously, use of pGUG enabled construction of a fusion protein expressed from its native promoter.

In summary, epitope tags and fluorescent protein fusions have become indispensable components of the molecular toolbox for studying the biochemistry and cell biology of all model organisms. Because of its unusual codon usage, most generic epitope tags will not function in *C. albicans*, an important human pathogen. The new cassettes constructed here extend the number of epitope tags available for use in biochemical studies of *C. albicans*. In addition, we produced a synthetic DsRFP cassette useful for fluorescent tagging and studying proteins alone or in double-labelling experiments (in combination with green or yellow fluorescent proteins). Furthermore, like codon-optimized mCherryRFP, codon-optimized DsRFP can be exploited to visually detect transformants 2 days after plating, using a simple fluorescence microscopy screen.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Mark McClellan and Aaron Christensen-Quick for web site creation and technical assistance. We thank Neta Dean for providing the mCherry GFP plasmid. This work was supported by funding from NIH, Grant Nos R01 AI0624273 and R01 AI075096, to J.B.

References

- Bahler J, Wu JQ, Longtine MS, et al. Heterologous modules for efficient and versatile PCR-based gene targeting in *Schizosaccharomyces pombe*. *Yeast*. 1998; 14:943–951. [PubMed: 9717240]
- Bertram G, Swoboda RK, Gooday GW, et al. Structure and regulation of the *Candida albicans ADH1* gene encoding an immunogenic alcohol dehydrogenase. *Yeast*. 1996; 12:115–127. [PubMed: 8686375]
- Bevis BJ, Glick BS. Rapidly maturing variants of the *Discosoma* red fluorescent protein (DsRed). *Nat Biotechnol*. 2002; 20:83–87. [PubMed: 11753367]
- Gerami-Nejad M, Berman J, Gale CA. Cassettes for PCR-mediated construction of green, yellow and cyan fluorescent protein fusions in *Candida albicans*. *Yeast*. 2001; 18:859–864. [PubMed: 11427968]
- Gerami-Nejad M, Berman J, Hausauer D, Gale CA. Cassettes for the PCR-mediated construction of regulatable alleles in *Candida albicans*. *Yeast*. 2004; 21:429–436. [PubMed: 15116343]
- Keppler-Ross S, Noffz C, Dean N. A new purple fluorescent color marker for genetic studies in *Saccharomyces cerevisiae* and *Candida albicans*. *Genetics*. 2008; 179:705–710. [PubMed: 18493083]
- Krawchuk MD, Wahls WP. High-efficiency gene targeting in *Schizosaccharomyces pombe* using a modular, PCR-based approach with long tracts of flanking homology. *Yeast*. 1999; 15:1419–1427. [PubMed: 10509024]
- Lavoie H, Sellam A, Askew C, et al. A toolbox for epitope-tagging and genome-wide location analysis in *Candida albicans*. *BMC Genomics*. 2008; 9:578. [PubMed: 19055720]
- Longtine MS, McKenzie AI, Demarini DJ, et al. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast*. 1998a; 14:953–961. [PubMed: 9717241]
- Longtine MS, McKenzie A III, Demarini DJ, et al. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast*. 1998b; 14:953–961. [PubMed: 9717241]

- Mao Y, Zhang Z, Gast C, Wong B. C-terminal signals regulate targeting of the GPI-anchored proteins to the cell wall or the plasma membrane in *Candida albicans*. *Eukaryot Cell*. 2008; 7:1906–1915. [PubMed: 18723603]
- Mao Y, Zhang Z, Wong B. Use of green fluorescent protein fusions to analyse the N- and C-terminal signal peptides of GPI-anchored cell wall proteins in *Candida albicans*. *Mol Microbiol*. 2003; 50:1617–1628. [PubMed: 14651643]
- Pfaller MA, Diekema DJ. Epidemiology of invasive candidiasis: a persistent public health problem. *Clin Microbiol Rev*. 2007; 20:133–163. [PubMed: 17223626]
- Schaub Y, Dunkler A, Walther A, Wendland J. New pFA-cassettes for PCR-based gene manipulation in *Candida albicans*. *J Basic Microbiol*. 2006; 46:416–429. [PubMed: 17009297]
- Schneider BL, Seufert W, Steiner B, et al. Use of polymerase chain reaction epitope tagging for protein tagging in *Saccharomyces cerevisiae*. *Yeast*. 1995; 11:1265–1274. [PubMed: 8553697]
- Shaner NC, Lin MZ, McKeown MR, et al. Improving the photostability of bright monomeric orange and red fluorescent proteins. *Nat Methods*. 2008; 5:545–551. [PubMed: 18454154]
- Shaner NC, Steinbach PA, Tsien RY. A guide to choosing fluorescent proteins. *Nat Methods*. 2005; 2:905–909. [PubMed: 16299475]
- Sherman, F. Getting started with yeast. In: Fink, CG., editor. *Methods in Enzymology; Guide to Yeast Genetics and Molecular Biology*. Harcourt Brace Jovanovich; San Diego: 1991. p. 3-20.
- Sung MK, Ha CW, Huh WK. A vector system for efficient and economical switching of C-terminal epitope tags in *Saccharomyces cerevisiae*. *Yeast*. 2008; 25:301–311. [PubMed: 18350525]
- Wilson RB, Davis D, Mitchell AP. Rapid hypothesis testing with *Candida albicans* through gene disruption with short homology regions. *J Bacteriol*. 1999; 181:1868–1874. [PubMed: 10074081]

Figure 1. PCR template cassettes used to generate tags at the 3' ends of *C. albicans* genes. Black boxes, *HA*, *MYC*, *GST*, *V5*, *HIS9*, *RFP* and *GFP* sequences; white boxes, *C. albicans* termination sequence (*TADH1*); grey boxes, *C. albicans* auxotrophic markers (*URA3*, *HIS1*). Oligonucleotide primers (F1, F2, F3, F4, F5, F6, R1, R2, R3) used to generate the tag cassettes by PCR are described in Materials and methods and Table 1

Figure 2.

Western blots and antibody detection of epitope-tagged strains. Lanes 1, 4, 6 and 9 correspond to the parent strain (BWP17). Lanes 2 and 3, HA-tagged strains YJB6751 (CDC3–HA *URA3*) and YJB9139 (CDC3–HA *HIS1*); lane 5, 6928 (CDC3–MYC); lanes 7 and 8, V5-tagged strains YJB8600 (Clb2–V5) and YJB8622 (Cln1–V5); lane 10, GST-tagged strain YJB8223 (CDC3–GST). Expected sizes of fusion protein products are as follows: 55 kDa, CDC3–HA; 69 kDa, CDC3–MYC; 81 kDa, CLB2–V5; 60 kDa, CLN1–V5; 75 kDa, CDC3–GST

Figure 3.

(A) Fluorescence is readily detected at the colony level after 2 days of growth. (B) Immunofluorescence micrographs of ENO1–RFP and ENO1–mCherry expressed from its native promoter. Construction of strains expressing ENO1–RFP is described in Materials and methods. Analysis of fluorescence intensities showed 5× stronger fluorescence of the strain tagged with ENO1–mCherry than with ENO1–RFP

Figure 4.

(A) Growth of HWP1–GF–URA3–FP colonies on FOA results in a restoration of the intact GFP tag. (B) Expression of *HWP1* internally tagged with *GFP* results in localization of GFP1p, like native HWP1p, to the cell periphery

Table 1

PCR primers used to amplify transformation cassettes

Primer	Epitope	Primer sequence
F1	HA, MYC, GST	5'-(gene-specific sequence) GGTGGTGGTCGGATCCCCGGGTTAATTAA-3'
F2	V5	5'-(gene-specific sequence) GGTGGTGGTGGTAAGCCTATCCCTAACCCCTCTC-3'
F3	His9	5'-(gene-specific sequence) GGTGGTGGTTTACGCCAAGCTTGGCAC-3'
F4	RFP	5'-(gene-specific sequence) GGTGGTGGTGATAAACTGAAGATGTTATT-3'
F5	GFP	5'-(gene-specific sequence) TCTAAAGGTGAAGAATTATTCACCTGG-3'
F6	M Cherry	5'-(gene-specific sequence) GGTGGTGGTGGTTTCAAAGGTGAAGAAGATAAT-3'
R1	URA3	5'-(gene-specific sequence) TCTAGAAGGACCACCTTTGATTG-3'
R2	His1	5'-(gene-specific sequence) GAATTCGGAATATTTATGAGAAAC-3'
R3	GFP	5'-(gene-specific sequence) TTTGTACAATTCATCCATAC-3'

Table 2

Strains used in this study

Strain ID	Relevant genotype	Reference
YJB9949 (BWP17)	<i>URA3Δ::λIMM434/URA3Δ::IMM434 HIS1::HISG/HIS1::HISG ARG4::HISG/ARG4::HISG</i>	Wilson, 1999
YJB6751	<i>BWP17 CDC3/CDC3::HA-URA3</i>	This study
YJB6928	<i>BWP17 CDC3/CDC3-MYC-URA3</i>	This study
YJB8223	<i>BWP17 CDC3/CDC3::GST-URA3</i>	This study
YJB8250	<i>BWP17 HWP1/HWP1::GFP</i>	This study
YJB8600	<i>BWP17 CLB2/CLB2::V5-URA3, CLN1/CLN1-MYC : HIS1</i>	This study
YJB8622	<i>BWP17 CLN1/CLN1-V5:URA3</i>	This study
YJB9139	<i>BWP17 MLC1/MLC1::YFP-URA3, CDC3/CDC3::HA-HIS1</i>	This study
YJB9452	<i>BWP17 ENO1/ENO1::RFP-URA</i>	This study
YJB11257	<i>BWP17 ENO1-mCherry-URA3</i>	This study