CORRESPONDENCE

The Prevention, Diagnosis, and Treatment of Dyslexia

by Prof. Dr. med. Gerd Schulte-Körne in volume 41/2010

Still Shameful

The article about prevention, diagnosis, and treatment of dyslexia is scientifically sound.

How then is it possible that the excellent therapeutic approach provided by the German Adult Education Association was not mentioned, which offers a free-of-charge online learning resource that has been set out brilliantly and takes learners “by the hand”, so to speak, by using multimedia images and audio (www.ich-will-lernen.de)? Subjects can use an anonymous password to log in exactly at the place where they stopped in their previous session. I have recommended this to many patients who would have normally not had the confidence to deal with this problem, which still bears the stigma of shame.

DOI: 10.3238/arztebl.2011.0262a

REFERENCES

1. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Wolfgang Ellenberger
Waldleiningen 18
69427 Mudau, Germany
Wolfgang.Ellenberger@schloss-waldleiningen.de

Auditory Perception Disorder

I was pleased to read the well structured article on this important topic.

In the context of diagnostic evaluation, it should be mentioned that in patients with symptoms of dyslexia, physical diagnostics should entail evaluation for auditory or visual impairments, and functional diagnostics should entail evaluation for auditory perception disorders (especially discrimination, phonological awareness, and seriality) as these need to be ruled out or treated concomitantly.

This is a more obvious range of conditions to evaluate diagnostically than diabetes, as mentioned in the article.

Children with auditory perception disorders and dyslexia are often found to have had speech therapy when their history is taken. If the correct pronunciation was achieved by means of oral motor exercises and tactile-kinesthetic feedback rather than by treating phonological awareness, then the problems encountered during speech acquisition will recur during the acquisition of writing skills.

In Lower Saxony, affected children can apply for in-

tegration grants according to paragraph 35a of the Child and Youth Welfare Act. The costs of educational support will then be covered for a limited time period.

The prerequisite for this is a complete diagnostic evaluation by a pediatrician, a specialist in pediatric audiology, and child and adolescent psychiatrist.

DOI: 10.3238/arztebl.2011.0262b

REFERENCES

1. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Dr. med. Jan Swart
Rehabilitationsklinik Werscherberg
Postfach 1161
49136 Bissendorf, Germany

Tests Were not Mentioned

Two lists of spelling and reading tests were printed with the article, which were said to have been standardized in the past 10 years. However, the authors omitted to mention the two tests that are most widely used in the educational setting for children and adolescents: the “Hamburger Schreibprobe” (*Hamburg spelling test*, HSP-9) was standardized in 2001 and is the most widely used test in Germany in the setting of educational diagnostic evaluation. The test has been used many millions of times in schools and has for a long time been the state-of-the-art test in this setting. In 2010 it was newly standardized; the new standards will be published in early 2011. Its electronic successor, the online test “Schreib-on” (“*write-on*”) has been available since 2007 and was also newly standardized in 2010. Both tests are widely used in the assessment of special educational needs and less so in the clinical setting, but many doctors and psychologists use it to diagnose dyslexia.

Similarly, the “Hamburger Leseprobe” (*Hamburg reading test*, HLP), which was standardized in 2003, has an important role in the assessing special education needs in individual children with reading difficulties.

Not to mention these tests in the published list represents a similar bias as if someone listing European currencies did mention the British pound sterling and the Danish krone but forgot the Euro. I assume that this is only an omission and not intentionally misleading, and I therefore recommend that the list be amended.

DOI: 10.3238/arztebl.2011.0262c

REFERENCES

1. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Dr. phil. Peter May, Dipl.-Psych.
Henriettenstr. 45
20259 Hamburg, Germany
post@peter-may.de

Professional Politics

The article does not mention that reading and writing are by necessity associated with vision, especially binocular vision, and will not be possible when these important functions are impaired. The retina is anatomically and historically the only part of the brain that is mobile and positioned in front of the top of the skull. We receive 80% of all information via the visual system, which uses up 50% of the brain's entire energy requirement. Vision is a motor activity consisting of pupil movement, accommodation, convergence, and fusion, all in the space of milliseconds. The idea that impairment in this area would not result in reading difficulties seems absurd. The usual tests for acuity of vision and stereo vision with the rapid diagnosis "eagle eyes" are not sufficient. What is needed is an analysis of associated heterophoria—"Winkelfehlsichtigkeit"—including the measurement of fixation disparity with polarized test images according to the measuring and correction method devised by Hasse. Deviations from the norm will then be corrected by using individually adjusted prism spectacles, more rarely with surgery to the ocular muscle.

Unfortunately, owing to professional considerations, this procedure is used only by very few highly specialized optometrists and ophthalmologists. On the basis of my positive experiences with many adolescents with learning difficulties, I think that this procedure is more helpful than permanent extra tuition or psychotherapy, since reading ability and thus overall performance will usually improve only once the visual function has been improved.

DOI: 10.3238/arztebl.2011.0263a

REFERENCES

1. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Dr. med. Fritz Gorzny
Schloßstr. 18
56068 Koblenz, Germany
dr.fritz_gornzy@web.de

Interdisciplinary Research

As someone working in language education I welcome the fact that *Deutsches Ärzteblatt* presented the problem of dyslexia as a CME article. Unfortunately, medical-psychological research—and usual teaching practice—do not take into consideration the results of current literacy research (1, 2).

The consensus of the diagnostic evaluation and therapy presented in the article is based on the following premises:

- Acquisition of written language skills follows the logic of a developmental model from writing true to sound to writing that is orthographically correct.
- The basis of language education in the lessons is appropriate for all groups of learners.

- Written language (orthography) is dependent on spoken language.

Current literacy research actually questions all these three premises. A reader's letter does not provide the scope to discuss the necessary revision of existing concepts in dyslexia. However, much would be gained from realizing that dyslexia is the subject of interdisciplinary research.

DOI: 10.3238/arztebl.2011.0263b

REFERENCES

1. Röber C: Der Mythos der Lauttreue. *Grundschule* 2001; (6): 40–2.
2. Hinney, G: Wortschreibungskompetenz und sprachbewusster Unterricht. Eine Alternativkonzeption zur herkömmlichen Sicht auf den Schriftspracherwerb. In: Bredel U, Müller A, Hinney G. (eds); *Schriftsystem und Schrifterwerb*. Berlin: De Gruyter 2010.
3. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Dr. phil. Gabriele Hinney
Marienburger Platz 22
31141 Hildesheim, Germany
ghinney@web.de

Important Cofactors

The article described very competently the diagnostic evaluation and therapy of dyslexia from the perspective of child and adolescent psychiatry.

In order to be able to assess the child in its entire cognitive-language learning development and not only with regard to the acquisition of written language competencies, the importance of peripheral auditory disorders, peripheral visual disorders, and auditory and visual processing and perception disorders should not be forgotten.

As an example, peripheral auditory disorders—especially unilateral or minimal hearing impairments—have been confirmed to affect the quality of performance at school (1), so that the relevant diagnostic evaluation of peripheral auditory ability should be included in the diagnostic evaluation of dyslexia.

Furthermore, close associations between specific language impairment and dyslexia are well known, so it seems advisable in a child with dyslexia to also test the current state of their expressive and receptive language skills development, in order to be able to include any deficits in these areas into the therapeutic concept.

Finally, in case of primarily perception related spelling mistakes, diagnostic evaluation of auditory processing and perception (2) may be helpful, including performance of phonological awareness, especially as training in differentiating and identifying phonemes can have a positive effect on spelling (3), whereas, according to the current state of research, merely language-free auditory training does not yield any transfer effects for reading-spelling performance (3).

DOI: 10.3238/arztebl.2011.0263c

REFERENCES

1. Nickisch A: Minimale Hörstörungen im Kindesalter. Bedeutung, Auswirkungen und Behandlung als Übersicht. Sprache-Stimme-Gehör 2009; 33: 110–5.
2. Nickisch A, Gross M, Schönweiler R, et al.: Auditive Verarbeitungs- und Wahrnehmungsstörungen. Konsensus-Statement der Deutschen Gesellschaft für Phoniatrie und Pädaudiologie. HNO 2007; 55: 61–72. PMID: 17211614
3. Nickisch A: Behandlungsmethoden von auditiven Verarbeitungs- und Wahrnehmungsstörungen: Bewertungskriterien. Kinderärztliche Praxis 2005; 76: 216–23.
4. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. Dtsch Arztebl Int 2010; 107(41): 718–27.

Dr. med. Andreas Nickisch
Heilghofstr. 63
81377 München, Germany
A.Nickisch@kinderzentrum-muenchen.de

In Reply:

The diagnostic evaluation of dyslexia according to the ICD-10 criteria requires the exclusion of auditory impairments; however, the ICD-10 does not specify which auditory impairments this includes. We presume that peripheral auditory impairments are included. Even though the rates of auditory impairments as mentioned by Nickisch are high, an association between these and dyslexia has not actually been shown. Auditory impairments can affect school performance, but the influence is an unspecific risk factor for learning performance.

The term auditory perception disorder is not clearly defined and includes a multitude of abilities for the processing of verbal and non-verbal stimuli. Such abilities to perceive and differentiate are often barely distinguishable from attention, memory capacity, phonological awareness, and intelligence. The association with dyslexia is often slight and depends on age. For this reason, there is currently no empirical basis for diagnostic procedures that reliably measure auditory perception functions specifically for dyslexia.

The importance of speech perception in dyslexia is undisputed, but distinction needs to be made between the neurobiological correlates of this speech perception, which is found to be changed in dyslexia, and the clinical diagnosis of a receptive or expressive speech impairment (1). Speech perception impairments in children with receptive and/or expressive speech disorders are different to those in children with dyslexia (2).

Skills such as phonological awareness need to be clearly defined in this context, as they are important for predicting reading and writing competence and are also used for preventive measures before the child starts school. Since phonological competencies represent an important factor for the development of reading and writing skills in some children with dyslexia beyond the preschool years, the BAKO (*Basiskompetenzen für Lese-Rechtschreibleistungen*, test battery measuring phonological processing) provides a diagnostic

procedure that measures these competencies. The diagnostic procedure is recommended for all children with dyslexia who have difficulties in perceiving and differentiating speech sounds at preschool stage. Current results from a meta-analysis of German-language intervention studies of dyslexia, however, show that supporting phonological awareness is effective only for the first year at primary school.

The importance of the diagnostic evaluation of eye movements for dyslexia is still the subject of controversial discussion. Using surgery to treat an apparent eye movement disorder in order to eliminate the cause of dyslexia has to be regarded as a treatment error and is not supported by studies (4). Gorzny's recommendation—to omit the only evaluated therapeutic approaches and treat dyslexic children with prism spectacles—is fatal (5).

Since dyslexia usually affects the affected person for their entire lifetime, supporting adults with dyslexia is important. The fact that Germany's adult education colleges have taken on supporting reading and writing skills in functionally illiterate adults is important and to be welcomed. However, this service is not specific to the needs of adults with dyslexia. Such adults mostly require specific support because—in contrast to most illiterate persons, who were just never taught reading and writing—they have received instruction but need specific help to improve their reading speed and spelling skills.

The “Hamburger Schreibprobe” is very popular, especially with teachers, and widely used in schools. However, its popularity does not enable any conclusions regarding its quality in terms of being a state-of-the-art tool. As soon as the new standards become available, the approach can be freshly evaluated. The HSP has been criticized by Tacke (6) for its lack of ability to differentiate. That means that the HSP on average results in higher performance ratings (percentage rankings) than other spelling tests—at least in those federal states where spelling competence is high compared with the rest of Germany. This effect hampers the test's clinical use in diagnostic evaluation. The HSP is an approach that, according to its author, from whom the test needs to be purchased, should become widely used. According to the author, the test was developed in 1993 and has been available with Germany-wide standards since 2000, and has now been around for a decade—a period after which a test has to be regarded as outdated.

DOI: 10.3238/arztebl.2011.0264

REFERENCES

1. Schulte-Körne G: Lese-Rechtschreibstörung und Sprachwahrnehmung. Psychometrische und neurophysiologische Untersuchungen zur Legasthenie. Münster: Waxmann Verlag 2001.
2. Robertson EK, Joanisse MF, Desroches AS, Ng S: Categorical speech perception deficits distinguish language and reading impairments in children. Dev Sci 2009; 12(5): 753–67.
3. Stock C, Marx P, Schneider W: Basiskompetenzen für Lese-Rechtschreibleistungen (BAKO 1–4). Göttingen: Hogrefe 2003.

4. American Academy of Ophthalmology: Policy Statement Learning Disabilities, Dyslexia, Vision. <http://www.aao.org/about/policy/upload/Learning-Disabilities-Dyslexia-Vision-2009.pdf>
5. Trauzettel-Klosinski S, Schäfer WD, Klosinski G: Legasthenie Grundlagen des Lesens – Lese-Rechtschreib-Störung – okuläre Lese-störung. *Der Ophthalmologe* 2002; 99(3): 208–27.
6. Tacke G, Völker R, Lohmüller R: Die Hamburger Schreibprobe: Probleme mit einem neuen Rechtschreibtest Psychologie in Erziehung und Unterricht 2001; 48(2): 135–45.
7. Schulte-Körne G: The prevention, diagnosis, and treatment of dyslexia. *Dtsch Arztebl Int* 2010; 107(41): 718–27.

Prof. Dr. med. Gerd Schulte-Körne

Direktor der Klinik für Kinder- und Jugendpsychiatrie, Psychosomatik und Psychotherapie Klinikum der Universität München
 Pettenkoferstr. 8a
 80336 München, Germany
 Gerd.Schulte-Koerne@med.uni-muenchen.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.