Published in final edited form as:

Ann Epidemiol. 2011 June ; 21(6): 399–406. doi:10.1016/j.annepidem.2011.03.003.

The Epidemiology of Hospitalized Postpartum Depression in New York State, 1995–2004

David A. Savitz, Ph.D.,

Departments of Community Health and Obstetrics and Gynecology, Brown University,
Providence, Rhode Island

Cheryl R. Stein, Ph.D.,

Department of Preventive Medicine, Mount Sinai School of Medicine, New York, New York

Fen Ye, M.S.,

Department of Medicine, Montefiore Medical Center, Bronx, New York

Lisa Kellerman, M.S., and

Department of Preventive Medicine, Mount Sinai School of Medicine, New York, New York

Michael Silverman, Ph.D.

Department of Psychiatry, Mount Sinai School of Medicine, New York, New York

Abstract

Purpose—The purpose of this study is to describe the patterns of hospitalization for depression in the year following delivery in relation to social, demographic, and behavioral characteristics.

Methods—Data on fetal deliveries were linked to hospitalizations for depression over the subsequent year in order to describe the frequency and patterns of hospitalized postpartum depression among 2,355,886 deliveries in New York State from 1995 – 2004. We identified “definite postpartum depression” based on ICD codes indicative of “mental disorders specific to pregnancy,” and “possible postpartum depression” by ICD codes for hospitalization with any depressive disorders.

Results—In New York State, we identified 1,363 women (5.8 per 10,000) who were hospitalized with definite postpartum depression, and 6,041 women (25.6 per 10,000) with possible postpartum depression, with lower risks in the New York City area. Postpartum depression was more common in later years and among mothers who were older, Black, smokers, lacking private insurance, and with multiple gestations, and was rarer among Asians. For possible postpartum depression, socioeconomic gradients were enhanced.

Conclusions—Risk of hospitalized postpartum depression is strongly associated with socioeconomic deprivation and varies markedly by ethnicity, with direct implications for screening and health services, also providing suggestions for etiologic studies.

© 2011 Elsevier Inc. All rights reserved.

Address for reprints: David A. Savitz, Ph.D., Department of Community Health, Brown University, Box G-S-121-2, Providence, RI 02912, David_savitz@brown.edu.

Location of work: Mount Sinai School of Medicine, New York, New York

None of the authors have a financial conflict of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

postpartum depression; pregnancy; ethnicity

INTRODUCTION

Postpartum depression, the most common complication associated with childbirth, affects an estimated 10 – 15% of women following delivery (1), with a wide range of severity. The Diagnostic and Statistical Manual of Mental Disorders, fourth edition (DSM-IV) (2) defines postpartum depression as major depression, using the same diagnostic criteria but with onset of depressive symptoms within four weeks after delivery. However, researchers often extend the time window for identifying pregnancy-related depression to the full year following birth (3). Given that pregnancy itself is a common event, postpartum depression constitutes a major public health concern deserving improved information on etiology and service needs. While there are multiple etiologic pathways that may explain the disorder, with genetic, endocrine hormone, neurotransmitter, psychosocial, and psychoanalytic hypotheses (4), the genesis of postpartum depression remains poorly understood.

Major depressive episodes subsequent to birth result in marked functional impairment and distress (5), as well as an increased risk for suicide (6). Maternal depression is also known to impact the health of the child through several pathways (7–9), including shortened duration of breastfeeding (10, 11) and renewal of maternal smoking (12, 13). Interactions of depressed mothers and their children are of lower quality than those of non-depressed mothers (7, 14, 15), adversely affecting neurobehavioral development and increasing the risk of conduct disorders and greater child insecurity in attachment relationships (16–21). The lack of energy and inability to cope effectively with the demands of the period after birth constitute a serious threat to the infant's well-being (22).

Much of the previous research on postpartum depression has employed small, convenience samples from clinics or hospitals that cannot generate normative data and are subject to varying diagnostic criteria. Additionally, most studies focus on *depressive symptomatology* rather than *clinical diagnoses of depression*, with markedly higher rates of depressive symptoms when minor or transient episodes are included (1). Of the studies examining diagnosed depression, few differentiate between minor and major depression (23). We report here on the frequency and pattern of severe depression leading to hospitalization in the year following delivery in a population of over 2.3 million women giving birth in New York State over a 10-year period.

METHODS

Study Population

Hospital discharge data from the New York State Department of Health's Statewide Planning and Research Cooperative System were used to identify maternal hospitalizations for delivery from 1995 – 2004, and linked on an individual basis to subsequent hospitalizations within the 12 months following delivery. Deliveries were identified by an *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM) (24) diagnosis code of V27 (outcome of delivery) or a federal or state diagnosis related group code of 370 – 375 (delivery services). Unique identifiers permitted extraction of all hospitalizations of the mother in New York State during the 12 months following delivery. For residents of New York City only (excluding residents of New York State outside New York City), we linked birth records from the New York City Department of Health and

Mental Hygiene to the hospital discharge data. Because of administrative restrictions, New York State birth records were not available for this purpose.

Starting with 2,526,504 deliveries, we excluded 64,197 (2.5%) records: 9,483 (0.4%) records with an AIDS flag that precluded use of identifiers needed to perform data linkages; 13,199 (0.5%) duplicates or non-unique identifiers indicative of database errors; and 41,515 (1.6%) non-New York residents who are not part of the population of interest. This resulted in 2,355,886 deliveries for New York residents from 1995 – 2004, and 106,421 episodes of maternal hospitalization in the year following delivery.

Study approval was obtained from the Mount Sinai School of Medicine Program for the Protection of Human Subjects, the New York State Department of Health Data Protection Review Board, the New York State Department of Health Institutional Review Board, and the New York City Department of Health and Mental Hygiene Institutional Review Board.

Measures

We include as cases hospitalized diagnoses of depression occurring within 12 months postpartum as ascertained by review of hospital discharge records. Using ICD-9-CM codes we created categories of “definite” and “possible” postpartum depression. Definite postpartum depression required an ICD-9-CM code of 648.40 – 648.42 or 648.44. These codes explicitly link “mental disorders specific to pregnancy” within the first 4 weeks postpartum, and if used properly, they reflect episodes of postpartum depression. The available data, however, included all hospitalization discharges with depression diagnoses within 12 months postpartum, allowing us to examine “possible” postpartum depression by also including ICD-9-CM associated with unipolar and bipolar depression: 296.20 – 296.26, 296.80 – 296.82, 296.89, 309.0, 309.1, 309.24, 309.28, 309.29, and 311. It is likely that this broader grouping sacrifices specificity for increased sensitivity, but the extent of this misclassification is unknown.

Statistical Analysis

Maternal demographic and health characteristics were identified from the hospital discharge record for all New York State residents, including delivery year, season of delivery, maternal age, maternal race/ethnicity, plurality, insurance type, and residential location, classified by New York State region and urban/rural status. Plurality was identified by the third digit of ICD-9-CM code V27 indicating a single or multiple gestation. Insurance type was determined by the expected reimbursement type and coded as “Private” (including workers’ compensation, Blue Cross, commercial insurance company, health maintenance organization, Champus/Veteran’s Administration, or self-administered plan), “Public” (including Medicare, Medicaid, other governmental plan, or federal, state or local corrections services), “Self-pay,” or “Other” (including records marked no charge, other, or no-fault). New York was divided into 6 administrative regions (New York City; Long Island; Hudson Valley; Central; Western; Northern) based on county of residence (25). Urban/rural status was categorized according to National Center for Health Statistics guidelines based on county population size (26) listed from most urban to most rural: metropolitan area: large central counties in metro area >1 million population; metropolitan area: large fringe counties in metro area >1 million population; medium metropolitan area: counties in metro area 250,000 – 999,999 population; small metropolitan area: counties in metro area 50,000 – 249,999 population; micropolitan counties; non-core counties.

For births to residents of New York City we restricted the analysis to singleton births to facilitate matching to the birth certificate. Of the 1,124,073 New York City deliveries, 900,726 were singletons with a matched birth certificate record. For this subset we were also

able to consider information obtained from the birth certificate, but not available from hospital discharge data: parity (0, 1, 2+), nativity (US or foreign born), education (<12, 12, 13 – 16, 17+ years), prenatal tobacco use (smoker, non-smoker), and trimester of prenatal care initiation (1st, 2nd, 3rd, no prenatal care).

We calculated the risk of definite and possible postpartum depression per 10,000 deliveries for the whole population and subsets defined by attributes of interest, noted above. Prevalence odds ratios and 95% confidence intervals were estimated for each covariate using logistic regression. Missing and/or unknown were included as a distinct category for the bivariate analyses. Multivariate prevalence odds ratios and 95% confidence intervals were also derived, with adjustment for all other covariates. We were not able to account for repeat births to the same mother so variance estimates are slightly underestimated due to non-independence of observations. For multivariate models, unknown race/ethnicity was included as a separate category, but the few individuals missing data on other characteristics were excluded. Because the demographic characteristics and availability of birth records differed between New York City and other parts of New York State, we analyzed and presented results for these subsets separately. All analyses were performed using SAS version 9.2 (SAS Institute, Cary, NC).

RESULTS

During the period 1995–2004 for New York State as a whole, we identified 1,363 women (5.8 per 10,000) who were hospitalized in the year following delivery specifically for depression related to childbirth. The broader definition of possible postpartum depression, increasing sensitivity at the expense of specificity, identified over four times as many hospitalizations (6,041 in total for a prevalence of 25.6 per 10,000) (data for all of New York State not shown). Results are presented separately for New York State outside of New York City and for New York City because of social and demographic differences and the availability of birth certificate information for New York City only. Focusing first on definite postpartum depression, some striking patterns were apparent in New York State outside of New York City (Table 1): increased frequency in the latter part of the time period, for multiple gestations (aOR = 2.1, 95% CI = 1.5–3.0), among Black women (aOR = 1.4, 95% CI = 1.2–1.7), mothers age 35 or older (aOR = 1.5, 95% CI = 1.3–1.8), and among women with public insurance (lower socioeconomic status) (aOR = 3.3, 95% CI = 2.8–3.8). Reduced risk was noted for Asian women (OR = 0.5, 95% CI = 0.2–1.0) and among residents of Long Island and the Hudson Valley. More modest and non-monotonic variation was found in relation to urban-rural residence. In New York City, most of the same trends were found as in New York State outside of New York City (Table 2), including increasing risk in more recent years, and elevated risks for African-American mothers (aOR = 1.7, 95% CI = 1.2–2.4) and mothers who had public insurance (aOR = 1.7, 95% CI = 1.3–2.3), the latter lower in magnitude than the increase found in New York State. Smokers were at markedly elevated risk (aOR = 3.0, 95% CI = 2.0–4.5). Modest decreases were noted for parous women and foreign-born women. Adjustment for confounding had little impact on most measures of association, except for a marked reduction in the risk for non-Hispanic black women and for Hispanic women (in New York City), which was largely due to adjustment for indicators of socioeconomic status which is a far stronger predictor than race/ethnicity.

Despite markedly greater numbers for *possible* postpartum depression, the patterns were generally quite similar as for definite postpartum depression. In New York State (exclusive of New York City), a higher risk of possible postpartum depression was found for more recent births, multiple gestations, and births to Black women and those lacking private insurance. Lower risk was again noted for Asian women and residents of Long Island and

the Hudson Valley (Table 1). The same contrast of possible and definite postpartum depression also generated a number of similarities in the New York City population (Table 2): elevated risks for the more recent time period, for Black women, women on public insurance, and smokers, and reduced risks for Asians and foreign-born women. In addition, however, the more inclusive definition identified elevated risk for Hispanic mothers, mothers <20 years old, women of parity 2 or greater, a sharper gradient by educational level, and increased risks for those with late or no prenatal care. In general, there seem to be more and perhaps stronger socioeconomic gradients for *possible* than for *definite* postpartum depression.

DISCUSSION

The proportion of women in the United States with postpartum depression severe enough to require hospitalization is unknown, with little opportunity to compare our population-based rates to other populations or time periods that focus on milder cases. Studies from Scotland (27), Denmark (3), and Sweden (28) yielded hospitalization rates in the first three months postpartum broadly comparable to the ones obtained here, ranging from 10.3–21.9 per 10,000, respectively. The differing time periods for inclusion following delivery, varying scope of psychiatric hospitalizations included, and different calendar time periods and geographic settings limit the ability to make meaningful comparisons with our frequencies of hospitalized depression.

While comparison of absolute rates between our study and other studies is tempered by the non-comparability of case definitions, examinations within our population may generate more comparable information. Frequency of postpartum depression has been reported to vary by race and ethnicity, with Hispanics having the lowest prevalence and Native Americans the highest (29–31). The elevated risk among Hispanics in New York City but not elsewhere in New York State, suggests heterogeneity among Hispanic subgroups. The proportion Hispanic differs considerably, 6.4% in New York State other than New York City and 27.0% in New York City, but the composition in broad categories is similar, with a sizable fraction from Puerto Rico (~37% of Hispanics in both regions) though the identity of “other Hispanics” (which constitute around half in both regions) cannot be compared. Recent immigration has also been noted as a risk factor, where new migrants may be socially isolated or unaware of how to access mental health services (32–35) whereas we found reduced risk for foreign-born women in New York City (though we did not have information on duration of time in the US). Alternatively, there may be distinctive cultural factors affecting rates of treated postpartum depression, with some studies among Asian-American women supporting this possibility (36, 37).

Urbanization has been associated with higher rates of depressive disorders after delivery; alternatively urban residents in need may simply be more likely to access care and be counted (38, 39). We did not find any clear association with urban/rural residence in New York State, and the markedly lower rates in New York City relative to New York State are counter to such a pattern.

The effect of maternal age is indeterminate, with some studies showing that adolescents have a higher risk of postpartum depression (40), and others showing a five-fold increased risk for women over the age of 30 as compared to teens (41). We found an association between older maternal age and definite postpartum depression in New York State, but not in New York City. Primiparous women are also thought to be at higher risk (42, 43), which we found for definite cases in New York City. It has been speculated that this may reflect selective childbearing in that women who have experienced postpartum depression may be reluctant to become pregnant subsequently.

Life stressors, including major events before, during, and after pregnancy, marital discord, and adverse socioeconomic conditions have been variably associated with increased risk of postpartum depression (35, 44–52). We could not examine psychosocial stress directly in our study, the clear and strong association with insurance type, a marker of economic stress, is consistent with the reported patterns (53). However, there are multiple pathways by which insurance status may predict hospitalization for depression, including serving as a marker of risk associated with lower socioeconomic status and providing a financial resource to cover costs associated with such hospitalization.

The broadly comparable results for women in New York State and New York City are notable, despite a nearly two-fold greater absolute risk in the former as compared to the latter. For the social and demographic factors that were available for comparison (age, race/ethnicity, insurance type), the patterns were quite comparable. Despite a markedly larger number of women with hospitalization for “possible” rather than “definite” postpartum depression, the patterns were similar by race/ethnicity, insurance type, nativity, education, and tobacco use. However the differences are notable for some of the attributes examined in New York City: possible postpartum depression showed increased risks for women of parity two or greater and women with later prenatal care initiation, not found for definite postpartum depression.

There are several limitations in our approach that should be recognized, most notably the potential for differential access to and use of inpatient care for depression. While the presence and severity of depression would ideally be the primary determinant of whether hospitalization occurs, the additional requirement is obtaining medical care for the condition. Women who have access to family resources, outpatient services, and other ways of managing their depression without hospitalization would be less likely to be admitted. On the other end, those women who are disconnected from medical care access due to geographic, cultural, or socioeconomic isolation would be less likely to access care and be hospitalized. In addition, the coding of postpartum depression by different service providers, a proclivity to use one diagnosis or another for similar conditions as a function of age, ethnicity, socioeconomic status, etc., will influence the patterns of risk in unpredictable ways.

We did not have information on prior episodes of depression or depression that began during rather than after the pregnancy, strong predictors of postpartum depression (54) and plausibly related to a number of the predictors of interest. The independent etiologic influence of these general markers of social and behavioral predictors is limited by the lack of detailed information on the more proximal mediators of risk for postpartum depression. Finally, we were not able to account for repeat births to the same mother, making our confidence intervals slightly narrower than they should have been, a problem that is mitigated by the extremely large population included in the study.

These data provide important new information to assess the magnitude of the public health burden associated with postpartum depression and point to potential determinants. The sizable variation in risk across geographic areas, especially across regions of New York State, and the dramatic increase in hospitalization over the period between 1995 and 2004, raises the possibility of differential use of health services across areas and time periods. It seems implausible that the frequency of the underlying problem could vary more than two-fold across adjacent locations and a narrow time period. A closer examination of the extent to which health care access accounts for these patterns may be warranted. The notable elevation in risk for women of lower socioeconomic status, African-Americans, and at least some Hispanic subgroups points towards the social stress of poverty, lack of resources, or possibly discrimination as potential contributors to the etiology of postpartum depression.

Given that nearly all women receive prenatal care and deliver in hospitals, there are clear opportunities to prevent or intervene early in the psychiatric disorders that follow. At minimum, our data identify the women who are at greatest risk, warrant closer follow-up, and should be considered for earlier or more intensive intervention.

Acknowledgments

David Savitz, Cheryl Stein, and Michael Silverman conceptualized the study plans, David Savitz, Cheryl Stein, Lisa Kellerman, Fen Ye, and Michael Silverman contributed to the analysis and interpretation of the results, David Savitz, Lisa Kellerman, and Michael Silverman contributed to the drafting of the manuscript, and David Savitz, Cheryl Stein, Lisa Kellerman, and Michael Silverman provided substantive review and comment on the draft manuscript. The research was supported by Grant R21HD58111 from the National Institute of Child Health and Human Development. David Savitz had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. O'Hara MW, Swain AM. Rates and risk of postpartum depression--a meta-analysis. *Int Rev Psychiatry*. 1996; 8(1):37–53.
2. American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-IV. 4. Washington, DC: American Psychiatric Association; 1994. Task Force on DSM-IV.
3. Munk-Olsen T, Laursen TM, Pedersen CB, Mors O, Mortensen PB. New parents and mental disorders: a population-based register study. *Jama*. 2006; 296(21):2582–9. [PubMed: 17148723]
4. Tovino SA. Scientific understandings of postpartum illness: improving health law and policy? *Harvard Journal of Law and Gender*. 2010; 33(1):99–173.
5. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder (revision). *Am J Psychiatry*. 2000; 157(4 Suppl):1–45.
6. Oates M. Suicide: the leading cause of maternal death. *Br J Psychiatry*. 2003; 183:279–81. [PubMed: 14519602]
7. Stein A, Gath DH, Bucher J, Bond A, Day A, Cooper PJ. The relationship between post-natal depression and mother-child interaction. *Br J Psychiatry*. 1991; 158:46–52. [PubMed: 2015451]
8. Chung EK, McCollum KF, Elo IT, Lee HJ, Culhane JF. Maternal depressive symptoms and infant health practices among low-income women. *Pediatrics*. 2004; 113(6):e523–9. [PubMed: 15173532]
9. Brand SR, Brennan PA. Impact of antenatal and postpartum maternal mental illness: how are the children? *Clin Obstet Gynecol*. 2009; 52(3):441–55. [PubMed: 19661760]
10. McCarter-Spaulding D, Horowitz JA. How does postpartum depression affect breastfeeding? *MCN Am J Matern Child Nurs*. 2007; 32(1):10–7. [PubMed: 17308452]
11. Alvik A, Haldorsen T, Lindemann R. Alcohol consumption, smoking and breastfeeding in the first six months after delivery. *Acta Paediatr*. 2006; 95(6):686–93. [PubMed: 16754549]
12. Suplee PD. The importance of providing smoking relapse counseling during the postpartum hospitalization. *J Obstet Gynecol Neonatal Nurs*. 2005; 34(6):703–12.
13. Solomon LJ, Higgins ST, Heil SH, Badger GJ, Thomas CS, Bernstein IM. Predictors of postpartum relapse to smoking. *Drug Alcohol Depend*. 2007
14. Fleming A, Flett G, Ruble D, Shaul D. Postpartum adjustment in first-time mothers: relations between mood, maternal attitudes and mother-infant interactions. *Dev Psychol*. 1988; 24:71–81.
15. Field T. Postpartum depression effects on early interactions, parenting, and safety practices: a review. *Infant Behav Dev*. 2010; 33(1):1–6. [PubMed: 19962196]
16. Murray L, Stein A. The effects of postnatal depression on the infant. *Baillieres Clin Obstet Gynaecol*. 1989; 3(4):921–33. [PubMed: 2700150]
17. Murray L. The impact of postnatal depression on infant development. *J Child Psychol Psychiatry*. 1992; 33(3):543–61. [PubMed: 1577898]
18. Murray L, Fiori-Cowley A, Hooper R, Cooper P. The impact of postnatal depression and associated adversity on early mother-infant interactions and later infant outcome. *Child Dev*. 1996; 67(5):2512–26. [PubMed: 9022253]

19. Murray L, Cooper P. Effects of postnatal depression on infant development. *Arch Dis Child*. 1997; 77(2):99–101. [PubMed: 9301345]
20. Civic D, Holt VL. Maternal depressive symptoms and child behavior problems in a nationally representative normal birthweight sample. *Matern Child Health J*. 2000; 4(4):215–21. [PubMed: 11272341]
21. Field T, Diego M, Hernandez-Reif M, Schanberg S, Kuhn C. Depressed mothers who are “good interaction” partners versus those who are withdrawn or intrusive. *Infant Behav Dev*. 2003; 26:238–52.
22. Wisner KL, Parry BL, Piontek CM. Clinical practice. Postpartum depression. *N Engl J Med*. 2002; 347(3):194–9. [PubMed: 12124409]
23. Gaynes BN, Gavin N, Meltzer-Brody S, Lohr KN, Swinson T, Gartlehner G, et al. Perinatal depression: prevalence, screening accuracy, and screening outcomes. *Evid Rep Technol Assess (Summ)*. 2005(119):1–88.
24. Official authorized errata for the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM). *J Am Med Rec Assoc*. 1983; 54(8):29–32.
25. Century Commission on Health Care Facilities in the 21st Century. 2007. [updated 2007; cited]; Available from: <http://www.nyhealthcarecommission.org/regions/>
26. Ingram, DD.; Franco, S. 2006 NCHS Urban-Rural Classification Scheme for Counties. National Center for Health Statistics; 2006.
27. Kendell RE, Rennie D, Clarke JA, Dean C. The social and obstetric correlates of psychiatric admission in the puerperium. *Psychol Med*. 1981; 11(2):341–50. [PubMed: 7267875]
28. Valdimarsdottir U, Hultman CM, Harlow B, Cnattingius S, Sparen P. Psychotic illness in first-time mothers with no previous psychiatric hospitalizations: a population-based study. *PLoS Med*. 2009; 6(2):e13. [PubMed: 19209952]
29. Baker L, Oswalt K. Screening for postpartum depression in a rural community. *Community Ment Health J*. 2008; 44(3):171–80. [PubMed: 18049894]
30. Wei G, Greaver LB, Marson SM, Herndon CH, Rogers J. Postpartum depression: racial differences and ethnic disparities in a tri-racial and bi-ethnic population. *Matern Child Health J*. 2008; 12(6): 699–707. [PubMed: 17955356]
31. Zittel-Palamara K, Rockmaker JR, Schwabel KM, Weinstein WL, Thompson SJ. Desired assistance versus care received for postpartum depression: access to care differences by race. *Arch Womens Ment Health*. 2008; 11(2):81–92. [PubMed: 18463943]
32. Sword W, Watt S, Krueger P. Postpartum health, service needs, and access to care experiences of immigrant and Canadian-born women. *J Obstet Gynecol Neonatal Nurs*. 2006; 35(6):717–27.
33. Abe-Kim J, Takeuchi DT, Hong S, Zane N, Sue S, Spencer MS, et al. Use of mental health-related services among immigrant and US-born Asian Americans: results from the National Latino and Asian American Study. *Am J Public Health*. 2007; 97(1):91–8. [PubMed: 17138905]
34. Jackson JS, Neighbors HW, Torres M, Martin LA, Williams DR, Baser R. Use of mental health services and subjective satisfaction with treatment among Black Caribbean immigrants: results from the National Survey of American Life. *Am J Public Health*. 2007; 97(1):60–7. [PubMed: 17138907]
35. Davey HL, Tough SC, Adair CE, Benzies KM. Risk Factors for Sub-Clinical and Major Postpartum Depression Among a Community Cohort of Canadian Women. *Matern Child Health J*. 2008
36. Klainin P, Arthur DG. Postpartum depression in Asian cultures: a literature review. *Int J Nurs Stud*. 2009; 46(10):1355–73. [PubMed: 19327773]
37. Lee DT, Yip AS, Leung TY, Chung TK. Ethnoepidemiology of postnatal depression. Prospective multivariate study of sociocultural risk factors in a Chinese population in Hong Kong. *Br J Psychiatry*. 2004; 184:34–40. [PubMed: 14702225]
38. Mueller DP. The current status of urban-rural differences in psychiatric disorder. An emerging trend for depression. *J Nerv Ment Dis*. 1981; 169(1):18–27. [PubMed: 7452218]
39. McGarry J, Kim H, Sheng X, Egger M, Baksh L. Postpartum depression and help-seeking behavior. *J Midwifery Womens Health*. 2009; 54(1):50–6. [PubMed: 19114239]

40. Lanzi RG, Bert SC, Jacobs BK. Depression among a sample of first-time adolescent and adult mothers. *J Child Adolesc Psychiatr Nurs.* 2009; 22(4):194–202. [PubMed: 19930300]
41. Luke S, Salihu HM, Alio AP, Mbah AK, Jeffers D, Berry EL, et al. Risk factors for major antenatal depression among low-income African American women. *J Womens Health (Larchmt).* 2009; 18(11):1841–6. [PubMed: 19951220]
42. Stewart DE, Gagnon A, Saucier JF, Wahoush O, Dougherty G. Postpartum depression symptoms in newcomers. *Can J Psychiatry.* 2008; 53(2):121–4. [PubMed: 18357931]
43. Dipietro JA, Costigan KA, Sipsma HL. Continuity in self-report measures of maternal anxiety, stress, and depressive symptoms from pregnancy through two years postpartum. *J Psychosom Obstet Gynaecol.* 2008; 29(2):115–24. [PubMed: 18655259]
44. O'Hara MW, Rehm LP, Campbell SB. Predicting depressive symptomatology: cognitive-behavioral models and postpartum depression. *J Abnorm Psychol.* 1982; 91(6):457–61. [PubMed: 7153419]
45. Righetti-Veltema M, Conne-Perreard E, Bousquet A, Manzano J. Risk factors and predictive signs of postpartum depression. *J Affect Disord.* 1998; 49(3):167–80. [PubMed: 9629946]
46. McGill H, Burrows VL, Holland LA, Langer HJ, Sweet MA. Postnatal depression: a Christchurch study. *N Z Med J.* 1995; 108(999):162–5. [PubMed: 7753510]
47. Beck CT, Gable RK. Further validation of the Postpartum Depression Screening Scale. *Nurs Res.* 2001; 50(3):155–64. [PubMed: 11393637]
48. Paykel ES, Emms EM, Fletcher J, Rassaby ES. Life events and social support in puerperal depression. *Br J Psychiatry.* 1980; 136:339–46. [PubMed: 7388241]
49. Hickey AR, Boyce PM, Ellwood D, Morris-Yates AD. Early discharge and risk for postnatal depression. *Med J Aust.* 1997; 167(5):244–7. [PubMed: 9315011]
50. Goyal D, Gay C, Lee KA. How much does low socioeconomic status increase the risk of prenatal and postpartum depressive symptoms in first-time mothers? *Women's Health Issues.* 2010
51. Hayes DK, Ta VM, Hurwitz EL, Mitchell-Box KM, Fuddy LJ. Disparities in self-reported postpartum depression among Asian, Hawaiian, and Pacific Islander women in Hawaii: Pregnancy Risk Assessment Monitoring System (PRAMS), 2004–2007. *Matern Child Health J.* 2009
52. Canady RB, Bullen BL, Holzman C, Broman C, Tian Y. Discrimination and symptoms of depression in pregnancy among African American and White women. *Womens Health Issues.* 2008; 18(4):292–300. [PubMed: 18590883]
53. Prevalence of self-reported postpartum depressive symptoms--17 states, 2004–2005. *MMWR Morb Mortal Wkly Rep.* 2008; 57(14):361–6. [PubMed: 18401329]
54. O'Hara MW. Postpartum depression: what we know. *J Clin Psychol.* 2009

Table 1

Maternal Hospitalization for Depression within 12 Months of Delivery, New York State (excluding New York City), 1995 – 2004, N = 1,231,813 Deliveries

Characteristic	Cases (N)	Definite Postpartum Depression			Possible Postpartum Depression		
		Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)	Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)
All Deliveries	907	7.4			31.8		
Year (delivery)							
1995 – 1998	209	4.4	1.0	1.0	24.9	1.0	1.0
1999 – 2001	240	6.3	1.4 (1.2–1.7)	1.4 (1.2–1.7)	30.0	1.2 (1.1–1.3)	1.2 (1.1–1.3)
2002 – 2004	458	12.2	2.8 (2.4–3.3)	2.7 (2.3–3.1)	42.4	1.7 (1.6–1.8)	1.7 (1.6–1.8)
Season (delivery)							
Spring (March – May)	207	6.6	0.9 (0.8–1.1)	0.9 (0.8–1.1)	29.8	1.0 (0.9–1.1)	1.0 (0.9–1.1)
Summer (June – August)	236	7.2	1.0	1.0	30.9	1.0	1.0
Fall (September – November)	243	8.0	1.1 (0.9–1.3)	1.1 (0.9–1.3)	33.8	1.1 (1.0–1.2)	1.1 (1.0–1.2)
Winter (December – February)	221	7.7	1.1 (0.9–1.3)	1.0 (0.9–1.2)	32.8	1.1 (1.0–1.2)	1.0 (0.9–1.1)
Plurality							
Single gestation	874	7.2	1.0	1.0	31.5	1.0	1.0
Multiple gestation	32	14.6	2.0 (1.4–2.9)	2.1 (1.5–3.0)	44.3	1.4 (1.1–1.7)	1.6 (1.3–1.9)
Unknown/unspecified gestation	1				4		
Maternal Age (years)							
<20	74	7.8	1.1 (0.9–1.4)	0.7 (0.5–0.9)	53.5	1.7 (1.6–1.9)	1.0 (0.9–1.1)
20 – 34	653	7.2	1.0	1.0	31.0	1.0	1.0
35+	180	8.0	1.1 (0.9–1.3)	1.5 (1.3–1.8)	25.7	0.8 (0.8–0.9)	1.2 (1.1–1.3)
Maternal Race/Ethnicity							
White non-Hispanic	538	6.9	1.0	1.0	29.5	1.0	1.0
Black non-Hispanic	150	14.2	2.1 (1.7–2.5)	1.4 (1.2–1.7)	65.6	2.2 (2.0–2.4)	1.6 (1.4–1.7)
Hispanic	45	5.6	0.8 (0.6–1.1)	0.6 (0.5–0.9)	24.1	0.8 (0.7–0.9)	0.6 (0.5–0.7)
Asian	7	3.1	0.4 (0.2–0.9)	0.5 (0.2–1.0)	13.6	0.5 (0.3–0.7)	0.5 (0.4–0.7)
Other	16	3.9	0.6 (0.3–0.9)	0.5 (0.3–0.8)	17.6	0.6 (0.5–0.8)	0.5 (0.4–0.7)
Unknown	151	7.4	1.1 (0.9–1.3)	1.0 (0.8–1.2)	30.7	1.0 (1.0–1.1)	1.0 (0.9–1.1)
Insurance Type							
Private	388	4.5	1.0	1.0	17.1	1.0	1.0

Characteristic	Definite Postpartum Depression			Possible Postpartum Depression		
	Cases (N)	Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)	Cases (N)	Proportion (per 10,000)
Public	483	14.7	3.2 (2.8–3.7)	3.3 (2.8–3.8)	2,288	69.5
Self-pay	27	8.7	1.9 (1.3–2.8)	2.1 (1.4–3.2)	130	41.7
Other	9	5.7	1.3 (0.6–2.4)	1.3 (0.7–2.6)	34	21.4
New York Region						
Long Island	152	4.7	0.5 (0.4–0.6)	0.7 (0.5–0.9)	592	18.4
Hudson Valley	120	4.6	0.5 (0.4–0.6)	0.6 (0.4–0.7)	630	24.3
Central	307	9.6	1.0	1.0	1,332	41.6
Western	191	11.1	1.2 (1.0–1.4)	1.1 (0.9–1.3)	744	43.1
Northern	127	9.0	0.9 (0.8–1.2)	1.1 (0.9–1.3)	592	42.1
Unknown	10	5.8	0.6 (0.3–1.1)		24	13.8
Urban/Rural Status						
Metro Area: Large Central Counties	209	11.3	1.5 (1.3–1.9)	1.3 (1.0–1.6)	702	37.9
Metro Area: Large Fringe Counties	264	4.9	0.7 (0.6–0.8)	1.0 (0.7–1.3)	1,173	21.8
Medium Metro Area	194	7.4	1.0	1.0	889	33.8
Small Metro Area	66	9.1	1.2 (0.9–1.6)	1.2 (0.9–1.6)	365	50.4
Metropolitan Counties	121	10.3	1.4 (1.1–1.8)	1.2 (0.9–1.5)	574	48.8
Non-core Counties	43	11.0	1.5 (1.1–2.1)	1.4 (1.0–1.9)	187	47.7
Unknown	10	5.8	0.8 (0.4–1.5)		24	13.8

* Adjusted ORs include adjustment for all other covariates

Table 2

Maternal Hospitalization for Depression within 12 Months of Delivery, New York City, 1995 – 2004, N = 900,726 Singleton Deliveries

Characteristic	Definite Postpartum Depression			Possible Postpartum Depression		
	Cases (N)	Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)	Cases (N)	Proportion (per 10,000)
All Deliveries	352	3.9			1,635	18.1
Year (delivery)						
1995 – 1998	106	2.9	1.0	1.0	583	16.1
1999 – 2001	98	3.7	1.3 (1.0–1.7)	1.4 (1.0–1.8)	459	17.4
2002 – 2004	148	5.4	1.8 (1.4–2.4)	2.0 (1.5–2.7)	593	21.6
Season (delivery)						
Spring (March – May)	88	4.0	1.0 (0.8–1.4)	1.1 (0.8–1.5)	400	18.0
Summer (June – August)	90	3.9	1.0	1.0	382	16.5
Fall (September – November)	85	3.7	1.0 (0.7–1.3)	1.1 (0.8–1.5)	426	18.7
Winter (December – February)	89	4.1	1.1 (0.8–1.4)	1.0 (0.7–1.4)	427	19.6
Maternal Age (years)						
<20	35	4.2	1.1 (0.8–1.5)	0.7 (0.4–1.0)	205	24.4
20 – 34	255	3.8	1.0	1.0	1,156	17.4
35+	62	4.1	1.1 (0.8–1.4)	1.1 (0.8–1.5)	274	18.0
Maternal Race/Ethnicity						
White non-Hispanic	62	2.9	1.0	1.0	247	11.5
Black non-Hispanic	121	6.2	2.2 (1.6–2.9)	1.7 (1.2–2.4)	529	27.2
Hispanic	77	4.3	1.5 (1.1–2.1)	1.2 (0.8–1.8)	452	25.3
Asian	15	2.0	0.7 (0.4–1.2)	0.7 (0.3–1.2)	51	6.8
Other	19	3.7	1.3 (0.8–2.1)	1.3 (0.8–2.3)	58	11.2
Unknown	58	3.1	1.1 (0.8–1.5)	1.1 (0.8–1.7)	298	15.9
Insurance Type						
Private	98	2.7	1.0	1.0	337	9.2
Public	246	5.0	1.9 (1.5–2.4)	1.7 (1.3–2.3)	1,231	24.8
Self-pay	7	2.2	0.8 (0.4–1.8)	0.7 (0.3–1.7)	64	20.2
Other	1	2.3	0.9 (0.1–6.1)	1.3 (0.2–9.7)	3	6.8
Parity						

Characteristic	Definite Postpartum Depression				Possible Postpartum Depression			
	Cases (N)	Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)	Cases (N)	Proportion (per 10,000)	OR (95% CI)	aOR* (95% CI)
0	183	4.5	1.0	1.0	618	15.2	1.0	1.0
1	75	2.8	0.6 (0.5–0.8)	0.6 (0.4–0.8)	424	15.6	1.0 (0.9–1.2)	1.0 (0.9–1.2)
2+	94	4.2	0.9 (0.7–1.2)	0.7 (0.5–0.9)	593	26.7	1.8 (1.6–2.0)	1.3 (1.1–1.5)
Unknown	0				0			
Nativity								
US born	180	4.4	1.0	1.0	1,024	25.1	1.0	1.0
Foreign born	171	3.5	0.8 (0.6–1.0)	0.8 (0.6–1.0)	592	12.2	0.5 (0.4–0.5)	0.5 (0.4–0.5)
Unknown	1	1.7	0.4 (0.1–2.7)		19	32.2	1.3 (0.8–2.0)	
Education (years)								
<12	109	5.0	1.3 (1.0–1.7)	1.2 (0.9–1.6)	634	28.9	1.7 (1.5–1.9)	1.3 (1.1–1.4)
12	117	3.9	1.0	1.0	527	17.5	1.0	1.0
13–16	98	3.6	0.9 (0.7–1.2)	1.0 (0.7–1.4)	382	14.1	0.8 (0.7–0.9)	1.0 (0.9–1.2)
17+	25	2.7	0.7 (0.4–1.1)	0.9 (0.6–1.5)	62	6.7	0.4 (0.3–0.5)	0.7 (0.5–0.9)
Unknown	3	2.1	0.5 (0.2–1.7)		30	20.9	1.2 (0.8–1.7)	
Prenatal Tobacco Use								
Non-smoker	318	3.7	1.0	1.0	1,414	16.3	1.0	1.0
Smoker	33	11.4	3.1 (2.2–4.5)	3.0 (2.0–4.5)	213	73.9	4.6 (3.9–5.3)	2.8 (2.4–3.3)
Unknown	1	2.7	0.7 (0.1–5.3)		8	21.8	1.3 (0.7–2.7)	
Start of Prenatal Care								
1st trimester	198	3.7	1.0	1.0	811	15.3	1.0	1.0
2nd trimester	81	3.9	1.1 (0.8–1.4)	1.0 (0.8–1.3)	403	19.6	1.3 (1.1–1.4)	1.1 (1.0–1.3)
3rd trimester	22	4.2	1.1 (0.7–1.7)	0.9 (0.6–1.5)	126	23.9	1.6 (1.3–1.9)	1.2 (1.0–1.5)
None	4	3.0	0.8 (0.3–2.2)	0.5 (0.2–1.7)	56	42.2	2.8 (2.1–3.6)	1.8 (1.4–2.4)
Unknown	47	4.8	1.3 (0.9–1.7)		239	24.2	1.6 (1.4–1.8)	

* Adjusted ORs include adjustment for all other covariates