Published in final edited form as:

J Abnorm Psychol. 2011 May ; 120(2): 483–489. doi:10.1037/a0021810.

Imagining the future: Degraded representations of future rewards and events in schizophrenia

Erin A Heerey¹, Tatyana M Matveeva², and James M Gold²

¹ School of Psychology, Bangor University, Bangor, UK

² Maryland Psychiatric Research Center, University of Maryland, Baltimore

Abstract

Over the course of life, most people work towards temporally distant rewards such as university degrees or work-related promotions. In contrast, many people with schizophrenia show deficits in behavior oriented toward long-term rewards, although they function adequately when rewards are more immediately present. Moreover, when asked about possible future events, individuals with schizophrenia show foreshortened future time perspectives relative to healthy individuals. Here, we take the view that these deficits are related and can be explained by cognitive deficits. We compared participants with schizophrenia (N=39) and healthy participants' (N=25) performance on tasks measuring reward discounting and future event representations. Consistent with previous research, we found that relative to healthy participants, those with schizophrenia discounted the value of future rewards more steeply. Furthermore, when asked about future events, their responses were biased toward events in the near future, relative to healthy participants' responses. Interestingly, although discounting and future representations were unrelated in healthy participants, we found significant correlations across the tasks among participants with schizophrenia, as well as correlations with cognitive variables and symptoms. Further analysis showed that statistically controlling working memory eliminated group differences in task performance. Together these results suggest that the motivational deficits characteristic of schizophrenia relate to cognitive deficits affecting the ability to represent and/or evaluate distant outcomes, a finding with important implications for promoting recovery from schizophrenia.

Keywords

Schizophrenia; reward; time perspectives; delay discounting; cognitive function

Despite decades of research, schizophrenia remains a mysterious illness, which is often characterized by significant disability (Lehman et al., 2004; Novick, Haro, Suarez, Vieta, & Naber, 2009). One of the major impediments to recovery for many individuals is a failure to sustain long-term patterns of goal-oriented behavior, especially when associated rewards are distant. For example, many people with schizophrenia find it difficult to maintain involvement with vocational rehabilitation programs that offer longer-term rewards (e.g., a pay-check after one or two weeks' work; Zito, Greig, Wexler, & Bell, 2007). In contrast,

Address for correspondence: Erin Heerey, Bangor University, School of Psychology, Brigantia Building, Bangor, Gwynedd, LL57 2AS, UK, Tel: +44(0)1248388804, e.heerey@bangor.ac.uk.

Publisher's Disclaimer: The following manuscript is the final accepted manuscript. It has not been subjected to the final copyediting, fact-checking, and proofreading required for formal publication. It is not the definitive, publisher-authenticated version. The American Psychological Association and its Council of Editors disclaim any responsibility or liabilities for errors or omissions of this manuscript version, any version derived from this manuscript by NIH, or other third parties. The published version is available at www.apa.org/pubs/journals/ABN

research shows that in situations such as token economies, in which behaviors and/or goals are rewarded on a daily or hourly basis, people with schizophrenia engage quite readily (Dickerson, Tenhula, & Green-Paden, 2005; Silverstein et al., 2009). Thus, the critical issue does not appear to be whether individuals with schizophrenia are sensitive to rewards per se, but rather that they have difficulty organizing and engaging in the behaviors required to obtain temporally distant rewards.

Accordingly, we have shown that in schizophrenia, rewards that are not immediately present in the environment quickly lose their ability to modulate behavioral responses and that the degree to which they do so correlates with working memory (Heerey & Gold, 2007). This idea suggests that individuals with schizophrenia may undervalue delayed relative to immediate rewards because of difficulties maintaining reward-value representations over time. Insofar as this ability is compromised, individuals should prefer the more clearly represented immediate choices. Indeed, we have reported that individuals with schizophrenia discount or devalue temporally distant rewards to a greater degree than do healthy individuals (Heerey, Robinson, McMahon, & Gold, 2007). Notably, although steeper discounting is usually interpreted as reflecting impulsivity (Petry, 2001; Reynolds, 2006), cognitive impediments, including introducing working memory loads, also degrade it (Hinson, Jameson, & Whitney, 2003).

Interestingly, the ability to represent future time, which is impaired in schizophrenia (e.g., Eack, George, Prasad, & Keshavan, 2008; Suto & Frank, 1994; Wallace, 1956), may also hinge upon the ability to maintain and utilize abstract representations (Klapproth, 2008), and therefore be influenced by working memory-related processes (Addis, Wong, & Schacter, 2007). Thus, according to this formulation, reduced preferences for distant rewards in schizophrenia should relate to foreshortened future time perspectives because events further in the future (e.g., distant rewards) are poorly represented relative to more proximal events (e.g., immediate rewards).

Notably, this prediction differs from the pattern observed in individuals with orbitofrontal cortex (OFC) lesions. Specifically, Fellows and Farah (2005) demonstrated that people with OFC injuries had impaired future time perspectives but did not discount delayed rewards more steeply than did other lesion groups or healthy participants. Thus, discounting and time perspectives were dissociable in the OFC group. In contrast, we expect a relationship between discounting and time perspectives in schizophrenia because we think that global impairments in the ability to represent non-immediate experiences, including distant rewards and events, may underpin both deficits.

The goals of this research were to: 1) replicate previous findings showing increased delay discounting (Heerey, et al., 2007) and foreshortened future time perspectives in schizophrenia (Wallace, 1956); 2) understand the relationship between delay discounting and future time perspectives in schizophrenia; and 3) link time perspectives and discounting to symptoms, cognitive ability and functional outcomes. We predict that in participants with schizophrenia, time perspectives and discounting will be correlated and will relate to cognition, symptoms and functioning.

Methods

Participants

Participants included 39 outpatients with schizophrenia (SC) and 25 healthy comparison (HC) participants. Schizophrenia diagnoses were confirmed with the Structured Clinical Interview for DSM-IV (SCID, First, Spitzer, Miriam, & Williams, 2002). SC participants were deemed clinically stable by their clinicians and received stable diagnoses of

antipsychotic medicine with no medication changes for >4 weeks before participation. Symptom and functioning assessments included the Scale for the Assessment of Negative Symptoms (SANS, Andreasen, 1989), the Brief Psychiatric Rating Scale (BPRS, Overall & Gorham, 1962), and the Level of Functioning Scale (LoFS, Strauss & Carpenter, 1972). HC participants were receiving no psychiatric medications and were confirmed free of psychiatric diagnoses based on SCID screening. Potential participants were excluded if there was evidence of neurological injury or disorder, substance abuse/dependence other than nicotine within the 4 months before study enrolment, or other disorder capable of affecting task performance. All participants gave written informed consent. The University of Maryland Institutional Review Board approved the study. Participant characteristics appear in Table 1.

Procedures

Participants completed two tasks in the context of a larger study. The first was a computerized delay-discounting task, measuring the degree to which rewards lose their value as they become further in the future. In delay-discounting tasks, participants choose between smaller, more immediate rewards and larger rewards at later times (e.g., \$10 today or \$20 in 28 days). Discounting is estimated according to the hyperbolic discounting

equation, $V = \frac{A}{1+kD}$, in which V is a future reward's discounted value, A is its undiscounted or immediate value, D is the delay interval and k is the discounting parameter, which varies according to how quickly rewards are discounted (Kirby, 2000). For example, assuming a k -value of .01 and a delay of 28 days, \$20 would have a discounted value of \$15.63. Higher k -values indicate steeper/faster discounting.

We have shown that SC participants discount distant rewards more steeply, which may indicate impulsivity. We therefore included a condition in which the smaller, sooner reward was not immediate. That is, on some trials of the task, participants were asked whether they preferred, for example, a smaller reward in 7 days or a larger reward in 14 days. This procedure, known as subadditive delay-discounting (Read & Roelofsma, 2003), equates the amount of time that lapses between the decision-point and the proximal reward and the proximal and distal rewards (delay intervals ranged from a minimum of 5 days to a maximum of 6 months). Theoretically, subadditive discounting procedures reduce impulsive choice by enforcing patience (i.e., a delay before reward receipt Green, Myerson, & Macaux, 2005; Read, 2001), as research in both humans (Kirby & Herrnstein, 1995) and animals shows (Ainslie, 1974). We therefore reasoned that a group-by-condition interaction, in which SC participants showed normal subadditive discounting but steeper immediate discounting would demonstrate that their immediate decisions were indeed more impulsive, whereas no interaction would point away from simple impulsivity as a factor in the SC deficit.

On each trial of the task, participants chose between a smaller, sooner reward (SSR) and a larger, later reward (LLR) using a key-press. Participants were told that they would not actually receive the rewards they chose but were nonetheless asked to choose as if each trial was a choice between genuine rewards. We used this procedure because many previous studies have failed to find differences between hypothetical and real rewards (e.g., Bickel, Pitcock, Yi, & Angtuaco, 2009; Lagorio & Madden, 2005; Madden, Begotka, Raiff, & Kastern, 2003).

In the immediate condition, participants chose between an SSR with no delay ("today") and an LLR available after a variable number of days. In the subadditive condition, they chose between rewards in which the delay between the SSR and LLR was the same as that between "today" and the SSR. To determine the point of subjective equality (PSE) for each

condition, participants were presented with choices at values of k ranging between 0 and 1, using separate adaptive “staircases” in which the k -values started at 1 and moved, step-by-step, toward 0 or vice versa, allowing us to gradually increase the precision of our k -estimates. We used two staircases per condition (one starting at $k=0$ and the other at $k=1$), each of which provided an independent estimate of PSE by reversing direction each time participants switched between choosing the SSR and the LLR (e.g., if k was increasing before the preference shift, it now decreased). For example, one staircase started at $k=0$ (no discounting—the SSR and the LLR held the same monetary value, e.g., “\$34 today or \$34 in 60 days”), meaning that participants should choose the SSR. On this staircase, k increased by .10 on each successive trial until participants chose the LLR, at which point k began to decrease, until there was another preference shift. The step size decreased by 20% with each preference shift (e.g., .10, .08, .064), allowing us to narrow the window within which participants’ k -estimates fell. There were 20 trials per staircase. Trials presenting choices from each of the four staircases were randomly intermixed. This procedure has three advantages over procedures using fixed k -intervals (e.g., Kirby, Petry, & Bickel, 1999). First, it allows precise estimates of discounting parameters by testing increasingly narrow k -intervals. Second, it identifies participants who adopt idiosyncratic response strategies such as choosing only the SSR. Third, this procedure is robust against inconsistent responding (e.g., accidentally pressing the wrong button), which is an issue for SC participants and can produce artificially inflated/deflated k -estimates.

Finally, our LLR values ranged from \$75 to \$85 as we have found that SC participants discount LLRs in this value range much more steeply than do HC participants. This meant that our SSR values had a large range – from \$0 to \$85.

The second task participants completed was a future time perspective interview, designed to produce an estimate of the average time window within which participants view future events. Using procedures adapted from Fellows and Farah (2005) and Wallace (1956), participants listed ten events that they believed would occur in their futures using a standard probe (“I’d like you to spend some time thinking about your own future. Please think of ten events that may happen to you during the rest of your life.”). Participants responded verbally and responses were recorded verbatim. Participants received general encouragement after each item and were encouraged to “keep thinking” until they had generated ten events. All participants except one SC participant generated 10 events. The experimenter then reviewed events one-at-a-time and asked participants to estimate how far in the future they thought the event would occur. To estimate average time horizon, we calculated the length of time between the date of the interview and the estimated time of each event’s occurrence in months and took the mean time estimate the events. We also report the maximum “extension” of participants’ time horizons (i.e., the most distant event).

In addition to the tasks above, participants completed the letter-number sequencing and Hopkins Verbal Learning tests to measure verbal working memory and verbal learning as recommended in the MATRICS consensus cognitive battery (<http://www.matrics.ucla.edu>), and the Wechsler Test of Adult Reading (Wechsler, 2001) to estimate current/premorbid function.

Data analysis

To estimate k in both discounting conditions (immediate, subadditive), we computed the geometric mean of the final k -values of the two staircases measuring that condition. There were 3 participants (2 SC, 1 HC) for whom k could not be estimated because they never varied their choices. These participants were excluded from discounting analyses. The log-transformed k -values were analyzed with a mixed model ANOVA with group (HC, SC) as the between-subjects variable and condition (immediate, subadditive) as the within-subjects

variable. Time horizons were not normally distributed even after log transformation, so group comparisons and correlations used non-parametric analyses. Post-hoc analyses use Bonferroni's correction for multiple comparisons.

Results

Delay Discounting

Figure 1 shows participants' estimated k -values in immediate and subadditive conditions. As predicted, we found a main effect of group such that SC participants showed steeper discounting than did HC participants, $F(1, 59) = 5.03$; $p = .03$; $\eta_p^2 = .08$, replicating our previous findings. There was also a strong main effect for condition such that all participants benefited in the subadditive condition, $F(1, 59) = 96.41$; $p < .001$; $\eta_p^2 = .62$. That is, discounting was less steep when choosing between two delayed choices rather than an immediate and a delayed choice. Post-hoc analyses showed that SC participants discounted more steeply in both immediate ($p = .04$) and subadditive conditions ($p = .02$), meaning that the introduction of a delay before the SSR improved discounting in both groups similarly. There was no group-by-condition interaction, $F(1, 59) < .01$; $p = .99$, suggesting that SC participants' steeper immediate discounting cannot simply be attributed to impulsivity.

Future Time Perspectives

Two coders (blind to diagnosis) classified events into a set of categories including household chores (e.g., spring cleaning), health-related events (e.g., potential illness/relapse, attending a swimming class), leisure (e.g., seeing a film, vacation), financial (e.g., reaching a savings target), religion (e.g., attend prayer group), hobbies/interests (e.g., photography class), and family events (e.g., nephew's graduation) (Cohen's k appas $> .72$). Interestingly, there were no differences in the frequency with which participants reported events across categories (p -values $> .17$). However, as anticipated, SC participants reported events that were both on average, Mann-Whitney $U = 292$, $p = .008$, and maximally, Mann-Whitney $U = 354$, $p = .036$, nearer in the future than HC participants' events. One reason that SC participants' report nearer-future events might be that their reports are weighted toward events that repeat daily/weekly and are therefore easily reported (e.g., "eat supper tonight", reported by 1 HC and 4 SC participants; "clean the house", reported by 2 HC and 5 SC participants). If this explanation is the cause of the time-horizon differences, participants' event distributions should be similar after removing immediate events. Alternatively, if the distributions are still different after immediate-event removal, then the cause of the difference cannot be simple ease of reporting. To examine this idea, we tallied events falling into 10 time bins for both groups (Figure 2). A χ^2 goodness-of-fit test confirmed that in general, SC participants' events were distributed differently than HC participants' events, $\chi^2 = 37.95$; $df = 9$; $p < .001$. To examine the degree to which this pattern held for non-immediate events, we did the same analysis excluding events in time bins one week or sooner in the future. Results again showed that SC participants' events were nearer than were HC participants' events, $\chi^2 = 23.24$; $df = 5$; $p < .001$. These results mirror the discounting results and suggest that SC participants' reports are biased toward more proximal events.

To support this idea, we correlated the log-transformed, average k -value with average log-transformed time horizon. Among HC participants, there was no relationship between discounting and time horizon, $r = .06$; $p = .76$. However, among SC participants we found a significant relationship between average time horizon and discounting, $r = -.35$; $p = .04$, such that longer time horizons correlated with less discounting impairment.

Cognitive and Symptom Predictors of Task Performance

To better understand the factors contributing to the group differences in both tasks together, we computed a composite score by converting the log-transformed average time horizon and the log-transformed average k -value to z -scores and averaging these. Among HC participants, the composite 'future-representation' score correlated with working memory, $\rho_{\text{Spearman}} = .44$; $p = .02$, and verbal learning, $\rho_{\text{Spearman}} = .40$; $p = .04$ and among SC participants, future-representation related to working memory, $\rho_{\text{Spearman}} = .45$; $p = .003$, but not verbal learning, $\rho_{\text{Spearman}} = .23$; $p = .15$, suggesting that better cognitive resources correspond to better representations of future time and events. Current/premorbid verbal function did not correlate with future-representation in either group (p -values $> .23$).

To better understand how cognitive differences underpin group differences in task performance, we conducted a hierarchical multiple regression with the composite future-representation score as the dependent variable and entered the cognitive variables (working memory and verbal learning) at step 1 and diagnostic status (0=HC; 1=SC) at step 2 as predictors. At step 1, working memory significantly predicted task performance, $\beta = .39$; $t(59) = 2.88$; $p = .005$, but verbal learning did not, $\beta = .19$; $t(59) = 1.36$; $p = .18$. However, after controlling for cognitive performance, the group difference at step 2 was no longer significant, $\Delta R^2 = .002$; $p = .70$.

Finally, we examined the relationships between the composite future-representation score, general functioning, positive symptoms and avolition/anhedonia in the SC group. Future-representation correlated with LoFS work-functioning, $\rho_{\text{Spearman}} = .33$; $p = .04$, and tended to relate to social-functioning, $\rho = .27$; $p = .10$, but did not correlate with positive symptoms, $\rho_{\text{Spearman}} = -.04$; $p = .82$. To examine avolition/anhedonia, we created a composite score using the avolition global score and the anhedonia global score from the SANS. Future representations correlated with avolition/anhedonia, $\rho_{\text{Spearman}} = -.32$, $p = .05$, such that participants with more avolition/anhedonia showed reduced performance.

Discussion

As anticipated, results showed that SC participants had foreshortened time perspectives and discounted rewards more steeply than did HC participants. We also found that these processes were related in SC but not HC participants, such that SC participants with shorter average time horizons valued future rewards less. After controlling for cognitive performance, specifically working memory, the group differences in task performance disappeared. These findings hint at the possibility that a common factor underpins both deficits. What might this factor be? We suggest two possibilities.

The most obvious suggestion for the relationship between discounting and time horizons in SC participants is that they are more impulsively reward-focused than HC participants. High levels of reward-focus reduce ability to attend to future rewards (Mitchell, Fields, D'Esposito, & Boettiger, 2005; Perry & Carroll, 2008) and to the future more generally (Bechara, Dolan, & Hindes, 2002). Although previous research concludes that schizophrenia is associated with impulsive decision-making (Enticott, Ogloff, & Bradshaw, 2008), the present results do not support this view. If the subadditive condition had normalized SC participants' discounting, this would have suggested that poor delay tolerance or impulsivity was responsible for the result (Ainslie, 1974; Kirby & Herrnstein, 1995). However, the delay enhanced both groups' performance equally. Moreover, avolition and anhedonia correlated with task performance, suggesting that simple impulsiveness is not sufficient to explain these results.

Alternatively, we argue that cognitive deficits present in the SC group may adversely affect the ability to represent both future time and future rewards. We found a relationship between working memory and performance in both tasks such that participants with better working memory showed less task-related impairment (for similar results see, Strauss et al., In Press). Working memory, which is impaired in schizophrenia (Barch, 2005), may support the integration of cognitive with affective information (Becerril & Barch, 2010), and serve to maintain reward representations over time. This idea is in keeping with other research showing deficits in the ability to represent future rewards in schizophrenia (Gard, Germans Gard, Kring, & John, 2006). Indeed, the results of our regression analysis suggest that working memory deficits may underpin difficulties in representing future rewards and events, which in turn may have functional consequences (Gard, Fisher, Garrett, Genevsky, & Vinogradov, 2009). Indeed, SC participants' work-related functioning declined as their task performance worsened.

Interestingly, the future-representation and negative symptom and cognitive correlations in the present study diverge slightly from our previously reported results (Heerey, et al., 2007). In that study, surprisingly, we found that more negative symptoms tended to predict less impaired discounting. In the present study, we found the opposite relationship. Specifically, increased negative symptoms correlated with more impaired future representations. This discrepancy may relate to SC participants' symptom-levels, which were lower in the present study, suggesting that the current participants suffered from fewer symptoms at the time of testing than did our previous participants. Thus, the present research may have under-sampled participants with extremely high negative symptom levels. Additionally, our previous report showed that verbal learning and spatial working memory predicted task performance and in the present study, verbal working memory alone accounted for the group differences in future-representation ability, although we note that the verbal learning correlation tends toward this without reaching statistical significance.

There are two important caveats to these findings. First, all of our SC participants were receiving antipsychotic medicine. We therefore remain agnostic about whether the present results are characteristic of schizophrenia or its treatment. Second, participants completed the task for hypothetical rewards. In some ways, the use of hypothetical rewards mirrors life in that people make decisions about future rewards that are only imagined until they actually occur (e.g., investment dividends). Although this may alter decision-making relative to tasks involving tangible rewards, especially among SC participants, it is also more compatible with the future time perspectives task in which participants speculated about possible but not certain future events. Regardless, these findings have consequences for understanding some of the motivational and functional deficits characteristic of the illness. Specifically, they suggest that treatment programs promoting functional recovery should place emphasis on short- and medium-term goals with gradual shifts to organizing and working toward longer-term outcomes. Interestingly, these results are fully compatible with the effectiveness of token economy treatment approaches documented 40 years ago (Lloyd & Abel, 1970). This suggests that the challenges faced by people with schizophrenia when tight contingency control is reduced must be replaced by the ability to produce and utilize self-generated mental representations to drive the push toward future goals.

The power of distant rewards to drive behavior is an important element in functional outcomes. In summary, we found that participants with schizophrenia showed foreshortened future time perspectives and steeper discounting, which related to working memory deficits. These results are important because they suggest that cognitive deficits may underpin the ability to represent future time and rewards, thereby impinging upon functional adjustment such as engagement with work and the pursuit of pleasurable activities. Thus, at least among

medicated individuals, improving the ability to look toward and act upon future rewards may improve functional outcomes.

Acknowledgments

Funding for this work was provided by NIMH Grants R01-MH080066 and R24-MH72647. We thank Sharon August, Kimberley Bell-Warren and Benjamin Robinson for their assistance with data collection. All authors declare that they have no conflicts of interest to report.

References

- Addis DR, Wong AT, Schacter DL. Remembering the past and imagining the future: common and distinct neural substrates during event construction and elaboration. *Neuropsychologia*. 2007; 45(7): 1363–1377. [PubMed: 17126370]
- Ainslie G. Impulse control in pigeons. *Journal of the Experimental Analysis of Behavior*. 1974; 21:485–489. [PubMed: 16811760]
- Andreasen NC. The Scale for the Assessment of Negative Symptoms (SANS): conceptual and theoretical foundations. *Br J Psychiatry Suppl*. 1989; (7):49–58. [PubMed: 2695141]
- Barch DM. The cognitive neuroscience of schizophrenia. *Annu Rev Clin Psychol*. 2005; 1:321–353. [PubMed: 17716091]
- Becerril K, Barch D. Influence of Emotional Processing on Working Memory in Schizophrenia. *Schizophr Bull*. 2010
- Bechara A, Dolan S, Hindes A. Decision-making and addiction (part II): myopia for the future or hypersensitivity to reward? *Neuropsychologia*. 2002; 40(10):1690–1705. [PubMed: 11992657]
- Bickel WK, Pitcock JA, Yi R, Angtuaco EJ. Congruence of BOLD response across intertemporal choice conditions: fictive and real money gains and losses. *J Neurosci*. 2009; 29(27):8839–8846. [PubMed: 19587291]
- Dickerson FB, Tenhula WN, Green-Paden LD. The token economy for schizophrenia: review of the literature and recommendations for future research. *Schizophr Res*. 2005; 75(2-3):405–416. [PubMed: 15885531]
- Eack SM, George MM, Prasad KM, Keshavan MS. Neuroanatomical substrates of foresight in schizophrenia. *Schizophr Res*. 2008; 103(1-3):62–70. [PubMed: 18603414]
- Enticott PG, Ogloff JR, Bradshaw JL. Response inhibition and impulsivity in schizophrenia. *Psychiatry Res*. 2008; 157(1-3):251–254. [PubMed: 17916385]
- Fellows LK, Farah MJ. Dissociable elements of human foresight: a role for the ventromedial frontal lobes in framing the future, but not in discounting future rewards. *Neuropsychologia*. 2005; 43(8): 1214–1221. [PubMed: 15817179]
- First, MB.; Spitzer, RL.; Miriam, G.; Williams, JBW. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition With Psychotic Screen. New York, NY: Biometrics Research, New York State Psychiatric Institute; 2002.
- Gard DE, Fisher M, Garrett C, Genevsky A, Vinogradov S. Motivation and its relationship to neurocognition, social cognition, and functional outcome in schizophrenia. *Schizophr Res*. 2009; 115(1):74–81. [PubMed: 19783407]
- Gard DE, Germans Gard MK, Kring AM, John OP. Anticipatory and consummatory experience of pleasure: A scale development study. *Journal of Research in Personality*. 2006; 40:1086–1102.
- Green L, Myerson J, Macaux EW. Temporal discounting when the choice is between two delayed rewards. *J Exp Psychol Learn Mem Cogn*. 2005; 31(5):1121–1133. [PubMed: 16248754]
- Heerey EA, Gold JM. Patients with schizophrenia demonstrate dissociation between affective experience and motivated behavior. *J Abnorm Psychol*. 2007; 116(2):268–278. [PubMed: 17516760]
- Heerey EA, Robinson BM, McMahon RP, Gold JM. Delay discounting in schizophrenia. *Cogn Neuropsychiatry*. 2007; 12(3):213–221. [PubMed: 17453902]
- Hinson JM, Jameson TL, Whitney P. Impulsive decision making and working memory. *J Exp Psychol Learn Mem Cogn*. 2003; 29(2):298–306. [PubMed: 12696817]

- Kirby, KN. Instructions for inferring discount rates from choices between immediate and delayed rewards. Williams College; 2000. p. 7
- Kirby KN, Herrnstein RJ. Preference reversals due to myopic discounting of delayed reward. *Psychological Science*. 1995; 6(2):83–89.
- Kirby KN, Petry NM, Bickel WK. Heroin addicts have higher discount rates for delayed rewards than non-drug-using controls. *J Exp Psychol Gen*. 1999; 128(1):78–87. [PubMed: 10100392]
- Klapproth F. Time and decision making in humans. *Cogn Affect Behav Neurosci*. 2008; 8(4):509–524. [PubMed: 19033245]
- Lagorio CH, Madden GJ. Delay discounting of real and hypothetical rewards III: steady-state assessments, forced-choice trials, and all real rewards. *Behav Processes*. 2005; 69(2):173–187. [PubMed: 15845306]
- Lehman AF, Kreyenbuhl J, Buchanan RW, Dickerson FB, Dixon LB, Goldberg R, et al. The Schizophrenia Patient Outcomes Research Team (PORT): updated treatment recommendations 2003. *Schizophr Bull*. 2004; 30(2):193–217. [PubMed: 15279040]
- Lloyd KE, Abel L. Performance on a token economy psychiatric ward: a two year summary. *Behav Res Ther*. 1970; 8(1):1–9. [PubMed: 5422892]
- Madden GJ, Begotka AM, Raiff BR, Kastern LL. Delay Discounting of Real and Hypothetical Rewards. *Experimental and Clinical Psychopharmacology*. 2003; 11(2):139–145. [PubMed: 12755458]
- Mitchell JM, Fields HL, D'Esposito M, Boettiger CA. Impulsive responding in alcoholics. *Alcohol Clin Exp Res*. 2005; 29(12):2158–2169. [PubMed: 16385186]
- Novick D, Haro JM, Suarez D, Vieta E, Naber D. Recovery in the outpatient setting: 36-month results from the Schizophrenia Outpatients Health Outcomes (SOHO) study. *Schizophr Res*. 2009; 108(1-3):223–230. [PubMed: 19070991]
- Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. *Psychological Reports*. 1962; 10:790–812.
- Perry JL, Carroll ME. The role of impulsive behavior in drug abuse. *Psychopharmacology (Berl)*. 2008; 200(1):1–26. [PubMed: 18600315]
- Petry NM. Pathological gamblers, with and without substance use disorders, discount delayed rewards at high rates. *J Abnorm Psychol*. 2001; 110(3):482–487. [PubMed: 11502091]
- Read D. Is time-discounting hyperbolic or subadditive? *Journal of Risk and Uncertainty*. 2001; 23(1): 5–32.
- Read D, Roelofsma PHMP. Subadditive versus hyperbolic discounting: A comparison of choice and matching. *Organizational Behavior and Human Decision Processes*. 2003; 91:140–153.
- Reynolds B. A review of delay-discounting research with humans: relations to drug use and gambling. *Behav Pharmacol*. 2006; 17(8):651–667. [PubMed: 17110792]
- Silverstein SM, Spaulding WD, Menditto AA, Savitz A, Liberman RP, Berten S, et al. Attention shaping: a reward-based learning method to enhance skills training outcomes in schizophrenia. *Schizophr Bull*. 2009; 35(1):222–232. [PubMed: 18212327]
- Strauss GP, Robinson BM, Waltz JA, Frank MJ, Kasanova Z, Herbener ES, et al. Patients With Schizophrenia Demonstrate Inconsistent Preference Judgments for Affective and Nonaffective Stimuli. *Schizophr Bull*. In Press.
- Strauss JS, Carpenter WT Jr. The prediction of outcome in schizophrenia. I. Characteristics of outcome. *Arch Gen Psychiatry*. 1972; 27(6):739–746. [PubMed: 4637891]
- Suto M, Frank G. Future time perspective and daily occupations of persons with chronic schizophrenia in a board and care home. *Am J Occup Ther*. 1994; 48(1):7–18. [PubMed: 8116786]
- Wallace M. Future time perspective in schizophrenia. *Journal of Abnormal Social Psychology*. 1956; 52:240–245.
- Wechsler, D. Wechsler Test of Adult Reading. San Antonio, TX: The Psychological Corporation; 2001.
- Zito W, Greig TC, Wexler BE, Bell MD. Predictors of on-site vocational support for people with schizophrenia in supported employment. *Schizophr Res*. 2007; 94(1-3):81–88. [PubMed: 17512172]

Figure 1. Discounting constants for HC (gray bars) and SC participants (white bars) in the immediate and subadditive conditions. Error bars show +/-1 standard error of the mean.

Figure 2. Percentage of responses across time for HC participants (dashed line) and SC participants (solid line). Note that time is shown on a logarithmic scale.

Table 1
Participant Characteristics

	Healthy Comparison Participants	Participants with Schizophrenia	p-value
N	25	39	---
Age	47.00 (10.18)	45.16 (10.03)	.47
Age at illness onset	---	23.41 (7.56)	---
Education	14.29 (1.98)	12.23 (1.94)	<.001
Paternal Education	12.42 (4.31)	12.50 (4.23)	.94
Gender (male:female) *	17:8	31:8	.36
Race *			.69
African American	9	11	
Caucasian	16	26	
Other	0	2	
Cognitive Performance			
Working Memory	16.15 (3.92)	11.38 (3.18)	<.001
Verbal Learning	28.46 (3.95)	20.43 (4.99)	<.001
Test of Adult Reading	106.04 (15.90)	94.77 (17.05)	.01
Antipsychotic medicine			
Atypical	---	32	---
Typical	---	6	---
Both	---	1	---
Clinical Symptoms			
SANS Avolition/Anhedonia	---	3.49 (1.85)	---
BPRS Total	---	27.76 (7.53)	---
Level of Functioning Scales			
Social	---	3.54 (2.53)	---
Work	---	2.79 (2.63)	---

Note: Table displays means (standard deviations). Except where noted, group differences were tested with t-tests.

* Group differences tested with chi-square