Two non-histone proteins are associated with the promoter region and histone H1 with the transcribed region of active *hsp-70* genes as revealed by UV-induced DNA – protein crosslinking *in vivo*

Sergei V.Belikov, Alexander I.Belgovsky, Olga V.Preobrazhenskaya, Vadim L.Karpov and Andrei D.Mirzabekov*

W.A.Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow 117984, Russia

Received April 27, 1992; Revised and Accepted January 11, 1993

ABSTRACT

We described here an approach for mapping proteins on any sequence of genomic DNA. UV-induced DNA – protein crosslinking within whole cells and the ‘protein image’ hybridization technique (1) were applied to test the proteins bound to different regions of the *D.melanogaster hsp-70* gene. The histone H1-DNA association with the coding region is shown to be maintained, even during very intensive transcription, but is absent in the promoter. Two non-histone proteins with apparent molecular masses of 50 kD (p50) and 100 kD (p100) are crosslinked only to the active *hsp-70* gene regulatory region and preferentially bind to its complementary and coding DNA strands, respectively.

INTRODUCTION

Regulation of gene activity is a result of the intricate interactions of several proteins with regulatory gene elements (2,3). Histone H1 could be involved in the regulation of transcription (4,5). The use of different techniques, such as DNaseI footprinting, Exo III protection mapping, sequence specific oligonucleotide affinity chromatography and gel retardation assay provide some important information on the role and mechanism of the interactions of such proteins with DNA *in vitro* or in isolated nuclei *in situ*. However regulation of gene activity consists of complicated and subtle processes that are affected by many factors and therefore may differ in some essential details in cells, isolated nuclei or in solution. The use of DNA – protein crosslinking within whole cells with subsequent analyses of proteins crosslinked to specific sequences provides a useful approach that evades these difficulties.

We applied here UV-crosslinking in whole cells together with the ‘protein image’ hybridization method to identify *in vivo* the proteins associated with the regulatory elements and demonstrate

the presence of histone H1 on the coding sequences of highly transcribed *hsp-70* gene.

MATERIALS AND METHODS

D.melanogaster tissue culture cells, subline K, were grown at 26°C in Eschaler medium (6) supplemented with 10% foetal calf thymus serum, up to a density of $6-8 \times 10^6$ cells per ml. For heat shock, the cell suspension was incubated at 37°C for 30 min and quickly chilled to 0°C. Tissue culture cells in the medium were transferred to a jacketed dish and irradiated by UV-light, as described by Gilmour and Lis (7). The source of UV-light was a Chromato-Vue transilluminator, model C-61 (Ultra-Violet Products, USA) without filter, placed 10 cm above the cell suspension, which provided approximately $350 \mu\text{W}/\text{cm}^2$ light intensity. The cell suspension (0.5 cm in depth) was irradiated for 12 min at 0°C. Nuclei were isolated from irradiated cells as described (7) then lysed in 0.15 M NaCl, 10 mM Tris-HCl (pH 8.0), 1 mM EDTA, 0.1% NP-40, 2% lauryl sarcosinate, sonicated to DNA size of about 5–10 kilobases and the material was loaded onto a preformed CsCl gradient (7). Centrifugation was carried out at $200,000 \times g$ in a Ti-50.2 fixed-angle rotor (Beckman) for 40 hr at 16°C. The DNA-containing fractions were collected and dialysed against 0.25% NP-40, 10 mM Tris-HCl (pH 8.0). After dialysis, MgCl_2 and DNase I were added (to 5 mM and to $0.01 \mu\text{g}$ per $50 \mu\text{g}$ of DNA, respectively) for random fragmentation of DNA. The digestion was carried out for 7–10 min at 37°C, according to preliminary analytical experiments to get DNA fragments about 200–500 nucleotides long. The reaction was stopped by addition of EDTA to 10 mM.

Most of the uncrosslinked DNA was removed by phenol-chlorophorm extraction as in (8). Two-dimensional diagonal gel electrophoresis of the DNA – protein complexes was done as described (8,9) to obtain ‘protein images’. The DNA fragments

* To whom correspondence should be addressed

Figure 1. 'Protein image' hybridization for active *hsp-70* gene (probes A and B) and nontranscribed type II ribosomal insertion (D). Electrophoresis was carried out in the 1st dimension from left to right, and in the 2nd dimension from top to bottom. The diagonal stripes represent uncrosslinked DNA (0), 'images' of histone HI (HI), p50 (50) and p100 (100). The vertical line is the internal standard DNA (M), applied onto the gel of second dimension. A map of *D.melanogaster hsp-70* (proximal, locus 87A) is shown to the right. Restriction sites are designated by the capital letters: X -Xho I, S -Sal I, B -Bam HI. Figures on the blot indicate the size of single-stranded DNA.

were transferred onto a Hybond nylon membrane (Amersham). The conditions for DNA labeling, hybridization and washing of the filter were as described (10). One-dimensional densitometric scanning of autoradiographs of the blots was performed as in (7).

RESULTS

The *D.melanogaster* tissue culture cells in medium were UV-irradiated to initiate low-level DNA-protein crosslinking, the isolated nuclei were lysed, then DNA and the crosslinked DNA-protein complexes were purified by centrifugation in a CsCl gradient. The DNA-containing fractions were collected, the DNA was fragmented randomly by DNase I. Most of the uncrosslinked DNA was removed by phenol-chloroform extractions. The enriched DNA-protein complexes were subjected to 'protein image' hybridization analysis (1,8,9) as follows. Upon gel electrophoresis in the first dimension, DNA and DNA-protein complexes were separated according to mass, the crosslinked proteins retarding the running of the bound DNA fragments. The proteins were then degraded by Pronase directly in the gel, and the DNA fragments were separated in the second dimension. The DNA fragments were electroblotted to a nylon membrane and repeatedly hybridized with several cloned ³²P-labeled specific gene probes. The additional diagonals (besides that of the uncrosslinked DNA) corresponded to the DNA to which the proteins had been initially bound and the inclinations of these diagonals were proportional to the molecular masses of the crosslinked proteins. Some model experiments with the individual proteins of known molecular masses were carried out

Figure 2. 'Protein image' hybridization for heat shock activated (HS) and inactive control (C) *hsp-70* gene (probe A, coding strand). Gels of the first and second dimension electrophoresis contained 8% acrylamide. For details, see Fig. 1.

Figure 3. 'Protein image' hybridization for active *hsp-70* gene. Probe A, coding strand (a) and probe A, complementary strand (b). Gels of the first and second dimensional electrophoresis contained 6% acrylamide. For details see Fig. 1.

with the purpose to estimate the apparent molecular masses of crosslinking proteins (accuracy of our method was $\pm 10\%$ for the proteins with molecular masses less than 40 kD and $\pm 15\%$ for the larger proteins). σ -subunit of RNA-polymerase from *E.coli* (90 kD), bovine serum albumine (67 kD), DNA-topoisomerase II from cord seedling (135 kD), histone HI from chicken erythrocytes (37 kD) were chemically crosslinked to fragmented DNA and separated by two-dimensional gel electrophoresis in the same conditions as for the experimental samples. Unfractionated reference DNA (as an internal standard) from irradiated cells was loaded on the second-dimension gel.

Firstly, this gives us a possibility to compare on the same blot the intensities of diagonals on autoradiographs of different hybridizations.

And secondly, this guarantees against any fortuitous underrepresentations of any particular DNA region of any gene occurs (in the course of cross-linking, fractionation, transfer, etc.). For details see ref.(8).

We investigated the *D.melanogaster hsp-70* gene from cytogenic locus 87A (proximal) (11), using as hybridization probes the following cloned sequences (Fig. 1).

Probe A (177 b.p.-long) is present only in locus 87A genes contains the nuclease-hypersensitive promoter region (12).

Probe B (900 b.p.-long) is a part of the *hsp-70* transcription unit.

Probe D is the one for inactive chromatin. It is a 670-b.p. long fragment of *D.melanogaster* ribosomal type II insertion (13). This sequence is repeated 35 times in the genome, the level of its transcription is negligible (14).

The results of UV-crosslinking in heat-shocked *Drosophila* cells are presented in Fig. 1. After hybridization with probe D for inactive chromatin, the 'image' of histone H1 was revealed, along with the diagonal of uncrosslinked DNA. The attribution was carried out by crosslinking of individual proteins to DNA (data not shown). The core histone crosslinking was negligible under chosen irradiation exposures. Hybridization of the same filter with the probe for the coding region of *hsp-70* genes (probe B) did not led to any essentially different crosslinking of histone H1 to actively transcribed sequences as compared with nontranscribed ones. The intensities of the diagonals after hybridizations with Probe B and Probe D were very similar for both DNA strands.

The hybridization with probe A (coding strand) showed that the promoter region of *hsp-70* was free of histone H1. The appearance of two additional diagonals witnesses for the crosslinking of two proteins with apparent molecular masses of 50 kD (p50) and 100 kD (p100), respectively.

It was of interest to know whether or not the proteins found in the promoter region of activated *hsp-70* genes can be crosslinked to inactive genes. The cell culture was halved and one part was heat-shocked. The heat-shocked and control cells were UV-irradiated, the DNA-protein complexes isolated, enriched and loaded onto neighbouring tracks of the first dimension gel. Electrophoresis in the second dimension was performed in the same block of PAGE and after electroblotting the filter was hybridized with probe A (coding strand). The proteins discovered were crosslinked to the promoter only of the heat shock-activated gene (Fig. 2).

The hybridization with different ³²P-labeled strands of probe A, cloned in M13 phage, shows that p100 predominantly binds to the coding strand, and p50 prefers the complementary one, whereas histone H1 does not discriminate between these two strands (Fig. 3; the data for probe B not shown). It should be mentioned here, that the diagonals corresponding to histone H1 (Fig. 3 a,b) can be found in the regions of DNA fragments longer than about 250 bases; this must be caused by interactions of histone H1 with the DNA regions bordering the nucleosome-free promoter region of about 200 bp in length.

DISCUSSION

Activation of *hsp* genes causes a substantial increase in transcription (15), the density of RNA-polymerase molecules rises to the level of 2–3 RNA-polymerases per 200 bp of nucleosomal length (9). Such a high density of large multimeric complex of RNA-polymerase, whose dimensions are comparable with the size of the nucleosomal DNA, should lead to some changes in histone-DNA contacts. However, the histone H1-DNA contacts fixed by UV-crosslinking appeared to be unaffected. An assumption that histone H1 is preserved in the coding strand during RNA-polymerase movement along the gene has not been confirmed as the degree of histone H1 crosslinking was the same in both strands (data not shown).

Our data are consistent with those of Nacheva et al.(8), who

used various modifications of crosslinking proteins to partially depurinated DNA with dimethylsulfate. In active *hsp-70* chromatin, the interaction between DNA and the 'globular' domains of histones was shown to be disrupted, but histones remained associated with DNA via their destructured N- or C-'tails', which seem to function as 'bridges' compartmentalizing histones on DNA. As it was demonstrated by Stefanovsky et.al.(16, 17) UV-light also crosslinks histone H1 to DNA predominantly through its C- and N-terminal tails.

In contrast from the earlier studies which suggested that the binding of H1 repress the gene activity(18, 19), more recent works has tended toward the view that histone H1 is apparently present in transcriptionally active regions but it may differently interact with DNA in active and inactive chromatin(20, 21, 22).

However, Postnikov et al.(23) using several crosslinking methods and isolation of specific complexes by antibodies showed a twofold reduced density of H1 and H5 on the transcribing *b* globin gene compared with the inactive ovalbumin and lysozyme genes of chicken embryonic erythrocyte. The authors in great detail discussed the contradictory of their data with those of Nacheva et.al.(8), so we haven't touched this problem here.

We found two nonhistone proteins characteristic of the promoter region of activated *hsp-70* gene. Earlier Wu, using the Exo III protection technique showed the presence of two proteins in the regions upstream of activated *Drosophila hsp-70* and *hsp-82* heat shock genes (24–26). One of these genes binds to the TATA-box sequence independently of gene activation. The other can be found only after heat shock stimulation and is an *hsp* gene transcription activator (HSTF, heat shock transcription factor). This protein was purified by Wu et al. and its molecular mass was found to be 110 kD (27). This suggests that the crosslinked protein with an approximate molecular mass of 100 kD is an *hsp* gene's transcription activating factor (see also (28)).

Although the functional role of p50 remains unclear, it may be a part of a large transcription complex, in particular transcription factors (IIA, IIB or IIF) (29). A conclusive answer to the question whether or not p50 can be identified with TFs could be given after mapping the crosslinking sites on DNA for p50 and p100. It would be interesting to find out if the preferential crosslinking of these proteins to different strands is an essential feature of this DNA-protein complex or it just reflects some specificity in crosslinking.

We were unable to detect DNA-RNA-polymerase contacts within the regions analysed, because of a rather low efficiency of RNA-polymerase crosslinking; the same could be true in case of some other proteins.

Protein mapping and genome sequencing

We develop the 'protein image' analysis which can be used for mapping proteins on any genomic DNA as far as its probe or sequence are available. Many sequences can be tested by repeatedly hybridizing the same blot with a number of probes. The mapping proteins along genomic DNA can be directly coupled with the program of genome sequencing and it will not need much additional efforts.

ACKNOWLEDGEMENTS

We thank Miss E.I.Novikova for help in manuscript preparation. This study was supported by a grant from the Russian Human Genome Program and Russian State program 'Frontiers in Genetics'.

REFERENCES

1. Karpov, V.L., Preobrazhenskaya, O.V., Mirzabekov, A.D. (1984) *Cell* 36, 423–431
2. Wasyluk, B. (1988) *Biochimica et Biophysica Acta* 951, 17–35.
3. Pederson, D.S., Thoma, F. and Simpson, R.T. (1986) *Ann. Rev. Cell Biol* 34, 1033–1042
4. Zlatanova, J. (1990) *Trends Biochem.Sci.* 15, 273–276
5. Daneholt, B. and Grossbach, U. (1990) *Trends Biochem.Sci.* 15, 463–470
6. Echallier, G. and Ohanessian, A. (1970) *In Vitro* 6, 162–172
7. Gilmour, D. and Lis, J. (1985) *Mol. Cell. Biol.* 5, 2009–2018
8. Nacheva, G.A., Guschin, D.Y., Preobrazhenskaya, O.V., Karpov, V.L., Ebralidse, K.K. and Mirzabekov, A.D. (1989) *Cell* 58, 27–36
9. Mirzabekov, A.D., Bavykin, S.G., Belyavsky, A.V., Karpov, V.L., Preobrazhenskaya, O.V., Shick, V.V., Ebralidse, K.K. (1989) *Meth. Enzymology* 170, 386–408
10. Church, G. and Gilbert, W. (1984) *Proc. Natl. Acad. Sci. USA* 81, 1991–1995
11. Holmgren, R., Livak, R., Morimoto, R., Freund, R. and Meselson, M. (1979) *Cell* 18, 1359–1370
12. Costlow, N and Lis, J.T (1984) *Mol. Cell. Biol.* 4, 1853–1863
13. Kolchinsky, A.M., Vashakidse, R.P., Evgen'ev, M.B. and Mirzabekov, A.D. (1982) *Molecular Biology (USSR)* 6, 302–314
14. Jamrich, M. and Miller, O.L. (1984) *EMBO J.* 3, 1541–1545
15. Ashburner, M. and Bonner, J.J. (1979) *Cell* 17, 241–254
16. Stefanovsky, V.Yu., Dimitrov, S.I., Angelov, D. and Pashev, I.G. (1989) *Biochem. Biophys. Res. Commun.* 164, 304–310.
17. Stefanovsky, V.Yu., Dimitrov, S.I., Russanova, V.R., Angelov, D. and Pashev, I.G. (1989) *Nucl.Acids Res.* 17, 10069–10081
18. Weintraub, H. (1985) *Cell* 42, 705–711.
19. Schlissel, M.S. and Brown, D.D. (1984) *Cell* 35, 903–913.
20. Garrard, W.T. (1991) *BioEssays* 13, 87–88.
21. Baldwin, J.P. (1992) *Current Opinion in Structural Biology*, 2, 78
22. van Holde, K.E., Lohr, D.E., Robert, C., *J.B.Chem.*, 267, 2837–2840.
23. Postnikov, Y.V., Shick, V.V., Belyavsky, A.V., Khrapko, K.R., Brodolin, K.L., Nikolskay, T.A., Mirzabekov, A.D. (1991) *NAR* 19, 717–725.
24. Wu, C. (1984a) *Nature* 309, 229–234.
25. Wu, C. (1984b) *Nature* 311, 81–84.
26. Wu, C. (1985) *Nature* 317, 84–87.
27. Wu, C, Wilson, S., Walker, B., Dawid, I., Paisley, T. and Zimarino, V. (1987) *Science* 238, 1247–1253.
28. Wiederrecht, G., Shuey, D.J., Kibbe, W.A. and Parker, C.S. (1987) *Cell* 48, 507–515.
29. Greenblatt, J. (1992) *Nature* 360, 16–17