Rare disease

Spontaneous spinal epidural haematoma due to arteriovenous malformation in a child

António Jorge Cabral,¹ Andreia Barros,² Cristina Aveiro,¹ Rui Vasconcelos¹¹Pediatrics Department, Funchal Central Hospital, Funchal, Portugal;²Pediatrics Department, Hospital Nélio Mendonça, Funchal, PortugalCorrespondence to Dr Andreia Barros, asofiabarrosgmail.com

Summary

Spontaneous spinal epidural haematoma (SSEH) is a rare clinical entity, especially in infants, in whom only a few cases have been reported. In a paediatric emergency setting, SSEH should be considered as part of the differential diagnosis for acute extremity weakness and paraesthesia. Epidural vascular malformations are often suspected in these cases but have rarely been demonstrated. The authors report herein a case of SSEH in a 9-year-old boy arising from an epidural vascular malformation. He initially presented with sudden intense cervicodorsal pain followed by hypotonic lower extremities and progressive motor weakness, with no sensory change. The MRI showed an acute extradural haematoma extending from C7 to T4 with compression of the spinal cord. After submission to decompression surgery, he presented full recovery in 1 month. The histopathological analysis revealed a vascular malformation.

BACKGROUND

Spontaneous spinal epidural haematomas (SSEH) are a rare clinical entity with an incidence of 0.1 per 100 000 per year.^{1–4} There is preponderance in older-aged patients, especially those receiving anticoagulant drugs and have bleeding and coagulation defect tendencies. During childhood, they are even more unlikely^{5–7} with only few cases reported^{8–19} and their causes are frequently unknown.^{5–20} The youngest case reported is on a 22-month-old male infant.²¹ We report a rare case of spontaneous haemorrhage from an epidural vascular malformation in a 9-year-old boy. The haematoma and spinal cord compression was diagnosed by MRI, and the pathology was confirmed by microscopic analysis.

CASE PRESENTATION

This 9-year-old white male presented with a sudden intense cervicodorsal pain when he rose at dawn. After roughly 30 min, he demonstrated progressive muscular weakness of the lower extremities. There was no history of trauma to the head or spine, nor any bleeding disorder or pertinent family history. He did not have any fever, rhinorrhoea or other upper respiratory tract symptoms.

On arrival at our institution, the pain had subsided but he was non-ambulatory. The neurological examination showed normal upper extremity motor and tone, and normal cranial nerve examination. He demonstrated hypotonic lower extremities, and motor weakness in the lower limbs with no antigravity movement. Deep tendon reflexes were absent in the legs. Sensory evaluation was normal. A complete laboratory work-up, including bleeding time, prothrombin and thromboplastin time, plaque count, serum electrolytes, blood glucose level and calcium, was within normal limits.

Plain radiographs of the chest were normal. The CT of the spine did not show any evidence of destructive change or fracture of the vertebral body and neural arch. MRI of

the spine revealed an extradural mass, compatible with acute haematoma, extending from the C7 to T4 levels, compressing the spinal cord (figures 1 and 2).

The patient underwent surgery with C7 to T4 laminectomies and decompression of the extradural haematoma at which time anomalous blood vessels were detected and sent for histopathological examination. Microscopy analysis of the surgical specimen revealed an arteriovenous vascular malformation.

After surgery, the patient began a rehabilitation programme with an uneventful and progressive recovery and was released 26 days after admission with a normal neurological exam.

INVESTIGATIONS

MRI of the spine revealed an extradural mass, compatible with acute haematoma, extending from the C7 to T4 levels, compressing the spinal cord.

Microscopy analysis of the surgical specimen revealed an arteriovenous vascular malformation.

TREATMENT

The patient underwent surgery with C7 to T4 laminectomies and decompression of the extradural haematoma at which time anomalous blood vessels were detected and sent for histopathological examination.

OUTCOME AND FOLLOW-UP

After surgery, the patient began a rehabilitation programme with an uneventful recovery and a normal neurological examination within a month.

DISCUSSION

SSEH in children are rare lesions of unclear origin^{5–7} and in only 10% of the patients is the underlying cause

Figure 1 MRI sagittal T1-weighted images: left posterolateral extradural isointense mass spanning C4 to T7 vertebral segments (arrow) and appears inhomogeneous with compression of the spinal cord.

identified.²² A vascular malformation that has been overlooked or obliterated is one possible explanation in idiopathic cases.²³

In the case presented, a vascular malformation caused the acute clinical features with no significant medical history, no history of trauma and no warning signs. The presenting symptoms of SSEH are usually a consequence of spinal cord compression and roots compression, resulting in a sudden onset of back and neck pain. In adults, the pain and neurological deficits are characteristically specific and localised, but in children, particularly those under the age of 2 years, the initial symptoms are non-specific. Older children may report back pain, weakness or paraesthesia, whereas infants may have irritability or excessive crying. In some children, there are no significant symptoms to

suggest SSEH. Therefore, accurate and early diagnosis may be difficult.^{8 14 20}

The majority of spinal arteriovenous malformations are dural (up to 70%).²⁴ Epidural vascular malformations are rare, particularly in children, and are fed by radicular vessels accompanying the exiting nerve root, more commonly in the lower cervical and thoracic spine, where radicular arteries are more prominent.^{5 22 23}

MRI is the choice of imaging modality and plays an important role in diagnosis and evaluation of SSEH^{14 25} by simplifying the diagnostic work-up of epidural compression, as it can differentiate soft tissue from vascular and bony pathology.^{22 26-28}

Rapid surgical evacuation has been recommended as a treatment of choice for symptomatic SSEH^{8 12 29}. The

Figure 2 MRI axial T1-weighted: compression of the spinal cord that is displaced anteriorly and to the right by the mass (arrow).

outcome of patients treated surgically is generally good^{5 20 23 26}. The most significant factors that determine neurological outcome are interval from symptom onset to surgery, and the degree of neurological deficits prior to surgery. The shorter interval from symptom onset to surgery and minor neurological deficit has been reported to result in more favourable outcome.²⁹ The overall good outcome associated with epidural arteriovenous malformations is in distinct contrast to the poorer prognosis associated with dural vascular malformations^{26 30} and is likely related to the alteration of spinal cord blood flow rather than to the spinal cord compression.³¹

Learning points

- ▶ Spontaneous epidural haematomas should be considered in acute extremity weakness and paraesthesia in children
- ▶ Prompt MRI of the brain and spine is essential for correct diagnosis
- ▶ Immediate surgical decompression offers the best chance for improved outcome
- ▶ Prognosis of epidural vascular malformations is favourable with complete recovery.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Groen RJ**, van Alphen HA. Operative treatment of spontaneous spinal epidural hematomas: a study of the factors determining postoperative outcome. *Neurosurgery* 1996;**39**:494–508; discussion 508–9.
2. **Holtås S**, Heiling M, Lönntoft M. Spontaneous spinal epidural hematoma: findings at MR imaging and clinical correlation. *Radiology* 1996;**199**:409–13.
3. **Aksay E**, Kiyan S, Yuruktumen A, *et al*. A rare diagnosis in emergency department: spontaneous spinal epidural hematoma. *Am J Emerg Med* 2008;**26**:835.e3–5.
4. **Riaz S**, Jiang H, Fox R, *et al*. Spontaneous spinal epidural hematoma causing Brown-Sequard syndrome: case report and review of the literature. *J Emerg Med* 2007;**33**:241–4.
5. **Foo D**, Rossier AB. Preoperative neurological status in predicting surgical outcome of spinal epidural hematomas. *Surg Neurol* 1981;**15**:389–401.
6. **Posnikoff J**. Spontaneous spinal epidural hematoma of childhood. *J Pediatr* 1968;**73**:178–83.
7. **Robertson WC Jr**, Lee YE, Edmonson MB. Spontaneous spinal epidural hematoma in the young. *Neurology* 1979;**29**:120–2.
8. **Caldarelli M**, Di Rocco C, La Marca F. Spontaneous spinal epidural hematoma in toddlers: description of two cases and review of the literature. *Surg Neurol* 1994;**41**:325–9.
9. **Hehman K**, Norrell H. Massive chronic spinal epidural hematoma in a child. *Am J Dis Child* 1968;**116**:308–10.
10. **JACKSON FE**. Spontaneous spinal epidural hematoma coincident with whooping cough. case report. *J Neurosurg* 1963;**20**:715–7.
11. **Lee JS**, Yu CY, Huang KC, *et al*. Spontaneous spinal epidural hematoma in a 4-month-old infant. *Spinal Cord* 2007;**45**:586–90.
12. **Liao CC**, Lee ST, Hsu WC, *et al*. Experience in the surgical management of spontaneous spinal epidural hematoma. *J Neurosurg* 2004;**100**(1 Suppl Spine):38–45.
13. **Licata C**, Zoppetti MC, Perini SS, *et al*. Spontaneous spinal haematomas. *Acta Neurochir (Wien)* 1988;**95**:126–30.
14. **Pai SB**, Maiya PP. Spontaneous spinal epidural hematoma in a toddler—a case report. *Childs Nerv Syst* 2006;**22**:526–9.
15. **Patel H**, Boaz JC, Phillips JP, *et al*. Spontaneous spinal epidural hematoma in children. *Pediatr Neurol* 1998;**19**:302–7.
16. **SHENKIN HA**, HORN RC Jr, GRANT FC. Lesions of the spinal epidural space producing cord compression. *Arch Surg* 1945;**51**:125–46.
17. **Vallée B**, Besson G, Gaudin J, *et al*. Spontaneous spinal epidural hematoma in a 22-month-old girl. *J Neurosurg* 1982;**56**:135–8.
18. **Lim JJ**, Yoon SH, Cho KH, *et al*. Spontaneous spinal epidural hematoma in an infant: a case report and review of the literature. *J Korean Neurosurg Soc* 2008;**44**:84–7.
19. **Lo MD**. Spinal cord injury from spontaneous epidural hematoma: report of 2 cases. *Pediatr Emerg Care* 2010;**26**:445–7.

20. **Beatty RM**, Winston KR. Spontaneous cervical epidural hematoma. A consideration of etiology. *J Neurosurg* 1984;**61**:143–8.
21. **Muhonen MG**, Piper JG, Moore SA, *et al.* Cervical epidural hematoma secondary to an extradural vascular malformation in an infant: case report. *Neurosurgery* 1995;**36**:585–7; discussion 587–8.
22. **D'Angelo V**, Bizzozero L, Talamonti G, *et al.* Value of magnetic resonance imaging in spontaneous extradural spinal hematoma due to vascular malformation: case report. *Surg Neurol* 1990;**34**:343–4.
23. **Emery DJ**, Cochrane DD. Spontaneous remission of paralysis due to spinal extradural hematoma: case report. *Neurosurgery* 1988;**23**:762–4.
24. **Lev N**, Maimon S, Rappaport ZH, *et al.* Spinal dural arteriovenous fistulae—a diagnostic challenge. *Isr Med Assoc J* 2001;**3**:492–6.
25. **Crisi G**, Sorgato P, Colombo A, *et al.* Gadolinium-DTPA-enhanced MR imaging in the diagnosis of spinal epidural haematoma. Report of a case. *Neuroradiology* 1990;**32**:64–6.
26. **Avrahami E**, Tadmor R, Ram Z, *et al.* MR demonstration of spontaneous acute epidural hematoma of the thoracic spine. *Neuroradiology* 1989;**31**:89–92.
27. **Dickman CA**, Zabramski JM, Sonntag VK, *et al.* Myelopathy due to epidural varicose veins of the cervicothoracic junction. Case report. *J Neurosurg* 1988;**69**:940–1.
28. **Nagel MA**, Taff IP, Cantos EL, *et al.* Spontaneous spinal epidural hematoma in a 7-year-old girl. Diagnostic value of magnetic resonance imaging. *Clin Neurol Neurosurg* 1989;**91**:157–60.
29. **Lawton MT**, Porter RW, Heiserman JE, *et al.* Surgical management of spinal epidural hematoma: relationship between surgical timing and neurological outcome. *J Neurosurg* 1995;**83**:1–7.
30. **Coats TJ**, King TT. The diagnosis of dural spinal vascular malformations. *Br J Neurosurg* 1991;**5**:609–15.
31. **Maurice-Williams RS**. Spinal dural arteriovenous malformations—a treatable cause of progressive paraparesis in elderly people. *Age Ageing* 1992;**21**:412–6.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Cabral AJ, Barros A, Aveiro C, Vasconcelos R. Spontaneous spinal epidural haematoma due to arteriovenous malformation in a child. *BMJ Case Reports* 2011; 10.1136/bcr.02.2011.3875, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow