Published in final edited form as:

J Am Acad Child Adolesc Psychiatry. 2010 December ; 49(12): 1189–1285. doi:10.1016/j.jaac.2010.08.017.

Neurobiology of the Adolescent Brain and Behavior

BJ Casey, PhD and Rebecca M. Jones, MS

Sackler Institute for Developmental Psychobiology, Weill Cornell Medical College

Introduction

Adolescence is a transitional period of development when there are many changes experienced concomitantly, including physical maturation, drive for independence, increased salience of social and peer interaction, and brain development^{1–3}. This developmental period is also a time characterized by an inflection in risky behaviors including experimentation with drugs and alcohol, criminal activity and unprotected sex. Understanding the neural basis of these risky behaviors is key in identifying which teens may be at risk for poor outcomes, such as substance dependence and abuse.

A number of hypotheses have been postulated for why adolescents may engage in impulsive and risky behaviors. Traditional accounts of adolescence suggest that it is a period of development associated with progressively greater efficiency of cognitive control capacities. This efficiency in cognitive control is described as dependent on maturation of the prefrontal cortex as evidenced by imaging^{4–7} and post mortem studies^{8–10} showing continued structural and functional development of this region well into young adulthood.

Improved cognitive control with development of the prefrontal cortex is consistent with a linear increase in this ability from childhood to adulthood. Yet suboptimal choices and actions observed during adolescence represent an inflection in development¹¹ that is unique from either childhood or adulthood, as evidenced by the National Center for Health Statistics on adolescent behavior and mortality¹². If cognitive control and an immature prefrontal cortex were the basis for suboptimal choice behavior alone, then children should look remarkably similar or presumably worse than adolescents, given their less developed prefrontal cortex and cognitive abilities². This review addresses the primary question of how the brain is changing during adolescence that may explain inflections in risky and impulsive behavior. In addition, we provide examples of how alcohol and drug use during this period of development may further exacerbate these changes and can lead to subsequent abuse and dependence.

To accurately capture cognitive and neurobiological changes during adolescence, this period must be treated as a transitional one rather than a single snapshot in time³. In other words, to understand this developmental period, transitions into and out of adolescence are necessary for distinguishing distinct attributes of this period relative to other time points in development. Therefore, empirical data that establishes developmental trajectories from childhood to adulthood for cognitive and neural processes are essential in characterizing these transitions and more importantly in constraining any interpretations about changes in brain or behavior in adolescence.

Second, accurate depictions of adolescence require a refinement in the phenotypic characterization of this period. For example, on a behavioral level, adolescents are often characterized as impulsive and greater risk-takers with these constructs used almost synonymously. Yet, these constructs are distinct and appreciating this distinction is important for describing their developmental trajectories and neural underpinnings. We provide behavioral, clinical and neurobiological evidence that suggest that risk-taking is more tightly coupled with sensitivity to environmental incentives (sensation-seeking) whereas impulsivity is associated with poor top down cognitive control.

To theoretically ground the empirical findings, we provide a plausible neurobiological model for adolescence and suggest how development during this time may lead to an enhancement in vulnerabilities for alcohol and drug abuse. The intention of this review is not to psychopathologize adolescence, but rather to explain why some teens but not others are vulnerable to substance abuse. As such, we attempt to identify potential biological and behavioral markers for early identification and for outcome assessments of interventions.

Neurobiological Model of Adolescence

A neurobiological model of adolescent development² that builds on rodent models^{13, 14} and recent imaging studies of adolescence^{6, 7, 15–20} is depicted Figure 1. This model illustrates how subcortical and prefrontal top-down control regions must be considered together as a circuit. The cartoon shows different developmental trajectories for signaling of these regions, with limbic projections developing earlier than prefrontal control regions. According to the model, the adolescent is biased by functionally mature subcortical relative to less mature cortical circuitry during adolescence (i.e., imbalance in reliance of systems), compared to children, for whom this frontolimbic circuitry is still developing; and compared to adults, for whom these systems are fully mature. With development and experience, the functional connectivity between these regions is strengthened and provides a mechanism for top down modulation of the subcortical systems⁷. Thus it is the frontostriatal circuitry, along with functional strengthening of connections within this circuitry, that may provide a mechanism to explain changes in both impulsivity and risk-taking observed across development.

This model is consistent with previous ones^{21–24} in that it provides a basis for nonlinear inflections observed in behavior from childhood to adulthood, due to earlier maturation of subcortical projections relative to less mature top down prefrontal ones. Specifically, the triadic model²¹ proposes that motivated behavior has three distinct neural circuits (approach, avoidance and regulatory). The approach system is largely controlled by the ventral striatum, avoidance system by the amygdala and lastly, the regulatory system by the prefrontal cortex²⁵. The current model differs largely from others in that it is grounded in empirical evidence for brain changes not only in the transition from adolescence to adulthood, but rather the transition *into* adolescence from childhood and later *out of* adolescence into adulthood. Moreover, the model does not suggest that the striatum and amygdala are specific to approach and avoidant behavior given recent studies showing valence independence of these structures²⁶, but rather are systems that are important in detecting motivationally and emotionally relevant cues in the environment that can bias behavior. In this review, we describe the most recent evidence from behavioral and human imaging studies of adolescence in the context of our model that illustrates the transition from childhood to adulthood.

Phenotypic Characterization of Adolescence

The ability to resist temptation in favor of long-term goals is a form of cognitive control. Lapses in this ability have been suggested to be at the very core of adolescent risky

behavior²⁷. Cognitive control, which includes resistance from temptation or delay of immediate gratification has been studied in the context of social, developmental and cognitive psychology. Developmentally, this ability has been measured by assessing how long a toddler can resist an immediate reward (e.g., a cookie) in favor of a larger reward later (e.g., two cookies)²⁸. Although individuals vary in this ability even as adults, developmental studies suggest windows of development when an individual may be particularly susceptible to temptations. This ability has been described as a form of impulse control²⁹ and it is multi-faceted^{30, 31}, but can be operationally defined as the ability to accomplish goal-directed behavior in the face of salient, competing inputs and actions³².

Historically, developmental studies have shown a steady improvement in cognitive control capacity from infancy to adulthood³³. This observation is supported by a wealth of behavioral evidence from experimental paradigms in controlled laboratory settings including paradigms such as the Go-NoGo task, Simon task, and task-switching paradigms that require participants to override a prepotent response in order to achieve a correct one^{32, 34}. However, when it is advantageous to suppress a response to incentive-related cues, cognitive control suffers²⁰. This reduced control is especially evident during the period of adolescence, when suboptimal choices in sexual and drug related behaviors peak^{3, 11, 12, 14}. These observations imply that developmental trajectories in cognitive control are complex and can be modulated by emotionally charged or reinforcing contexts (e.g., social and sexual interactions), in which cognitive control demands interact with motivational drives or processes.

Motivation can modulate cognitive control in at least two ways. First, being rewarded for performance on a given task can make people work harder and ultimately perform better than when not rewarded¹⁷. Second, the capacity to exert control can be challenged when required to suppress thoughts and actions toward appetitive cues²⁰. Recent studies of adolescent development have begun to compare cognitive control capacity in relatively neutral versus motivational contexts. These studies suggest a change in sensitivity to environmental cues, especially reward-based ones at different points in development, and suggest a unique influence of motivation on cognition during the adolescent years.

In the following section, we highlight some of the most recent studies of how adolescent behavior is differentially biased in emotionally charged contexts relative to adults.

For example, Ernst and colleagues^{35, 36} examined performance on an antisaccade task with a promise of financial reward for accurate performance on some trials but not others. Results showed that promise of a reward, facilitated adolescent cognitive control behavior more than for adults, a finding that has been replicated¹⁷ and recently been extended to social rewards (e.g. happy faces²⁰).

While the previous examples provide instances of enhanced performance in teens with incentives, rewards can also diminish performance when suppressing responses to rewards that lead to high gain. For example, using a gambling task in which reward feedback was provided immediately during decision-making (“hot” trials which heightened task-elicited affective arousal) or withheld until after the decision (“cold” deliberate decision making trials), Figner and colleagues³⁷ showed that adolescents made disproportionately more risky gambles compared to adults but only in the “hot” condition. Using a similar task, the Iowa Gambling Task, Cauffman and colleagues³⁸ have shown that this sensitivity to rewards and incentives actually peaks during adolescence, with a steady increase from late childhood to adolescence in tendency to play with more advantageous decks of cards and then a subsequent decline from late adolescence to adulthood. These findings illustrate a curvilinear function, peaking roughly between 13 and 17, and then declining²⁷. While prior

findings with the Iowa Gambling task have shown a linear increase in performance with age³⁹, these studies did not look at age continuously nor did they examine only trials with advantageous decks of cards.

Recent studies have suggested that social contexts, particularly peers, may also serve as a motivational cue and can diminish cognitive control during adolescence. It has been shown that the degree to which an adolescent's peers are using substances is directly proportional to the amount of alcohol or illegal substances that the adolescent themselves will use⁴⁰. Using a simulated driving task, Gardner and colleagues⁴¹ have shown that adolescents make riskier decisions in the presence of peers than when alone and that these risky decisions decrease linearly with age^{23, 40}.

Taken together, these studies suggest that during adolescence, motivational cues of potential reward are particularly salient and can lead to improved performance when provided as a reinforcer or rewarded outcome, but to riskier choices or suboptimal choices when provided as a cue. In the latter case, the motivational cue can diminish effective goal-oriented behavior. Furthermore, these studies suggest that sensitivity to rewards and sensation-seeking behavior are distinct from impulsivity with very different developmental patterns (curvilinear function versus a linear function, respectively). This distinction is further evident in a recent study by Steinberg et al.⁴² using self-report measures of sensation-seeking and impulsivity. They tested whether the often-conflated constructs of sensation-seeking and impulsivity develop along different timetables in nearly 1000 individuals between the ages of 10 and 30. The results showed that differences in sensation-seeking with age followed a curvilinear pattern, with peaks in sensation-seeking increasing between 10 and 15 years and declining or remaining stable thereafter. In contrast, age differences in impulsivity followed a linear pattern, with decreasing impulsivity with age in a linear fashion (see Figure 2 panel A). These findings together with the laboratory based findings, suggest heightened vulnerability to risk-taking in adolescence "may be due to the combination of relatively higher inclinations to seek excitement and relatively immature capacities for self-control that are typical of this period of development"⁴².

Neurobiology of Adolescence

As denoted in our model of adolescence, two key regions implicated in cognitive and motivational behavior are the prefrontal cortex, known to be important for cognitive control⁴³ and the striatum critical in detecting and learning about novel and rewarding cues in the environment⁴⁴. We highlight recent animal and human imaging work on neurobiological changes supporting these motivational and cognitive systems across development in the context of the previous behavioral findings on the development of sensation-seeking and impulsivity. We use the previously described imbalance model of linear development of top down prefrontal regions relative to a curvilinear function for development of bottom-up striatal regions involved in detecting salient cues in the environment to ground the findings. The importance of examining circuitry rather than specific regional change, especially within frontostriatal circuits that underlie different forms of goal-oriented behavior is key. This perspective moves the field away from examination of how each region matures in isolation to how they may interact in the context of interconnected circuits.

Seminal animal and human work has shown how striatal and prefrontal cortical regions shape goal-directed behavior^{7, 27, 37, 38, 44}. Using single-unit recordings in monkeys, Pasupathy & Miller⁴⁵ demonstrated that when flexibly learning a set of reward contingencies, very early activity in the striatum provides the foundation for reward-based associations, whereas later, more deliberative prefrontal mechanisms are engaged to

maintain the behavioral outputs that can optimize the greatest gains, these findings have been replicated in lesion studies^{46–48}. A role for the striatum in early temporal coding of reward contingencies prior to the onset of activation in prefrontal regions has also been extended to humans⁴⁹. These findings suggest that understanding the interactions between regions (along with their component functions); within frontostriatal circuitry is critical for developing a model of cognitive and motivational control in adolescence.

Frontostriatal circuits undergo considerable elaboration during adolescence^{50–53} that are particularly dramatic in the dopamine system. Peaks in the density of dopamine receptors, D1 and D2 in the striatum occur early in adolescence, followed by a loss of these receptors by young adulthood^{54–56}. In contrast, the prefrontal cortex does not show peaks in D1 and D2 receptor density until late adolescence and young adulthood^{57, 58}. Similar developmental changes have been shown in other reward related systems including cannabinoid receptors⁵⁹. It remains unclear how changes in the dopamine systems may relate to motivated behavior as controversy remains as to whether reward sensitivity is modulated by dopamine systems (e.g.,^{60, 61}) and whether it is a result of less active or hypersensitive dopamine systems (e.g.,^{62, 63}). However, given the dramatic changes in dopamine rich circuitry during adolescence, it is likely to be related to changes in sensitivity to rewards distinct from childhood or adulthood^{50, 64}. Beyond the significant changes in dopamine receptors, there are also dramatic hormonal changes which occur during adolescence that lead to sexual maturity, and influence functional activity in frontostriatal circuits⁶⁵, however, a detailed discussion is beyond the scope of this paper, see^{66, 67} for detailed reviews on the subject.

Human imaging studies have begun to provide support for strengthening in the connections of dopamine rich frontostriatal circuitry, across development. Using diffusion tensor imaging and functional magnetic resonance (fMRI), Casey and colleagues^{68, 69} and others⁷⁰ have shown greater strength in distal connections within these circuits across development and have linked connection strength between prefrontal and striatal regions with the capacity to effectively engage cognitive control in typically and atypically developing individuals^{68, 69}. These studies illustrate the importance of signaling within corticostriatal circuitry which support the capacity to effectively engage in cognitive control.

Likewise, there is mounting evidence from human functional neuroimaging studies on how subcortical systems like the striatum and the prefrontal cortex interact to give rise to risky behavior observed in adolescents⁷¹. The majority of imaging studies have focused on one or the other regions showing that the prefrontal cortex, thought to subserve age-related improvement in cognitive control^{72–78} undergoes delayed maturation^{4, 79, 80} while striatal regions sensitive to novelty and reward manipulations develop earlier^{74, 81}. Several groups have shown that adolescents show heightened activation of the ventral striatum in anticipation and/or receipt of rewards compared to adults^{6, 15, 17, 18}, but others report a hypo-responsiveness⁸².

One of the first studies to examine reward related processes across the full spectrum of development from childhood to adulthood was completed by Galvan and colleagues⁶ in 6 to 29 year olds. They showed that ventral striatal activation was sensitive to varying magnitudes of monetary reward⁴⁹ and that this response was exaggerated during adolescence, relative to children and adolescents⁶ (see Figure 3), indicative of signal increases⁶ or more sustained activation⁸³. In contrast to the pattern in the ventral striatum, orbital prefrontal regions, showed protracted development across these ages (Figure 2b).

But, how does this enhancement of signaling in the ventral striatum relate to behavior? In a follow-up study, Galvan and colleagues¹⁶ examined the association between activity in the

ventral striatum to large monetary reward with personality trait measures of risk-taking and impulsivity. Anonymous self-report rating scales of risky behavior, risk perception and impulsivity were acquired in her sample of 7 to 29 year olds. Galvan et al. showed a positive association between ventral striatal activity to large reward and the likelihood of engaging in risky behavior (see Figure 3) These findings are consistent with adult imaging studies showing ventral striatal activity with risky choices ^{84, 85}.

To further support an association between adolescents' risky behavior and sensitivity to reward as indexed by an exaggerated ventral striatal response, Van Leijenhorst and colleagues ¹⁸ tested this association using a gambling task. The task included Low-Risk gambles with a high probability of obtaining a small monetary reward and High-Risk gambles with a smaller probability of obtaining a larger monetary reward. The fMRI results confirmed that High-Risk choices were associated with ventral striatal recruitment whereas Low-Risk choices were associated with activation in ventral medial prefrontal cortex. These findings are consistent with the hypothesis that risky behavior in adolescence is associated with an imbalance caused by different developmental trajectories of subcortical reward and prefrontal regulatory brain regions consistent with our neurobiological model of adolescence.

While there appears to be an association between risk-taking behavior and ventral striatal activation, in the Galvan study ¹⁶ no correlation was reported between ventral striatal activity with impulsivity. Rather, impulsivity ratings were correlated with age, consistent with numerous imaging studies showing linear development with age in prefrontal cortical recruitment during impulse control tasks ^{7, 75, 77} (and see reviews by ^{34, 86}). Moreover, recent studies have shown that impulsivity ratings inversely correlate with volume of the ventral medial prefrontal cortex in a sample of healthy boys (7–17yrs) ⁸⁷. Finally, studies of clinical populations characterized by impulsivity problems like ADHD, show impaired impulse control and reduced activity in prefrontal regions compared to controls, ^{88, 89} but do not show heightened responses to incentives ⁹⁰.

These findings provide neurobiological empirical support for a dissociation of the constructs related to risk-taking and reward sensitivity from that of impulsivity with the former showing an curvilinear pattern and the latter a linear pattern (see Figure 2 B). Thus adolescent choices and behavior cannot be explained by impulsivity or protracted development of the prefrontal cortex alone. Rather, motivational subcortical regions must be considered to elucidate why adolescent behavior is not only different from adults, but from children as well. Thus, the ventral striatum appears to play a role in levels of excitement ^{82, 91} and positive affect ¹⁵ when receiving rewards, as well as the propensity for sensation-seeking and risk-taking ^{16, 91}. More importantly, these findings suggest that during adolescence, some individuals may be more prone to engage in risky behaviors due to developmental changes in concert with variability in a given individual's predisposition to engage in risky behavior, rather than to simple changes in impulsivity.

A scientific area that has received less attention is determining how cognitive control and motivational systems interact over the course of development. As mentioned earlier, Ernst and colleagues ^{35, 36} showed that promise of a monetary reward facilitated adolescent cognitive control behavior more than for adults. Geier et al. ¹⁷ recently identified the neural substrates of this cognitive up-regulation using a variant of an antisaccade task during functional brain imaging. In adolescents and adults, trials for which money was at stake speeded performance and facilitated accuracy, but this effect was larger in adolescents. Following a cue that the next trial would be rewarded, adolescents showed exaggerated activation in the ventral striatum while preparing for and subsequently executing the antisaccade. An exaggerated response was observed in adolescents within prefrontal regions

along the precentral sulcus, important for controlling eye movements, suggesting a reward-related up-regulation in control regions as well.

Rewards, as suggested above, can enhance as well as diminish goal-directed behavior. The observation that adolescents take more risks when appetitive cues are present versus absent during gambling tasks makes this point (e.g., ³⁷). In a recent imaging study ²⁰, Somerville et al. identified the neural substrates of down-regulation of control regions with appetitive cues. Somerville et al. tested child, adolescent, and adult participants while they performed a go nogo task with appetitive social cues (happy faces) and neutral cues. Task performance to neutral cues showed steady improvement with age on this impulse control task. However, on trials for which the individual had to resist approaching appetitive cues, adolescents failed to show the expected age-dependent improvement. This performance decrement during adolescence was paralleled by enhanced activity in the striatum. Conversely, activation in the inferior frontal gyrus was associated with overall accuracy and showed a linear pattern of change with age for the nogo versus go trials. Taken together, these findings implicate exaggerated ventral striatal representation of appetitive cues in adolescents in the absence of a mature cognitive control response.

Collectively, these data suggest that although adolescents as a group are considered risk-takers ⁴¹, some adolescents will be more prone than others to engage in risky behaviors, putting them at potentially greater risk for negative outcomes. These findings underscore the importance of considering individual variability when examining complex brain-behavior relationships related to risk-taking and impulsivity in developmental populations. Further, these individual and developmental differences may help to explain vulnerability in some individuals to risk-taking which is associated with substance use, and ultimately, addiction ⁶⁴.

Substance Use and Abuse in Adolescents

Adolescence marks a period of increased experimentation with drugs and alcohol ⁹², with alcohol being the most abused of illegal substances by teens ^{11, 93, 94}. Early use of these substances, such as alcohol, is a reliable predictor of later dependence and abuse ⁹⁵. Given the surge in alcohol dependence between adolescence and adulthood that is unequaled at any other developmental stage ⁹⁶, we focus predominantly on a select review here of its use and abuse in adolescents and motivational properties.

Alcohol as well as other substances of abuse, including cocaine and cannabinoids, have been shown to have reinforcing properties. These substances influence mesolimbic dopamine transmission with acute activations of neurons in frontolimbic circuitry rich in dopamine, including the ventral striatum ^{97–99}. As suggested by Hardin and Ernst (2009) ⁹², the use of these substances may exacerbate an already enhanced ventral striatum response resulting in heightened or strengthening of reinforcement properties to the drug. Robinson and Berridge ^{61, 63, 100} have suggested that these drugs of abuse can “hijack” the systems associated with drug incentives like the ventral striatum, thus down regulating top down prefrontal control regions.

The majority of empirical work on adolescent use of alcohol has been done in animals, given ethical constraints in performing such studies in human adolescents. Animal models of ethanol also provide the most evidence for differential effects of alcohol in adolescents relative to adults and are consistent with human findings of adolescents having relative insensitivity to ethanol effects. Spear and colleagues have shown that adolescent rats relative to adults, are less sensitive to the social, motor, sedation, acute withdrawal and “hangover effects” of ethanol ^{101–103}. These findings are significant in that many of these effects serve as cues to limit intake in adults ¹¹. Likewise, at the same time when adolescents are

insensitive to cues that may help to limit their alcohol intake, positive influences of alcohol such as social facilitation may further encourage alcohol use¹⁰⁴. Most risky behaviors in humans- including alcohol abuse- occur in social situations²³, potentially pushing adolescents towards greater use of alcohol and drugs when this behavior is valued by their peers.

How is the brain altered with alcohol use and abuse in adolescence relative to adults? Whereas adolescents may be less sensitive to some of the behavioral effects of alcohol, they appear to be more sensitive to some of the neurotoxic effects⁹⁴. For example, physiological studies (e.g.,¹⁰⁵) show greater ethanol induced inhibition of NMDA-mediated synaptic potentials and long-term potentiation in hippocampal slices in adolescents than in adults. Repeated exposure of intoxicating doses of ethanol also produces greater hippocampal dependent memory deficits^{106, 107} and prolonged ethanol exposure has been associated with increased dendritic spine size¹⁰⁸. These latter findings of dendritic spine changes are suggestive of modification of brain circuitry that may stabilize addictive behavior⁹⁴.

Data from brain imaging studies provide parallel evidence in humans of neurotoxic effects of alcohol on the brain. A number of studies have reported altered brain structure and function in alcohol-dependent or -abusing adolescents and young adults compared to healthy individuals. These studies show smaller frontal and hippocampal volumes, altered white matter microstructure and poorer memory^{109–113}. Moreover, these studies show positive associations between hippocampal volumes and age of first use¹⁰⁹ suggesting that early adolescence may be a period of heightened risk to alcohol's neurotoxic effects. Duration, which was negatively correlated with hippocampal volume, may compound this effect.

Currently, only a few studies have examined functional brain activity to drug or alcohol related stimuli (i.e. pictures of alcohol) in adolescents¹¹⁴, although this is an area of future research (see¹¹⁵). Studies of high-risk populations (e.g. familial load of alcohol dependence) suggest impairments in frontal functioning are apparent prior to drug use exposure (e.g.^{116, 117}) and can predict later substance use^{118, 119}. However, in an early behavioral study of the effects of alcohol in 8 to 15 year old boys of low and high familial risk¹²⁰, the most significant finding was little if any behavioral change or problem on tests of intoxication -even after given doses which had been intoxicating in an adult population were observed. These neurotoxic effects together with increased sensitivity to the motivational effects of alcohol and evidence of poorer top down prefrontal control apparent even prior to drug use exposure¹¹⁶ may set up a long-term course of alcohol and drug abuse well beyond adolescence^{118, 119}.

Conclusions

Together, the studies described support a view of adolescent brain development as characterized by a tension between early emerging “bottom-up” systems that express exaggerated reactivity to motivational stimuli and later maturing “top-down” cognitive control regions. This bottom-up system which is associated with sensation- seeking and risk-taking behavior gradually loses its competitive edge with the progressive emergence of “top-down” regulation (e.g.,^{2, 7, 15, 23, 64, 121–123}). This imbalance between these developing systems during adolescence may lead to heightened vulnerability to risk-taking behaviors and increased susceptibility to the motivational properties of substances of abuse.

This review provides behavioral, clinical and neurobiological evidence for dissociating these subcortico-cortico systems developmentally. Behavior data from laboratory tasks and self-report ratings administered to children, adolescents and adults (e.g.,^{18, 20, 37, 42}) suggest curvilinear development of sensation-seeking with a peak inflection roughly between 13 and

17 years, while impulsivity decreases across development in a linear fashion from childhood to young adulthood. Human imaging studies show patterns of activity in subcortical brain regions sensitive to reward (ventral striatum) that parallel the behavioral data. Specifically, they show a curvilinear pattern of development in these regions and the magnitude of their response is associated with risk-taking behaviors. In contrast, prefrontal regions, important in top down regulation of behavior, show a linear pattern of development that parallels those seen in behavioral studies of impulsivity. Moreover, clinical disorders with impulse control problems show less prefrontal activity, further linking neurobiological substrates with the phenotypic construct of impulsivity.

The tension between subcortical regions relative to prefrontal cortical regions during this period may serve as a possible mechanism for the observed heightened risk-taking, including use and abuse of alcohol and drugs. The majority of adolescents have tried alcohol⁹³, but this does not necessarily lead to abuse. Individuals with less top-down regulation may be particularly susceptible to alcohol and substance abuse as suggested by studies of high-risk populations showing impairments in frontal functioning prior to alcohol and drug exposure (e.g.^{116, 117}). In the context of our neurobiological model of adolescence, these individuals would have an even greater imbalance in cortico-subcortical control. These findings are also in accordance with clinical findings in ADHD populations who show reduced prefrontal activity and are four times as likely to develop a substance use disorder compared to healthy controls¹²⁴. This imbalance in cortico-subcortical control would be further compounded by the insensitivity of adolescence to the motor and sedative effects of alcohol that otherwise may help to limit intake, and the positive influences of alcohol in social facilitation which may further encourage alcohol use¹⁰⁴. As shown by Steinberg and colleagues^{23, 41}, most risky behaviors- including alcohol and substance abuse- occur in social situations. Thus use of alcohol and drugs may be encouraged and maintained by peers when this behavior is valued.

One of the challenges in addiction related work is the development of biobehavioral markers for early identification of risk for substance abuse and/or for outcomes assessments for interventions/treatments. Our findings suggest that behavioral challenges that require both cognitive control in the presence of tempting appetitive cues may be useful potential markers. Example of such behavioral assays include gambling tasks with high and low risk or “hot” and “cold” conditions described in this review^{18, 37} or simple impulse control tasks that require suppressing a response to an appetitive/tempting cue²⁰. These tasks are reminiscent of the delay of gratification task developed by Mischel¹²⁵. In fact, performance on simple impulse control tasks such as these in adolescents and adults has been associated with their performance as toddlers on the delay of gratification task^{28, 29}. Mischel and colleagues have shown the high level of stability and predictive value of this task in later life. Relevant to substance abuse, they showed that the ability to delay gratification as a toddler, predicted less substance abuse (e.g., cocaine) later in life¹²⁶. In our current work, we are beginning to use a combination of these tasks to identify the neural substrates of this ability to further understand potential risk factors for substance abuse.

Collectively, these data suggest that although adolescents as a group are considered risk-takers⁴¹, some adolescents will be more prone than others to engage in risky behaviors, putting them at potentially greater risk for negative outcomes. However, risk-taking can be quite adaptive in the right environments. So rather than trying to eliminate adolescent risk-taking behavior that has not been a successful enterprise to date²³, a more constructive strategy may be to provide access to risky and exciting activities (e.g., after school programs with in-door wall climbing) under controlled settings and limit harmful risk-taking opportunities. As the adolescent brain is a reflection of experiences, with these safe risk-taking opportunities, the teenager can shape long-term behavior by fine tuning the

connections between top-down control regions and bottom-up drives with maturity of this circuitry. Other successful strategies are cognitive behavioral therapies which focus on refusal skills, or cognitive control, to reduce risky behaviors¹²⁷. The findings underscore the importance of considering individual variability when examining complex brain-behavior relationships related to risk-taking and impulsivity in developmental populations. Further, these individual and developmental differences may help explain vulnerability in some individuals to risk-taking associated with substance use, and ultimately, addiction.

Acknowledgments

This work was supported in part by NIDA R01 DA018879, NIDA Pre-Doctoral training grant DA007274, the Mortimer D. Sackler family, the Dewitt-Wallace fund, and by the Weill Cornell Medical College Citigroup Biomedical Imaging Center and Imaging Core.

References

1. Blakemore S-J. The social brain in adolescence. *Nature Reviews Neuroscience*. 2008; 9:267–277.
2. Casey BJ, Getz S, Galvan A. The adolescent brain. *Dev Rev*. 2008; 28(1):62–77. [PubMed: 18688292]
3. Casey BJ, Jones RM, Hare TA. The adolescent brain. *Ann N Y Acad Sci*. 2008 Mar; 1124:111–126. [PubMed: 18400927]
4. Sowell ER, Peterson BS, Thompson PM, Welcome SE, Henkenius AL, Toga AW. Mapping cortical change across the human life span. *Nat Neurosci*. 2003 Mar; 6(3):309–315. [PubMed: 12548289]
5. Gogtay N, Giedd JN, Lusk L, et al. Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences, USA*. 2004; 101(21):8174–8179.
6. Galvan A, Hare TA, Parra CE, et al. Earlier development of the accumbens relative to orbitofrontal cortex might underlie risk-taking behavior in adolescents. *Journal of Neuroscience*. 2006 Jun 21; 26(25):6885–6892. [PubMed: 16793895]
7. Hare TA, Tottenham N, Galvan A, Voss HU, Glover GH, Casey BJ. Biological substrates of emotional reactivity and regulation in adolescence during an emotional go-nogo task. *Biol Psychiatry*. 2008 May 15; 63(10):927–934. [PubMed: 18452757]
8. Bourgeois JP, Goldman-Rakic PS, Rakic P. Synaptogenesis in the prefrontal cortex of rhesus monkeys. *Cerebral Cortex*. 1994; 4:78–96. [PubMed: 8180493]
9. Huttenlocher PR. Synaptic density in human frontal cortex - developmental changes and effects of aging. *Brain Research*. 1979; 163:195–205. [PubMed: 427544]
10. Rakic P. Synaptic development of the cerebral cortex: implications for learning, memory and mental illness. *Prog. Brain Res*. 1994; 102:227–243. [PubMed: 7800815]
11. Windle M, Spear LP, Fuligni AJ, et al. Transitions into underage and problem drinking: developmental processes and mechanisms between 10 and 15 years of age. *Pediatrics*. 2008; 121:S273–S289. [PubMed: 18381494]
12. Eaton LK, Kann L, Kinchen S, et al. Youth Risk Behavior Surveillance - United States, 2007, surveillance summaries. *Morbidity and Mortality Weekly Report*. 2008; 57(SS04):1–131. [PubMed: 18185492]
13. Laviola G, Adriani W, Terranova ML, Gerra G. Psychobiological risk factors for vulnerability to psychostimulants in human adolescents and animal models. *Neurosci Biobehav Rev*. 1999 Nov; 23(7):993–1010. [PubMed: 10580313]
14. Spear LP. The adolescent brain and age-related behavioral manifestations. *Neuroscience and Biobehavioral Reviews*. 2000; 24(4):417–463. [PubMed: 10817843]
15. Ernst M, Nelson EE, Jazbec S, et al. Amygdala and nucleus accumbens in responses to receipt and omission of gains in adults and adolescents. *Neuroimage*. 2005 May 1; 25(4):1279–1291. [PubMed: 15850746]
16. Galvan A, Hare T, Voss H, Glover G, Casey BJ. Risk-taking and the adolescent brain: who is at risk? *Dev Sci*. 2007 Mar; 10(2):F8–F14. [PubMed: 17286837]

17. Geier CF, Terwilliger R, Teslovich T, Velanova K, Luna B. Immaturities in Reward Processing and Its Influence on Inhibitory Control in Adolescence. *Cereb Cortex*. 2009 Oct 29.
18. Van Leijenhorst L, Moor BG, Op de Macks ZA, Rombouts SA, Westenberg PM, Crone EA. Adolescent risky decision-making: Neurocognitive development of reward and control regions. *Neuroimage*. 2010 Feb 24.
19. Van Leijenhorst L, Zanolie K, Van Meel CS, Westenberg PM, Rombouts SA, Crone EA. What motivates the adolescent? Brain regions mediating reward sensitivity across adolescence. *Cereb Cortex*. 2010 Jan; 20(1):61–69. [PubMed: 19406906]
20. Somerville LH, Hare TA, Casey BJ. Frontostriatal maturation predicts behavioral regulation failures to appetitive cues in adolescence. *Journal of Cognitive Neuroscience*. In Press.
21. Ernst M, Pine DS, Hardin M. Triadic model of the neurobiology of motivated behavior in adolescence. *Psychol Med*. 2006 Mar; 36(3):299–312. [PubMed: 16472412]
22. Ernst M, Romeo RD, Andersen SL. Neurobiology of the development of motivated behaviors in adolescence: a window into a neural systems model. *Pharmacol Biochem Behav*. 2009 Sep; 93(3):199–211. [PubMed: 19136024]
23. Steinberg L. A social neuroscience perspective on adolescent risk-taking. *Developmental Review*. 2008; 28:78–106. [PubMed: 18509515]
24. Geier C, Luna B. The maturation of incentive processing and cognitive control. *Pharmacol Biochem Behav*. 2009 Sep; 93(3):212–221. [PubMed: 19593842]
25. Hare TA, Tottenham N, Davidson MC, Glover GH, Casey BJ. Contributions of amygdala and striatal activity in emotion regulation. *Biol Psychiatry*. 2005 Mar 15; 57(6):624–632. [PubMed: 15780849]
26. Levita L, Hare TA, Voss HU, Glover G, Ballon DJ, Casey BJ. The bivalent side of the nucleus accumbens. *Neuroimage*. 2009 Feb 1; 44(3):1178–1187. [PubMed: 18976715]
27. Steinberg L, Graham S, O'Brien L, Woolard J, Cauffman E, Banich M. Age differences in future orientation and delay discounting? *Child Dev*. 2009 Jan-Feb; 80(1):28–44. [PubMed: 19236391]
28. Mischel W, Shoda Y, Rodriguez MI. Delay of gratification in children. *Science*. May 26; 1989 244(4907):933–938. [PubMed: 2658056]
29. Eigsti IM, Zayas V, Mischel W, et al. Predicting cognitive control from preschool to late adolescence and young adulthood. *Psychol Sci*. 2006 Jun; 17(6):478–484. [PubMed: 16771797]
30. Barratt, E.; Patton, J. Impulsivity: cognitive, behavioral, and psychophysiological correlates. In: Zuckerman, M., editor. *Biological bases of sensation seeking, impulsivity, and anxiety*. NJ: Erlbaum, Hillsdale; 1983. p. 77-122.
31. Evenden JL. Varieties of impulsivity. *Psychopharmacology (Berl)*. 1999 Oct; 146(4):348–361. [PubMed: 10550486]
32. Casey, B. Frontostriatal and frontocerebellar circuitry underlying cognitive control. In: Mayr, U.; Owh, E.; Keele, SW., editors. *Developing individuality in the Human Brain*. Washington D.C.: American Psychological Association; 2005.
33. Davidson MC, Amso D, Anderson LC, Diamond A. Development of cognitive control and executive functions from 4 to 13 years: evidence from manipulations of memory, inhibition, and task switching. *Neuropsychologia*. 2006; 44(11):2037–2078. [PubMed: 16580701]
34. Casey BJ, Tottenham N, Liston C, Durston S. Imaging the developing brain: what have we learned about cognitive development? *Trends Cogn Sci*. 2005 Mar; 9(3):104–110. [PubMed: 15737818]
35. Hardin MG, Mandell D, Mueller SC, Dahl RE, Pine DS, Ernst M. Inhibitory control in anxious and healthy adolescents is modulated by incentive and incidental affective stimuli. *J Child Psychol Psychiatry*. 2009 Dec; 50(12):1550–1558. [PubMed: 19573033]
36. Jazbec S, Hardin MG, Schroth E, McClure E, Pine DS, Ernst M. Age-related influence of contingencies on a saccade task. *Exp Brain Res*. 2006 Oct; 174(4):754–762. [PubMed: 16733706]
37. Figner B, Mackinlay RJ, Wilkening F, Weber EU. Affective and deliberative processes in risky choice: age differences in risk taking in the Columbia Card Task. *J Exp Psychol Learn Mem Cogn*. 2009 May; 35(3):709–730. [PubMed: 19379045]
38. Cauffman E, Shulman EP, Steinberg L, et al. Age differences in affective decision making as indexed by performance on the Iowa Gambling Task. *Dev Psychol*. 2010 Jan; 46(1):193–207. [PubMed: 20053017]

39. Crone EA, van der Molen MW. Developmental changes in real life decision making: performance on a gambling task previously shown to depend on the ventromedial prefrontal cortex. *Dev Neuropsychol.* 2004; 25(3):251–279. [PubMed: 15147999]
40. Chassin, L.; Hussong, A.; Barrera, M.; Molina, B.; Trim, R.; Ritter, J. Adolescent substance use. In: Lerner, R.; Steinberg, L., editors. *Handbook of adolescent psychology*. 2nd ed.. New York: Wiley; 2004. p. 665-696.
41. Gardner M, Steinberg L. Peer influence on risk taking, risk preference, and risky decision making in adolescence and adulthood: an experimental study. *Dev Psychol.* 2005 Jul; 41(4):625–635. [PubMed: 16060809]
42. Steinberg L, Albert D, Cauffman E, Banich M, Graham S, Woolard J. Age differences in sensation seeking and impulsivity as indexed by behavior and self-report: evidence for a dual systems model. *Dev Psychol.* 2008 Nov; 44(6):1764–1778. [PubMed: 18999337]
43. Casey BJ, Giedd JN, Thomas KM. Structural and functional brain development and its relation to cognitive development. *Biol Psychol.* 2000 Oct; 54(1–3):241–257. [PubMed: 11035225]
44. Delgado MR. Reward-related responses in the human striatum. *Ann N Y Acad Sci.* 2007 May. 1104:70–88. [PubMed: 17344522]
45. Pasupathy A, Miller EK. Different time courses of learning-related activity in the prefrontal cortex and striatum. *Nature.* 2005 Feb 24; 433(7028):873–876. [PubMed: 15729344]
46. Buckley MJ, Mansouri FA, Hoda H, et al. Dissociable components of rule-guided behavior depend on distinct medial and prefrontal regions. *Science.* 2009 Jul 3; 325(5936):52–58. [PubMed: 19574382]
47. Cardinal RN, Pennicott DR, Sugathapala CL, Robbins TW, Everitt BJ. Impulsive choice induced in rats by lesions of the nucleus accumbens core. *Science.* 2001 Jun 29; 292(5526):2499–2501. [PubMed: 11375482]
48. Gill TM, Castaneda PJ, Janak PH. Dissociable Roles of the Medial Prefrontal Cortex and Nucleus Accumbens Core in Goal-Directed Actions for Differential Reward Magnitude. *Cereb Cortex.* 2010 Apr 1.
49. Galvan A, Hare TA, Davidson M, Spicer J, Glover G, Casey BJ. The role of ventral frontostriatal circuitry in reward-based learning in humans. *J Neurosci.* 2005 Sep 21; 25(38):8650–8656. [PubMed: 16177032]
50. Brenhouse HC, Sonntag KC, Andersen SL. Transient D1 dopamine receptor expression on prefrontal cortex projection neurons: relationship to enhanced motivational salience of drug cues in adolescence. *J Neurosci.* 2008 Mar 5; 28(10):2375–2382. [PubMed: 18322084]
51. Benes FM, Taylor JB, Cunningham MC. Convergence and plasticity of monoaminergic systems in the medial prefrontal cortex during the postnatal period: implications for the development of psychopathology. *Cereb Cortex.* 2000 Oct; 10(10):1014–1027. [PubMed: 11007552]
52. Cunningham MG, Bhattacharyya S, Benes FM. Increasing Interaction of amygdalar afferents with GABAergic interneurons between birth and adulthood. *Cereb Cortex.* 2008 Jul; 18(7):1529–1535. [PubMed: 17971342]
53. Tseng KY, O'Donnell P. Dopamine modulation of prefrontal cortical interneurons changes during adolescence. *Cereb Cortex.* 2007 May; 17(5):1235–1240. [PubMed: 16818475]
54. Seeman P, Bzowej NH, Guan HC, et al. Human brain dopamine receptors in children and aging adults. *Synapse.* 1987; 1(5):399–404. [PubMed: 3505371]
55. Tarazi FI, Baldessarini RJ. Comparative postnatal development of dopamine D(1), D(2) and D(4) receptors in rat forebrain? *Int J Dev Neurosci.* 2000 Feb; 18(1):29–37. [PubMed: 10708903]
56. Teicher MH, Krenzel E, Thompson AP, Andersen SL. Dopamine receptor pruning during the peripubertal period is not attenuated by NMDA receptor antagonism in rat. *Neurosci Lett.* 2003 Mar 20; 339(2):169–171. [PubMed: 12614921]
57. Andersen SL, Thompson AT, Rutstein M, Hostetter JC, Teicher MH. Dopamine receptor pruning in prefrontal cortex during the periadolescent period in rats. *Synapse.* 2000 Aug; 37(2):167–169. [PubMed: 10881038]
58. Weickert CS, Webster MJ, Gondipalli P, et al. Postnatal alterations in dopaminergic markers in the human prefrontal cortex. *Neuroscience.* 2007 Feb 9; 144(3):1109–1119. [PubMed: 17123740]

59. Fonseca, Rd; Ramos, JA.; Bonnin, A.; Fernandez-Ruiz, JJ. Presence of cannabinoid binding sites in the brain from early postnatal ages. *NeuroReport*. 1993; 4(2):135–138. [PubMed: 8453049]
60. Gardner, EL. The neurobiology and genetics of addiction: implications of the "reward deficiency syndrome" for therapeutic strategies in chemical dependency. In: Elster, J., editor. *Addiction: entries and exists*. New York: Russell Sage; 1999. p. 57-119.
61. Robinson TE, Berridge KC. The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Brain Res Rev*. 1993 Sep-Dec; 18(3):247–291. [PubMed: 8401595]
62. Volkow ND, Swanson JM. Variables that affect the clinical use and abuse of methylphenidate in the treatment of ADHD. *Am J Psychiatry*. 2003 Nov; 160(11):1909–1918. [PubMed: 14594733]
63. Robinson TE, Berridge KC. Addiction. *Annu Rev Psychol*. 2003; 54:25–53. [PubMed: 12185211]
64. Spear, L. *The Behavioral Neuroscience of Adolescence*. New York: W.W. Norton & Company; 2009.
65. Forbes EE, Ryan ND, Phillips ML, et al. Healthy adolescents' neural response to reward: associations with puberty, positive affect, and depressive symptoms. *J Am Acad Child Adolesc Psychiatry*. 2010 Feb; 49(2):162–172. e161–e165. [PubMed: 20215938]
66. Romeo RD. Puberty: a period of both organizational and activational effects of steroid hormones on neurobehavioural development. *J Neuroendocrinol*. 2003 Dec; 15(12):1185–1192. [PubMed: 14636181]
67. Forbes EE, Dahl RE. Pubertal development and behavior: hormonal activation of social and motivational tendencies. *Brain Cogn*. 2010 Feb; 72(1):66–72. [PubMed: 19942334]
68. Liston C, Watts R, Tottenham N, et al. Frontostriatal microstructure modulates efficient recruitment of cognitive control. *Cerebral Cortex*. 2006 Apr; 16(4):553–560. [PubMed: 16033925]
69. Casey BJ, Epstein JN, Buhle J, et al. Frontostriatal connectivity and its role in cognitive control in parent-child dyads with ADHD. *Am J Psychiatry*. 2007 Nov; 164(11):1729–1736. [PubMed: 17974939]
70. Asato MR, Terwilliger R, Woo J, Luna B. White Matter Development in Adolescence: A DTI Study. *Cereb Cortex*. 2010 Jan 5.
71. Durston S, Davidson MC, Tottenham N, et al. A shift from diffuse to focal cortical activity with development. *Dev Sci*. 2006 Jan; 9(1):1–8. [PubMed: 16445387]
72. Luna B, Padmanabhan A, O'Hearn K. What has fMRI told us about the development of cognitive control through adolescence. *Brain and Cognition*. 2010
73. Astle DE, Scerif G. Using developmental cognitive neuroscience to study behavioral and attentional control. *Developmental Psychobiology*. 2009; 51(2):107–118. [PubMed: 18973175]
74. Luna B, Thulborn KR, Munoz DP, et al. Maturation of widely distributed brain function subserves cognitive development. *Neuroimage*. 2001 May; 13(5):786–793. [PubMed: 11304075]
75. Bunge SA, Dudukovic NM, Thomason ME, Vaidya CJ, Gabrieli JD. Immature frontal lobe contributions to cognitive control in children: evidence from fMRI. *Neuron*. 2002 Feb 17; 33(2):301–311. [PubMed: 11804576]
76. Bitan T, Burman DD, Lu D, et al. Weaker top-down modulation from the left inferior frontal gyrus in children. *Neuroimage*. 2006; 33:991–998. [PubMed: 16978881]
77. Tamm L, Menon V, Reiss AL. Maturation of brain function associated with response inhibition. *J Am Acad Child Adolesc Psychiatry*. 2002 Oct; 41(10):1231–1238. [PubMed: 12364845]
78. Stevens MC, Skudlarski P, Pearlson GD, Calhoun VD. Age-related cognitive gains are mediated by the effects of white matter development on brain network integration. *Neuroimage*. 2009 Dec; 48(4):738–746. [PubMed: 19577651]
79. Giedd JN, Blumenthal J, Jeffries NO, et al. Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci*. 1999 Oct; 2(10):861–863. [PubMed: 10491603]
80. Huttenlocher PR. Morphometric study of human cerebral cortex development. *Neuropsychologia*. 1990; 28(6):517–527. [PubMed: 2203993]
81. Casey BJ, Thomas KM, Davidson MC, Kunz K, Franzen PL. Dissociating striatal and hippocampal function developmentally with a stimulus-response compatibility task. *The Journal of Neuroscience*. 2002; 22(19):8647–8652. [PubMed: 12351738]

82. Bjork JM, Knutson B, Fong GW, Caggiano DM, Bennett SM, Hommer DW. Incentive-elicited brain activation in adolescents: similarities and differences from young adults. *Journal of Neuroscience*. 2004 Feb 25; 24(8):1793–1802. [PubMed: 14985419]
83. Delgado MR, Nystrom LE, Fissell C, Noll DC, Fiez JA. Tracking the hemodynamic responses to reward and punishment in the striatum. *Journal of Neurophysiology*. 2000 Dec; 84(6):3072–3077. [PubMed: 11110834]
84. Kuhnen CM, Knutson B. The neural basis of financial risk taking. *Neuron*. 2005 Sep 1; 47(5):763–770. [PubMed: 16129404]
85. Matthews SC, Simmons AN, Lane SD, Paulus MP. Selective activation of the nucleus accumbens during risk-taking decision making. *Neuroreport*. 2004 Sep 15; 15(13):2123–2127. [PubMed: 15486494]
86. Casey BJ, Galvan A, Hare TA. Changes in cerebral functional organization during cognitive development. *Current Opinion in Neurobiology*. 2005 Apr; 15(2):239–244. [PubMed: 15831409]
87. Boes AD, Bechara A, Tranel D, Anderson SW, Richman L, Nopoulos P. Right ventromedial prefrontal cortex: a neuroanatomical correlate of impulse control in boys. *Soc Cogn Affect Neurosci*. 2009 Mar; 4(1):1–9. [PubMed: 19015086]
88. Vaidya CJ, Austin G, Kirkorian G, et al. Selective effects of methylphenidate in attention deficit hyperactivity disorder: a functional magnetic resonance study. *Proc Natl Acad Sci U S A*. 1998 Nov 24; 95(24):14494–14499. [PubMed: 9826728]
89. Epstein JN, Casey BJ, Tonev ST, et al. ADHD- and medication-related brain activation effects in concordantly affected parent-child dyads with ADHD. *Journal of Child Psychology and Psychiatry*. 2007; 48(9):899–913. [PubMed: 17714375]
90. Scheres A, Milham MP, Knutson B, Castellanos FX. Ventral striatal hypo-responsiveness during reward anticipation in attention-deficit/hyperactivity disorder. *Biol Psychiatry*. 2007 Mar 1; 61(5):720–724. [PubMed: 16950228]
91. Bjork JM, Knutson B, Hommer DW. Incentive-elicited striatal activation in adolescent children of alcoholics. *Addiction*. 2008 Aug; 103(8):1308–1319. [PubMed: 18851716]
92. Hardin MG, Ernst M. Functional brain imaging of development-related risk and vulnerability for substance use in adolescents. *J Addict Med*. 2009 Jun 1; 3(2):47–54. [PubMed: 20161036]
93. Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE. Monitoring the future national results on adolescent drug use: overview of key findings, 2008. Bethesda, MD: National Institute on Drug Abuse; 2009. NIH Publication No. 09-7401 ed
94. Witt ED. Research on alcohol and adolescent brain development: opportunities and future directions. *Alcohol*. 2010 Feb; 44(1):119–124. [PubMed: 20113880]
95. Grant BF, Dawson DA. Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: results from the National Longitudinal Alcohol Epidemiologic Survey. *J Subst Abuse*. 1997; 9:103–110. [PubMed: 9494942]
96. Li TK, Hewitt BG, Grant BF. Is there a future for quantifying drinking in the diagnosis, treatment, and prevention of alcohol use disorders? *Alcohol Alcohol*. 2007 Mar-Apr; 42(2):57–63. [PubMed: 17307789]
97. Volkow ND, Fowler JS, Wang GJ, Goldstein RZ. Role of dopamine, the frontal cortex and memory circuits in drug addiction: insight from imaging studies. *Neurobiol Learn Mem*. 2002 Nov; 78(3):610–624. [PubMed: 12559839]
98. Maldonado R, Rodriguez de Fonseca F. Cannabinoid addiction: behavioral models and neural correlates. *J Neurosci*. 2002 May 1; 22(9):3326–3331. [PubMed: 11978807]
99. French ED, Dillon K, Wu X. Cannabinoids excite dopamine neurons in the ventral tegmentum and substantia nigra. *Neuroreport*. 1997 Feb 10; 8(3):649–652. [PubMed: 9106740]
100. Robinson TE, Berridge KC. The incentive sensitization theory of addiction: some current issues. *Philos Trans R Soc Lond B Biol Sci*. 2008 Oct 12; 363(1507):3137–3146. Review. [PubMed: 18640920]
101. Doremus TL, Brunell SC, Varlinskaya EI, Spear LP. Anxiogenic effects during withdrawal from acute ethanol in adolescent and adult rats. *Pharmacol Biochem Behav*. 2003 May; 75(2):411–418. [PubMed: 12873633]

102. Spear LP, Varlinskaya EI. Adolescence. Alcohol sensitivity, tolerance, and intake. *Recent Dev Alcohol*. 2005; 17:143–159. [PubMed: 15789864]
103. Pautassi RM, Myers M, Spear LP, Molina JC, Spear NE. Adolescent but not adult rats exhibit ethanol-mediated appetitive second-order conditioning. *Alcohol Clin Exp Res*. 2008 Nov; 32(11):2016–2027. [PubMed: 18782343]
104. Varlinskaya EI, Spear LP. Acute effects of ethanol on social behavior of adolescent and adult rats: role of familiarity of the test situation. *Alcohol Clin Exp Res*. 2002 Oct; 26(10):1502–1511. [PubMed: 12394283]
105. White AM, Swartzwelder HS. Age-related effects of alcohol on memory and memory-related brain function in adolescents and adults. *Recent Dev Alcohol*. 2005; 17:161–176. [PubMed: 15789865]
106. Sircar R, Basak AK, Sircar D. Repeated ethanol exposure affects the acquisition of spatial memory in adolescent female rats. *Behav Brain Res*. 2009 Sep 14; 202(2):225–231. [PubMed: 19463705]
107. Sircar R, Sircar D. Adolescent rats exposed to repeated ethanol treatment show lingering behavioral impairments. *Alcohol Clin Exp Res*. 2005 Aug; 29(8):1402–1410. [PubMed: 16131847]
108. Carpenter-Hyland EP, Chandler LJ. Adaptive plasticity of NMDA receptors and dendritic spines: implications for enhanced vulnerability of the adolescent brain to alcohol addiction. *Pharmacol Biochem Behav*. 2007 Feb; 86(2):200–208. [PubMed: 17291572]
109. De Bellis MD, Clark DB, Beers SR, et al. Hippocampal volume in adolescent-onset alcohol use disorders. *Am J Psychiatry*. 2000 May; 157(5):737–744. [PubMed: 10784466]
110. Nagel BJ, Schweinsburg AD, Phan V, Tapert SF. Reduced hippocampal volume among adolescents with alcohol use disorders without psychiatric comorbidity. *Psychiatry Res*. 2005 Aug 30; 139(3):181–190. [PubMed: 16054344]
111. Brown SA, Tapert SF. Adolescence and the trajectory of alcohol use: basic to clinical studies. *Ann N Y Acad Sci*. 2004 Jun; 1021:234–244. [PubMed: 15251893]
112. Medina KL, McQueeney T, Nagel BJ, Hanson KL, Schweinsburg AD, Tapert SF. Prefrontal cortex volumes in adolescents with alcohol use disorders: unique gender effects. *Alcohol Clin Exp Res*. 2008 Mar; 32(3):386–394. [PubMed: 18302722]
113. McQueeney T, Schweinsburg BC, Schweinsburg AD, et al. Altered white matter integrity in adolescent binge drinkers. *Alcohol Clin Exp Res*. 2009 Jul; 33(7):1278–1285. [PubMed: 19389185]
114. Tapert SF, Cheung EH, Brown GG, et al. Neural response to alcohol stimuli in adolescents with alcohol use disorder. *Arch Gen Psychiatry*. 2003 Jul; 60(7):727–735. [PubMed: 12860777]
115. Pulido C, Brown SA, Cummins K, Paulus MP, Tapert SF. Alcohol cue reactivity task development. *Addict Behav*. 2010 Feb; 35(2):84–90. [PubMed: 19800172]
116. Monti PM, Miranda R Jr, Nixon K, et al. Adolescence: booze, brains, and behavior. *Alcohol Clin Exp Res*. 2005 Feb; 29(2):207–220. [PubMed: 15714044]
117. Schweinsburg AD, Paulus MP, Barlett VC, et al. An fMRI study of response inhibition in youths with a family history of alcoholism. *Ann N Y Acad Sci*. 2004 Jun; 1021:391–394. [PubMed: 15251915]
118. Deckel AW, Hesselbrock V. Behavioral and cognitive measurements predict scores on the MAST: a 3-year prospective study. *Alcohol Clin Exp Res*. 1996 Oct; 20(7):1173–1178. [PubMed: 8904966]
119. Tarter RE, Kirisci L, Mezzich A, et al. Neurobehavioral disinhibition in childhood predicts early age at onset of substance use disorder. *Am J Psychiatry*. 2003 Jun; 160(6):1078–1085. [PubMed: 12777265]
120. Behar D, Berg CJ, Rapoport JL, et al. Behavioral and physiological effects of ethanol in high-risk and control children: a pilot study. *Alcohol Clin Exp Res*. 1983 Fall; 7(4):404–410. [PubMed: 6318590]
121. Dahl RE. Affect regulation, brain development, and behavioral/emotional health in adolescence. *CNS Spectr*. 2001 Jan; 6(1):60–72. [PubMed: 17008832]

122. Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *American Journal of Psychiatry*. 2003 Jun; 160(6):1041–1052. [PubMed: 12777258]
123. Nelson EE, Leibenluft E, McClure EB, Pine DS. The social re-orientation of adolescence: a neuroscience perspective on the process and its relation to psychopathology. *Psychol Med*. 2005 Feb; 35(2):163–174. [PubMed: 15841674]
124. Mannuzza S, Klein RG. Long-term prognosis in attention-deficit/hyperactivity disorder. *Child Adolesc Psychiatr Clin N Am*. 2000 Jul; 9(3):711–726. [PubMed: 10944664]
125. Mischel W, Underwood B. Instrumental ideation in delay of gratification. *Child Dev*. 1974 Dec; 45(4):1083–1088. [PubMed: 4143856]
126. Ayduk O, Mendoza-Denton R, Mischel W, Downey G, Peake PK, Rodriguez M. Regulating the interpersonal self: strategic self-regulation for coping with rejection sensitivity. *J Pers Soc Psychol*. 2000 Nov; 79(5):776–792. [PubMed: 11079241]
127. Tripodi SJ, Bender K, Litschge C, Vaughn MG. Interventions for reducing adolescent alcohol abuse: a meta-analytic review. *Arch Pediatr Adolesc Med*. 2010 Jan; 164(1):85–91. [PubMed: 20048247]
128. Somerville LH, Casey B. Developmental neurobiology of cognitive control and motivational systems. *Curr Opin Neurobiol*. 2010 Feb 16.

Figure 1. Cartoon model of ventral striatal and prefrontal cortex (PFC) interactions across development. Deeper color indicates greater regional signaling. Line represents functional connectivity, with solid line indicating mature connection and dotted line indicating immaturity (from ¹²⁸).

Figure 2.

Illustration of different developmental courses for sensation-seeking and impulsivity. Panel A. Plot of sensation-seeking and impulsivity as a function of age (adapted from ⁴²). Panel B. Plot of patterns of activity in brain regions sensitive to reward outcomes during a cognitive control task across development (adapted from ^{6, 16}).

Figure 3. Ventral Striatal activity to reward and association with risk- taking. Localization of ventral striatum in axial plane (left panel), which is activated with reward (middle panel) and correlated with risk-taking (right panel) (adapted from ^{6, 16})