PamI and PamII restriction endonucleases from *Phormidium ambiguum*

Stanislaw Piechula, Sun C.Kim¹ and Anna J.Podhajska

Department of Microbiology, University of Gdansk, 80-822 Gdansk, ul. Kladki 24, Poland and

¹McArdle Laboratory for Cancer Research, The University of Wisconsin, 1400 University Avenue, Madison, WI 53076, USA

Submitted November 27, 1991

PamI and PamII are a type II restriction endonucleases from the Cyanobacterial strain *Phormidium ambiguum* GOM (CCALA Hindak 1965/117). PamI and PamII are isoschizomers of MstI and AcyI respectively (1). The enzymes were purified using two chromatographic steps: 1) phosphocellulose, 2) DEAE-sephadex G-25. The enzymes were free of contaminating nucleases activity. All digestions were performed at 37°C in a buffer containing 50 mM NaCl, 10 mM Tris-HCl (pH 7.5), 10 mM MgCl.

Parallel digestion of a standard substrate (Figure 1) and sequencing (Figures 3 and 4) confirmed that PamI is an isoschizomer of MstI (5'-TGC/GCA-3') and PamII is an isoschizomer of AcyI (5'-GR/CGYC-3'). The cleavage sites of PamI and PamII were determined as follows:

PamI: Two fragments (of about 1000 and 1663 bp) were isolated from pUC19 digested with PamI enzyme (Figure 2), and both fragments were treated with PolIK and dNTP. Both untreated- and filled-in DNA fragments were used to clone into SmaI in M13mp18 and both DNA fragments were clonable. The enzyme must therefore generate blunt ends. The clones containing the inserts were sequenced by the Sanger *et al.* (1977) method. The sequence immediately after 5'-CCC of the SmaI cut site was always 5'-GCA and that before 5'-GGG of the SmaI cut was 5'-TGC. The recognition site must therefore be 5'-TGCGCA-3' and the cut site is in the middle of the sequence, resulting in blunt ends as the sequences of both ends of the cut are symmetric (Figure 3).

PamII: Three different fragments (of about 300, 400 and 2000 bp) were isolated from pUC19 digested with PamII (Figure 2). The DNA fragments were treated in exactly the same way as the PamI digested fragments. However, only the DNA fragments treated which PolIK and dNTP were clonable which indicated that the enzyme does not generate blunt ends. The sequence immediately after 5'-CCC of the SmaI cut site was consistently either 5'-GACG or 5'-CGCC and the sequence before 5'-GGG of the SmaI cut site was either 5'-CGTC or 5'-CGCC (Figure 4). The CG sequence was repeated. The enzyme must therefore produce 2-nt 5' protruding cohesive ends, and considering the neighbouring sequence of the repeated sequence 5'-CG, the enzyme cut site must be GR\CGYC.

REFERENCE

1. Roberts, J.R. (1990) *Nucleic Acids Res.* **18**, 2331-2365.

Figure 1. Agarose gel electrophoresis of lambda DNA digested by: lane 1, PamI; lane 2, FspI (FspI is a commercial available isoschizomer of MstI); lane 3, PamII; lane 4, AhaII (AhaII is a commercial available isoschizomer of AcyI); lane 5, HindIII.

Figure 2. Polyacrylamide gel electrophoresis of pUC19 DNA digested by: lane 1, PamI; lane 2, FspI; lane 3, PamII; lane 4, AhaII; lane 5, TaqI.

Figure 3. Determination of PamI restriction specificity: autoradiography of the sequencing ladders representing the nucleotide sequence in the junction of the cloned fragment and M13mp18 DNA.

Figure 4. Determination of PamII restriction specificity: autoradiography of the sequencing ladders representing the nucleotide sequence in the junction of the cloned fragment and M13mp18 DNA.