CORRESPONDENCE

Malnutrition in Hospital: The Clinical and Economic Implications

by Prof. Dr. med. Christian Löser in volume 51–52/2010

Not Hungry Today

This article was long overdue, and the authors deserve congratulations. Among the causes of possible malnutrition, the lack of attention of the nursing staff should have been explicitly mentioned, or the fact that a person who can devote time to feeding patients was lacking. The reasons for why patients did not touch their food were not elicited. The food was cleared away by staff saying, “Oh, are we not hungry today?” Possible reasons: the patient was not able to reach the food (bed-side table on the wrong side of the bed, too far away). S/he was not able to butter their own bread (arm in plaster, infusion), or the bite-sized portions were too large. The headboard of the bed was not elevated; the patient did not see the food; the patient did not want to spill food over themselves (disgust, not wanting to annoy the nurses); perhaps the patient was not given a large napkin or such like. Furthermore, for patients lying in bed it is easier and safer to consume some foods with a tablespoon, rather than a teaspoon or fork. Tough bread rolls and unseasoned or cold meals are difficult to swallow. The patient would like to eat the fish, but someone “forgot” to put their spectacles on for them to enable them to see the small bones—fish fingers would offer a way out of this dilemma. Perhaps the patient’s choice of food was not considered and they were given exactly what they have disliked for the preceding 20 years. Paying attention to all these details does not incur any additional costs, creates satisfaction in patients and nurses, and improves the reputation of a hospital. A win-win situation for everyone.

DOI: 10.3238/arztebl.2011.0354a

REFERENCES

1. Löser C: Malnutrition in hospital—the clinical and economic implications. Dtsch Arztebl Int 2010; 107(51–52): 911–7.

Dr. med. R. Michael Stiebing
Bielefeld, Germany
rmstie@web.de

Bioimpedance Analysis

I read the article on malnutrition in hospital with great interest—I work in a hospital that has inpatients that have intentionally entered a state of malnutrition: patients with eating disorders such as bulimia or anorexia.

In order to assess the severity of the malnutrition, bioimpedance analysis has become the trusted standard

in our hospital. By measuring body cell mass (BCM, metabolically active mass or muscle mass), body fat, and body water it is possible to ascertain the extent to which body fat reserves are depleted, whether the patient is in a catabolic state with compensatory water retention or whether they are acting in a manipulative, self harming manner. By means of this test we then decide systematically on the basis of a certain stepwise scheme (phase angle, ECM/BCM index) whether a protein-rich diet (*Figure 2* Steps II or III in the article) is sufficient or whether parenteral nutrition (*Figure 2* Steps V or VI in the article) is required.

DOI: 10.3238/arztebl.2011.0354b

REFERENCES

1. Löser C: Malnutrition in hospital—the clinical and economic implications. Dtsch Arztebl Int 2010; 107(51–52): 911–7.

Dr. med. Monika Gebel
Psychotherapeutisches Zentrum
Kitzberg-Klinik
Bad Mergentheim, Germany
m.gebel@ptz.de

Treatment Principles for Decubitus Ulcers

In our hospital we have an intensive focus on malnutrition especially in the context of patients with decubitus ulcers, in particular geriatric or paralyzed patients. The “six treatment principles” for treating chronic decubitus ulcers (2) described by our working group is based on interdisciplinary, multiprofessional collaboration between conservative and surgical specialties—not least because of the multifactorial etiology of decubitus ulcers—and its main principle is the investigation of the causes and **causative** treatment of possibly comorbid malnutrition, which has to be dealt with for the decubitus ulcers to be treated successfully. We determine patients’ nutritional status before plastic surgery in the shape of a so called mini-lympho-nutrogram, which includes the values for albumin, transferrin, zinc, vitamin B12, folic acid, and the lymphocyte count (2). On the basis of the experiences gained in our hospital we can highly recommend the use of the nutrogram, as described by Seiler et al (3) in clinical practice because of our therapeutic successes in our specialty. In our hospital we have no experience with either the SGA (Subjective Global Assessment Score) or NRS (Nutritional Risk Score), but we see clear overlaps with our own assessment of a patient’s nutritional status, because the combination of nutrogram and medical history probably results in a comparable assessment. We mostly administer liquid foods and additional foods perioperatively, percutaneous endoscopic gastrostomy (PEG) tubes are used only rarely (for example, in patients having chemotherapy who have developed decubitus ulcers).

In the context of post-bariatric body contouring after massive weight loss, especially after obesity surgery to induce weight loss, these patients—often still classed as overweight—are often subject to substantial malnutrition, which has to be compensated perioperatively before possible surgical interventions in order to avoid complications.

In sum, I can only agree with Löser: nutritional medicine touches on almost all medical specialties and will have to be made an integral part of medical training for students and specialty doctors.

DOI: 10.3238/arztebl.2011.0354c

REFERENCES

1. Löser C: Malnutrition in hospital—the clinical and economic implications. *Dtsch Arztebl Int* 2010; 107(51–52): 911–7.
2. Rieger U, et al.: Six treatment principles of the basle pressure sore concept. *Handchir Mikrochir Plast Chir* 2007; 39: 206–14.
3. Seiler WO, Stahelin HB: Search for factors preventing wound healing. A motivating therapy concept in chronic decubitus ulcers. *Krankenpf J* 1999; 37: 317–23.

PD Dr. med. Ulrich Michael Rieger

Univ.-Klinik für Plastische, Rekonstruktive & Ästhetische Chirurgie
Medizinische Universität Innsbruck
Innsbruck, Tirol, Österreich
ulrich.rieger@i-med.ac.at

In Reply:

The many positive responses to my article confirm the great clinical relevance of the subject and the interest in the problem area that is malnutrition in hospital. This is true not only for the known clinical sequelae but also the proven consequences in terms of the business finances, and it shows convincingly that modern nutritional medicine is a highly effective and integral part of medical therapy and prevention, for which awareness and acceptance in Germany are fortunately rising.

I thank Rieger, Stiebing, and Gebel for their additional comments. Dr Monika Gebel emphasizes rightly that in patients with chronic diseases with severe malnutrition and especially in pronounced eating disorders, bioimpedance analysis (BIA) provides a very detailed

description of the individual body composition and forms a basis and observational method for the targeted nutritional medical care for these patients. With this indication it should be used generously in clinical practice; but BIA is too elaborate to be used as a screening tool in everyday clinical practice.

Dr Michael Stiebing rightly reminds us that doctors as well as nursing staff and family members need to pay attention to the complex factors that lead to malnutrition and that have to be borne in mind if individual treatment is to be adequate. In everyday clinical practice, it is often possible to achieve enormous successes by using measures that may strike one as banal at first glance. Many scientific studies have shown a high efficacy for such simple, basic measures. These need to be heeded and implemented by hospitals, old people's residential homes, and nursing and care homes. Dr Ulrich Rieger's comments confirm how important methodical nutritional medical care can be as an integral part of the complex interdisciplinary treatment principles especially in the surgical disciplines. Interdisciplinary treatment for chronic decubitus ulcers and perioperative conditioning are only two of the scientifically well supported examples for many relevant clinical situations.

I hope that my article, which has been so positively received, is a further contribution to making nutritional medical interventions gain the clinical importance in Germany that is long overdue in the context of modern, evidence based, medical care for patients, as is supported by a multitude of scientific data.

DOI: 10.3238/arztebl.2011.0355

REFERENCES

1. Löser C: Malnutrition in hospital—the clinical and economic implications. *Dtsch Arztebl Int* 2010; 107(51–52): 911–7.

Prof. Dr. med. Christian Löser

Medizinischen Klinik
Rotes Kreuz Krankenhaus, Kassel
chr.loeser@rkh-kassel.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.