Published in final edited form as:

Org Lett. 2011 June 17; 13(12): 3114–3117. doi:10.1021/ol2010227.

Oppolzer-Type Intramolecular Diels-Alder Cycloadditions via Isomerizations of Allenamides

John B. Feltenberger and Richard P. Hsung*

Division of Pharmaceutical Sciences and Department of Chemistry, University of Wisconsin, Madison, WI 53705

Abstract

A new approach to Oppolzer's intramolecular Diels-Alder cycloaddition [IMDA] through γ -isomerization of readily available *N*-tethered allenamides is described. These IMDA reactions are carried out in tandem with the allenamide isomerization or 1,3-H shift, leading to complex nitrogen heterocycles in a highly stereoselective manner.

Intermolecular Diels-Alder cycloadditions of 1-amino- or 1-amido-dienes^{1–7} have proven to be a powerful reaction manifold with some notable examples shown in Scheme 1. Although Oppolzer^{8,9} had elegantly demonstrated the concept of *N*-tethered 1-amido-dienes in intramolecular Diels-Alder cycloadditions [IMDA],¹⁰ relative to the intermolecular pursuit, it has shown limited applications in the last forty years in part due to a lack of efficient synthetic access to these cycloaddition precursors.^{10,11} We recently reported¹² that γ -isomerizations of allenamides **1** provide a facile entry to 1-amido-dienes **3** via a stereoselective 1,3-H shift [Scheme 1].¹³ This finding provokes a unique synthetic design in which the allenamide isomerization could be rendered in tandem with the Oppolzer-type intramolecular Diels-Alder cycloaddition when using *N*-tethered allenamides **4**. If successful [**4**→**5**→**6**], such an endeavor would allow rapid assembly of structural complexity from simple allenamides and further underscores the significance of developing chemistry of allenamides.^{14–16} We communicate here an Oppolzer-type IMDA through γ -isomerizations of allenamides.

While the general plan was laid out concisely in Scheme 2 with allenamides **11** and **12**¹⁷ readily attainable through amidative cross-coupling^{1f,18} of allenyl iodide **7** with respective amides **8** and **9**, developing an effective γ -isomerization in conjunction with Oppolzer's IMDA was not trivial; and a wide range of conditions had to be examined [Table 1]. While the modest yields attained under a number of conditions [entries 2–8] could still be

* rhsung@wisc.edu .

Supporting Information Available: Experimental procedures as well as NMR spectra, characterizations, and X-ray structural files are available for all new compounds and free of charge via Internet <http://pubs.acs.org>.

considered useful given the context of constructing multiple bonds and stereocenters in a tandem sequence, ultimately, it appeared that a base, optimally being proton sponge [entry 11] or Et₃N [entry 14], was needed to avoid significant hydrolysis of starting allenamides. Other amine bases, inorganic bases, or drying agents screened were not effective [entries 15–20].

In addition, a *para*-nosyl substituent on the nitrogen atom provided higher yield of the respective cycloadduct **16** [entry 14], relative to cycloadduct **15** containing an *N*-Ts group [entry 13]. Diastereoselectivity or the *endo-exo* ratio for **16** [*dr* = 10:1] was also slightly higher than that of **15** [*dr* = 8:1]. Relative stereochemistry of the major isomer of **16** was unambiguously assigned using single crystal X-ray structure [Figure 1], thereby suggesting a favored *endo*-transition state with respect to the carbon tethering.

A quick comparison was drawn through pursuing this Oppolzer-type IMDA sequentially in conjunction with the γ -isomerization of allenamides. As shown in Scheme 3, acid catalyzed isomerization of allenamide **12** gave *N*-tethered 1-amido diene **14** in near quantitative yield. The ensuing cycloaddition using either Et₃N or proton sponge as the additive led to cycloadduct **16** with a combined yield for either condition being very comparable to those attained in a tandem manner [in italics].

Having established the feasibility, various examples could be readily produced including tricycles **21–23** [see product Figure 2]. In general, the *para*-nosyl substitution gave better yields, and both stereoselectivity and overall yields are in the synthetically useful range and could be considered excellent in the context of constructing multiple bonds and stereocenters. It is noteworthy that additional substitutions on the olefin such as Ph group could be included, leading to cycloadduct **23** with an *endo to exo* ratio of 10:3 [the *endo* isomer was assigned using NOE experiments]. In addition, the original *cis* configuration of the olefin was preserved during the cycloaddition.

We then turned our attention toward constructing heterocycles with greater complexity. However, when using allenamide **24** containing a cinnamate motif in an attempt to prepare indolines through the tandem γ -isomerization-IMDA sequence, we encountered an unexpected enamide 1,4-addition [see **25**], leading to the formation of amido diene **27** [Scheme 4]. This new diene was only fully recognized after we were attempting an aromatization process using DDQ, which led to another Diels-Alder cycloadduct **28**.

Contemplating to take advantage of this new observation we prepared allenamide **29** containing an alloc group with the hope that another IMDA could take place through diene **30** leading to tetracycle **31** [Scheme 5]. Surprisingly, we found that 1,4-addition was even occurring prior to the allenamide isomerization, leading to the formation of cyclobutane **33**. Cyclobutane **33** most likely was derived from two consecutive 1,4-additions with latter involving vinyl iminium ion **32**. On the other hand, although being the major product, 1-amido-diene **31** did not undergo further IMDA to give the anticipated tetracyclic quinoline **31**. Fortunately, this issue could be circumvented by using allenamide **34** and **34'**, which gave respective indolines **36** and **36'** in modest overall yield through the expected sequence of tandem γ -isomerization–Oppolzer-type IMDA. Unfortunately, the *endo-exo* ratios for these cases were not high.

The power of this tandem sequence can be manifested in the quinoline synthesis. As shown in Scheme 6, a number of quinolines such as **41–43** could be accessed rapidly in an efficient manner from simple *N*-allyl-aniline **37** in four steps. These rapid assemblies feature aza-Claisen, copper-catalyzed amidative cross-coupling, and tandem γ -isomerization–Oppolzer-type intramolecular Diels-Alder cycloaddition. In particular, the tandem sequence is stereoselective for the synthesis of the tetracyclic quinoline **43**, which was isolated in 74%

yield with a diastereomeric ratio of 10:1. The major isomer of **43** was concisely assigned using NOE experiment.

We have described here an Oppolzer-type intramolecular Diels-Alder cycloaddition [IMDA] through γ -isomerization of readily available *N*-tethered allenamides. These IMDA reactions are carried out in tandem with the allenamide isomerization, which consists of a 1,3-H-shift, leading to complex nitrogen heterocycles in a highly stereoselective manner. Applications of this new tandem process in alkaloid synthesis are underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Authors thank NIH [GM066055] for support and Dr. Victor Young [University of Minnesota] for X-ray structural analysis.

References

1. For reviews on chemistry of dienamides, see: (a) Overman LE. *Acc. Chem. Res.* 1980; 13:218.. (b) Petrzilka M, Grayson JJ. *Synthesis*. 1981:753.. (c) Campbell AL, Lenz GR. *Synthesis*. 1987:421.. Also see: (d) Krohn K. *Angew. Chem. Int. Ed.* 1993; 32:1582.. (e) Enders D, Meyer O. *Liebigs Ann.* 1996:1023.. (f) Tracey MR, Hsung RP, Antoline J, Kurtz KCM, Shen L, Slafer BW, Zhang Y, Weinreb, Steve M. Chapter 21.4. *Science of Synthesis, Houben-Weyl Methods of Molecular Transformations*. 2005 Georg Thieme Verlag KG
2. For early applications of *N*-substituted dienes, see: (a) Hünig S, Kahanek H. *Chem. Ber.* 1957; 90:238.. (b) Terada A, Murata K. *Bull. Chem. Soc. Jpn.* 1967; 40:1644.. (c) Dauben WG, Kozikowski AP. *J. Am. Chem. Soc.* 1974; 96:3664.
3. Oppolzer W, Bieber L, Francotte E. *Tetrahedron Lett.* 1979:4537.
4. (a) Overman LE, Clizbe LA. *J. Am. Chem. Soc.* 1976; 98:2352. (b) Overman LE, Freerks RL, Petty CB, Clizbe LA, Ono RK, Taylor GF, Jssup. *J. Am. Chem. Soc.* 1981; 103:2816. references cited therein.
5. (a) Smith AB III, Wexler BA, Tu C-Y, Konopelski JP. *J. Am. Chem. Soc.* 1985; 107:1308. (b) Schlessinger RH, Pettus TRR, Springer JP, Hoogsteen K. *J. Org. Chem.* 1994; 59:3246. (c) Kozmin SA, Rawal VH. *J. Am. Chem. Soc.* 1997; 119:7165. and references cited therein. (d) Snowden RL. *Tetrahedron Lett.* 1984; 25:3835.
6. (a) Huang Y, Iwama T, Rawal VH. *J. Am. Chem. Soc.* 2000; 122:7843. (b) Kozmin SA, Iwama T, Huang Y, Rawal VH. *J. Am. Chem. Soc.* 2002; 124:4628. [PubMed: 11971711]
7. (a) Grauvry N, Huet F. *J. Org. Chem.* 2001; 66:583. [PubMed: 11429833] (b) Robiette R, Cheboub-Benchaba K, Peeters D, Marchand-Brynaert J. *J. Org. Chem.* 2003; 68:9809. [PubMed: 14656111] (c) von Wangelin AJ, Neumann H, Goirides D, Spannenberg A, Beller M. *Org. Lett.* 2001; 3:2895. [PubMed: 11529784]
8. For examples of tandem benzocyclobutene ring-opening-1-amidodiene Diels-Alder sequences, see: Oppolzer W. *J. Am. Chem. Soc.* 1971; 93:3834.. (b) Oppolzer W, Keller K. *J. Am. Chem. Soc.* 1971; 93:3836.. (c) Oppolzer W, Robbani C. *Helv. Chim. Acta.* 1983; 66:1119.. For a recent example and for references cited therein, see: (d) Feltenberger JB, Hayashi R, Tang Y, Babiash ESC, Hsung RP. *Org. Lett.* 2009; 11:3666. [PubMed: 19624104]
9. (a) Oppolzer W, Fröstl W. *Helv. Chim. Acta.* 1975; 58:590. (b) Oppolzer W, Flaskamp E, Bieber LW. *Helv. Chim. Acta.* 2001; 84:141.
10. For recent reviews for intramolecular Diels-Alder reactions for natural product synthesis, see: (a) Takao K, Munakata R, Tadano K. *Chem. Rev.* 2005; 105:4779. [PubMed: 16351062] . (b) Padwa A, Bur SK. *Tetrahedron.* 2007; 63:5341. [PubMed: 17940591]
11. For some examples of IMDA using systems related to 1-*N*-substituted-1,3-dienes over the last thirty years, see: (a) Stork G, Morgans DJ Jr. *J. Am. Chem. Soc.* 1979; 101:7110.. (b) Witiak DT,

- Tomita K, Patch RJ, Enna SJ. *J. Med. Chem.* 1981; 24:788. [PubMed: 6268788] . (c) Keck GE, Boden E, Sonnewald U. *Tetrahedron Lett.* 1981; 22:2615.. (d) Martin SF, Tu C, Kimura M, Simonsen SH. *J. Org. Chem.* 1982; 47:3634.. (e) Hwang G, Magnus P. *J. Chem. Soc. Chem. Commun.* 1983; 693. (f) Hayakawa K, Yasukouchi T, Kanematsu K. *Tetrahedron Lett.* 1986; 27:1837.. (g) Boeckman RK Jr. Goldstein SW, Walters MA. *J. Am. Chem. Soc.* 1988; 110:8250.. (h) Martin SF, Li W. *J. Org. Chem.* 1989; 54:265.. (i) Boger DL, Zhang M. *J. Am. Chem. Soc.* 1991; 113:4230.. (j) Yamada H, Sodeoka M, Shibasaki M. *J. Org. Chem.* 1991; 56:4569.. (k) Rawal VH, Iwasa S. *J. Org. Chem.* 1994; 59:2685.. (l) Keck GE, McHardy SF, Murry JA. *J. Am. Chem. Soc.* 1995; 117:7289.. (m) Lee M, Ikeda I, Kawabe T, Mori S, Kanematsu K. *J. Org. Chem.* 1996; 61:3406.. (n) Padwa A, Brodney MA, Dimitroff M. *J. Org. Chem.* 1998; 63:5304.. (o) Boonsombat J, Zhang H, Chughtai MJ, Hartung J, Padwa A. *J. Org. Chem.* 2008; 73:3539. [PubMed: 18376864] . (p) Friedrich A, Jainta M, Nising CF, Bräse S. *Synlett.* 2008:589.
12. Hayashi R, Hsung RP, Feltenberger JB, Lohse AG. *Org. Lett.* 2009; 11:2125. [PubMed: 19371081]
13. For a few notable examples of allenamide isomerizations, see: (a) Overman LE, Clizbe LA, Freerks RL, Marlowe CK. *J. Am. Chem. Soc.* 1981; 103:2807.. Also see. (b) Farmer ML, Billups WE, Greenlee RB, Kurtz AN. *J. Org. Chem.* 1966; 31:2885.. (c) Kinderman SS, van Maarseveen JH, Schoemaker HE, Hiemstra H, Rutjes FPJT. *Org. Lett.* 2001; 3:2045. [PubMed: 11418045] . (d) Trost BM, Stiles DT. *Org. Lett.* 2005; 7:2117. [PubMed: 15901148]
14. For a leading review on allenamide chemistry, see: Hsung RP, Wei L-L, Xiong H. *Acc. Chem. Res.* 2003; 36:773. [PubMed: 14567711]
15. Also see: (a) Lohse AG, Hsung RP. *Chem. A Eur. J.* 2011; 17:3812.. (b) Standen PE, Kimber MC. *Curr. Opin. Drug Discov. & Devel.* 2010; 13:645.. (c) Deagostino A, Prandi C, Tabasso S, Venturello P. *Molecules.* 2010; 15:2667. [PubMed: 20428072] . For general reviews on allenes, see: (d) Krause N, Hashmi ASK. *Modern Allene Chemistry.* 2004; Vol 1 and 2 Wiley-VCH Verlag GmbH & Co. KGaA Weinheim
16. Given the large volume of recent activities in allenamide chemistry, for reports in 2010–2011, see: (a) Lohse AG, Hsung RP, Leider MD, Ghosh SK. *J. Org. Chem.* 2011; 76 ASAP. . (b) Faustino H, López F, Castedo L, Mascareñas JL. *Chem. Sci.* 2011 ASAP. . (c) Zhu Y, Yin G, Hong D, Lu P, Wang Y. *Org. Lett.* 2011; 13:1024. [PubMed: 21268594] . (d) Yin G, Zhu Y, Zhang L, Lu P, Wang Y. *Org. Lett.* 2011; 13:940. [PubMed: 21348533] . (e) Lohse AG, Krenske EH, Antoline JE, Houk KN, Hsung RP. *Org. Lett.* 2010; 12:5506. [PubMed: 21049917] . (f) Hayashi R, Walton MC, Hsung RP, Schwab JH, Yu X. *Org. Lett.* 2010; 12:5768. [PubMed: 21090590] . (g) Hayashi R, Feltenberger JB, Hsung RP. *Org. Lett.* 2010; 12:1152. [PubMed: 20170149] . (h) Beccalli EM, Bernasconi A, Borsini E, Broggin G, Rigamonti M, Zecchi G. *J. Org. Chem.* 2010; 75:6923. [PubMed: 20863085] . (i) Hill AW, Elsegood MRJ, Kimber MC. *J. Org. Chem.* 2010; 75:5406. [PubMed: 20670040] . (j) Persson AKÅ, Bäckvall J-E. *Angew. Chem. Int. Ed.* 2010; 49:4624.. (k) Krenske EK, Houk KN, Lohse AG, Antoline JE, Hsung RP. *Chem. Science.* 2010; 1:387.. (l) Danowitz AM, Taylor CE, Shrikian TM, Mapp AK. *Org. Lett.* 2010; 12:2574. [PubMed: 20438110] . (m) Zbieg JR, E. McInturff EL, Krische MJ. *Org. Lett.* 2010; 12:2514. [PubMed: 20459077] . (n) Cordier P, Aubert C, Malacria M, Gandon V, Lacôte E. *Chem. Eur. J.* 2010; 16:9973.. (o) Kimber MC. *Org. Lett.* 2010; 12:1128. [PubMed: 20143846] . (p) Hashimoto K, Horino Y, Kuroda S. *Heterocycles.* 2010; 80:187.
17. See Supporting Information
18. (a) Trost BM, Stiles DT. *Org. Lett.* 2005; 7:2117. [PubMed: 15901148] (b) Shen L, Hsung RP, Zhang Y, Antoline JE, Zhang X. *Org. Lett.* 2005; 7:3081. [PubMed: 15987210] (c) Persson AKÅ, Johnston EV, Bäckvall J-E. *Organic Lett.* 2009; 11:3814.

Figure 1.
X-Ray Structure of **16**.

Figure 2. Tandem γ -Isomerization–Oppolzer-Type IMDA.^a

a. All reactions were run for 40–48 h in a screw-cap vial with Teflon coated caps at temp = 165–175 °C using the xylene/decane mixed solvent system, and 3.0 equiv Et₃N as additive. Cycloaddition for preparing **23** was run at 210 °C. The diastereomeric ratios were determined using ¹H and/or ¹³C NMR.

Scheme 1.
Oppolzer's IMDA Via Allenamide Isomerization.

Scheme 2.
Facile Assembly of *N*-Tethered Allenamides.

Scheme 3.
Sequential Pursuit Versus Tandem Process.

Scheme 4.
An Unexpected Enamide 1,4-Addition.

Scheme 5.
Assembly of Indolines.

Scheme 6.
Rapid Constructions of Quinolines.

Table 1

Tandem γ -Isomerization–Oppolzer-Type IMDA.

entry	W =	solvent ^a	additive [equiv]	temp [°C]	time [h]	yield [%] ^b
1	11 : Ts	toluene	-	135	18	trace ^c
2	11 : Ts	toluene	-	170	26	13 ^d
3	12 : <i>p</i> -Ns	xylene	-	150	40	42
4	12 : <i>p</i> -Ns	xylene	-	180	19	33
5	12 : <i>p</i> -Ns	decane	-	180	15	25
6	12 : <i>p</i> -Ns	1,2-DiClPh	-	165	24	33
7	12 : <i>p</i> -Ns	Ph ₂ O	-	165	24	32
8	12 : <i>p</i> -Ns	xy]/decane	-	165	24	38
9	11 : Ts	toluene	BSA [0.3–5]	165	24	- ^e
10	12 : <i>p</i> -Ns	xy]/decane	BSA [3]	165	24	- ^e
11	12 : <i>p</i> -Ns	xy]/decane	proton sponge [3]	165	24	77
12	12 : <i>p</i> -Ns	xy]/decane	1-Me-imidazole [3]	165	24	63
13	11 : Ts	xy]/decane	Et ₃ N [3]	165	24	34
14	12 : <i>p</i> -Ns	xy]/decane	Et ₃ N [3]	165	24	76
15	12 : <i>p</i> -Ns	xy]/decane	DBU [3]	165	24	trace ^f
16	12 : <i>p</i> -Ns	xy]/decane	DABCO [3]	165	24	trace ^f
17	12 : <i>p</i> -Ns	xy]/decane	Cs ₂ CO ₃ [3]	165	24	48
18	12 : <i>p</i> -Ns	xy]/decane	K ₂ CO ₃ [3]	165	24	- ^g
19	12 : <i>p</i> -Ns	xy]/decane	4Å MS [3]	165	24	- ^g
20	12 : <i>p</i> -Ns	xy]/decane	Na ₂ SO ₄ [3]	165	24	- ^g

^a All reactions were run in a screw-cap vial with Teflon coated caps. Concn = 0.05 M except for entries 1–3, and 10: concn = 0.02 M.^b Isolated yields for **15** or **16**.^c The only product was *N*-tethered 1-amidodiene **13** in 78% yield.^d The respective secondary amide was isolated in 15% yield as a result of hydrolysis along with 26% of **13**.

^e A complex mixture.

^f Mostly was decomposition of **12**.