The Child's Rumbling Slumber

David Hiestand, M.D., Ph.D.¹; Barbara Phillips, M.D., M.S.P.H.²¹Department of Medicine, Division of Pulmonary, Critical Care, & Sleep Disorders Medicine, University of Louisville, Louisville, KY;²Department of Medicine, Division of Pulmonary, Critical Care, & Sleep Disorders Medicine, University of Kentucky College of Medicine, Lexington, KY

A 4-year-old Caucasian male presents to a sleep disorders clinic for evaluation. The child has recently started preschool, and the parents report that he has significant difficulty with daytime activities, such as following directions and completing assigned tasks. In addition, he has begun to snore almost nightly in the last 6 months. His mother notes that he sweats profusely while sleeping and his sleep is described as restless ("flops all over" the bed and sleeps in odd body positions). He goes to bed at 8 pm and rises at 7 am. The bedtime ritual is fairly easy, and he does not commonly resist sleep.

His daytime activity is characterized as "more active than other children." He still takes one afternoon nap of approximately 2 hours. Despite the nap he also commonly falls asleep on the 30-minute ride home from daycare.

The birth history is unremarkable: born at 40 weeks, without complications, and normal early growth and development. He takes occasional antihistamines for cough and rhinitis.

The patient has no siblings. His father has obstructive sleep apnea. There are no family members with other cardiovascular, respiratory, or neurologic disease. He is not exposed to passive smoke. He lives with both parents and tends to sleep in his own bed.

A review of systems is positive for intermittent upper airway allergy symptoms. The remainder of the review of systems is unremarkable. He does have occasional episodes of bed wetting, but these are decreasing over time.

On examination the child is active but follows direction. Height and weight are at the 25th percentile for age. Vital signs are normal for age. Nares demonstrate modest nasal edema. The oropharyngeal exam demonstrates a Mallampati class II oropharynx, with tonsils graded at 3+. The palate and jaw are normal in size and dimension. The tongue is normal in size and position. The remainder of the examination is unremarkable.

QUESTION:

Which of the following would be most helpful in the evaluation of OSA in this 4-year-old?

- a. Nighttime home sleep study
- b. Daytime (nap) in-lab PSG
- c. Nighttime in-lab PSG
- d. Nighttime home video recording of child sleeping
- e. No test is needed. History is adequate to diagnose disorder

ANSWER: c. Nighttime in-lab PSG

Obstructive sleep apnea occurs in approximately 2% of young children with a peak prevalence at about age 5,¹ while habitual snoring occurs in 8% to 12%. The clinical consequences of childhood obstructive sleep apnea have been recently reviewed and include behavior problems, academic performance deficits, growth limitations, and cardiovascular consequences.²

Differentiating OSA from habitual snoring can be difficult based on clinical history. Overnight polysomnography provides the most information on extent of cardiorespiratory events and is currently the gold standard for evaluation of pediatric obstructive sleep apnea. In-lab polysomnography allows for the characterization of sleep architecture and assures adequate evaluation of sleep stages, including REM sleep. Currently, both the American Academy of Pediatrics and American Thoracic Society recommend overnight polysomnography for evaluation of children with signs and symptoms of OSA.^{3,4}

Potential limitations to overnight in-lab studies include night-to-night variability, first night effect with disrupted sleep architecture, and patient/family apprehension about the study. Older children do demonstrate some difference in sleep architecture when studied on consecutive nights; however, respiratory differences are not clinically significant.^{5,6}

Nighttime home sleep studies are quite varied in scope of and content. The American Academy of Pediatrics listed home sleep studies as an option for diagnostic testing.³ While there have been some studies showing reasonable positive predictive value,⁷ many insurance companies consider home studies investigational and do not reimburse for these. Furthermore, children tend to pull off leads, making data difficult to interpret. Home testing based on oximetry alone is not adequate for identification of OSA in otherwise healthy children.⁸ The American Academy of Sleep Medicine's clinical guidelines on portable monitoring specifically exclude children due to paucity of data.⁹

Nap studies are inadequate since REM sleep is limited or may not be achieved during sleep, thus underestimating the degree and severity of respiratory events.^{10,11}

Nighttime home video recording is listed as an option for evaluation of certain children in the AAP consensus statement.³ In a trial of 58 children comparing PSG and home video recordings, the sensitivity of overall investigator judgment was 94% with a specificity of 68%, with the authors suggesting a possible role in screening for those needing formal PSG.¹² This method of evaluation is not, however, reimbursed by most insurance companies and is not commonly used as an isolated testing method.

Clinical history has been shown to be inadequate for diagnosis of OSA in children. A systematic review of the literature found very little correlation between clinical exam and PSG-proven OSA.¹³ Common nighttime symptoms include snoring; labored breathing with gasping, gagging, or choking; noisy breathing; witnessed apnea; and sleeping with open mouth and hyperextended neck. Sweating during sleep is also a commonly noted symptom but has very little correlation with OSA.

Norms for polysomnography in children have been established by 4 studies.¹⁴⁻¹⁷ These studies demonstrated that normal children have central and obstructive apnea indices of one or fewer per hour. Defining the lower limit of clinically meaning-

ful sleep apnea, however, remains the subject of some debate, and most clinicians would favor more liberal standards than an apnea hypopnea index of 5/h as the lower limit of disease. One clinical severity index defines mild OSA as an AHI $\geq$ 1-5 without significant desaturations, moderate OSA as AHI $\geq$ 5-10 without significant desaturation, and severe OSA as AHI $\geq$ 10 or significant, sustained desaturations below 92%. The significance of such severity scoring as it relates to treatment options or long-term sequelae is uncertain.

REFERENCES

1. Redline S, Tishler PV, Schluchter M, Aylor J, Clark K, Graham G. Risk factors for sleep-disordered breathing in children. Associations with obesity, race, and respiratory problems. *Am J Respir Crit Care Med* 1999;159:1527-32.
2. Halblower AC, Ishman SL, McGinley, BM. Childhood obstructive sleep-disordered breathing: a clinical update and discussion of technological innovations and challenges. *Chest* 2007;132:2030-41.
3. American Academy of Pediatrics. Clinical practice guideline: diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 2002;109:703-22.
4. American Thoracic Society. Standards and indications for cardiopulmonary sleep studies in children. *Am J Respir Crit Care Med* 1996;153:866-78.
5. Li AM, Wing YK, Cheung A, et al. Is a 2-night polysomnographic study necessary in childhood sleep-related disordered breathing? *Chest* 2004;126:1467-72.
6. Scholle S, Scholle HC, Kemper A, et al. First night effect in children and adolescents undergoing polysomnography for sleep-disordered breathing. *Clin Neurophysiol* 2003;114:2138-45.
7. Jacob SV, Morielli A, Mograss MA, Ducharme FM, Schloss MD, Brouillette RT. Home testing for pediatric obstructive sleep apnea syndrome secondary to adenotonsillar hypertrophy. *Pediatr Pulmonol* 1995;20:241-52.
8. Kirk VG, Bohn SG, Flemons WW, Remmers JE. Comparison of home oximetry monitoring with laboratory polysomnography in children. *Chest* 2003;124:1702-8.
9. Collop NA, Anderson WM, Boehlecke B, et al. Clinical guidelines for the use of unattended portable monitors in the diagnosis of obstructive sleep apnea in adult patients. Portable Monitoring Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2007;3:737-47.
10. Marcus CL, Keens TG, Ward SL. Comparison of nap and overnight polysomnography in children. *Pediatr Pulmonol* 1992;13:16-21.
11. Saeed MM, Keens TG, Stabile MW, Bolokowicz J, Davidson Ward SL. Should children with suspected obstructive sleep apnea syndrome and normal nap sleep studies have overnight sleep studies. *Chest* 2000;118:360-5.
12. Sivan Y, Kornecki A, Schonfeld T. Screening obstructive sleep apnoea syndrome by home videotape recording in children. *Eur Respir J* 1996;9:2127-31.
13. Brietzke SE, Katz ES, Roberson DW. Can history and physical examination reliably diagnose pediatric obstructive sleep apnea/hypopnea syndrome? A systematic review of the literature. *Otolaryngol Head Neck Surg* 2004;131:827-32.
14. Uliel S, Tauman R, Greenfeld M, Sivan Y. Normal polysomnographic respiratory values in children and adolescents. *Chest* 2004;125:872-8.
15. Traeger N, Schultz B, Pollock AN, Mason T, Marcus C, Arens R. Polysomnography values in children 2-9 years old: Additional data and review of the literature. *Pediatr Pulmonol* 2005;40:22-30.
16. Montgomery-Downs HE, O'Brien LM, Gulliver TE, Gozal D. Polysomnographic characteristics in normal preschool and early school-aged children. *Pediatrics* 2006;117:741-53.
17. Marcus CL, Omlin KJ, Basinki DJ, et al. Normal polysomnographic values for children and adolescents. *Am Rev Respir Dis* 1992;146:1235-9.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication March, 2011

Accepted for publication March, 2011

Address correspondence to: David Hiestand, M.D., Ph.D., University of Louisville School of Medicine, Division of Pulmonary, Critical Care, and Sleep Disorders Medicine, 550 S. Jackson St, ACB, A3R40, Louisville, KY 40202; Tel: (502) 852-5841; Fax: (502) 852-1359; E-mail: david.hiestand@louisville.edu

DISCLOSURE STATEMENT

The authors have indicated no financial conflicts of interest.