Published in final edited form as:

Chemistry. 2011 May 27; 17(23): 6326–6329. doi:10.1002/chem.201100444.

Nickel-Catalyzed Cyclizations of Enoates and Chiral Allenes: An Approach to Domoic Acid

Ahmad S. ElDouhaibi, Refaie M. Kassab, Minsoo Song, and John Montgomery

Department of Chemistry University of Michigan 930 N. University Ave., Ann Arbor, MI 48109-1055

Keywords

allenes; alkynes; C-C coupling; nickel; cyclization

The kainoid family of marine natural products has attracted considerable interest due to the potent pharmacological properties of compounds in this class.^[1] By serving as conformationally restricted analogues of glutamate, this group of natural products displays potent neuroexcitatory activity, and the more complex members of the domoic acid type have been responsible for outbreaks of poisoning from mussel toxins.^[2] The kainic acid and domoic acid families of natural products include a diverse range of structures that possess highly functionalized pyrrolidine di- or triacid motifs (Scheme 1).^[3] Kainic and allokainic acids possess a simple C4 propenyl substituent, whereas domoic acid and its isomers possess more complex C4 functionality. Our interest in the synthesis of members of this class of compounds has led to the development of a number of nickel-catalyzed processes, including the alkylative cyclization of enoate-alkyne and enoate-allene substrates. These processes have served as key steps in the total syntheses of kainic acid (**1**), allokainic acid (**2**), and isodomoic acids G (**3**) and H (**4**).^[4–6] We envisioned that the development of a related cyclization method involving cyclization of an enoate tethered with a chiral allene would allow synthesis of the remaining members of the domoic acid class, including domoic acid (**5**), that do not possess the C4 exocyclic functionality seen in isodomoic acids G and H. Herein, we describe an initial investigation of this strategy, and the synthesis of a model substructure of domoic acid.

Our first-generation strategy for the synthesis of domoic acid (**5**) features an ambitious dimethyl zinc-mediated cyclization of enoate-allene **6**.^[7] As depicted (Scheme 2), we anticipated that the allene chirality of **6** would ultimately establish stereochemistry in the generation of the C3 and C4 stereocenters and the C1' trisubstituted alkene of **9** during the cyclization.^[8] Our mechanistic hypothesis involves formation of a Ni(0) π -complex **7** with an eclipsed orientation of the reactive π -systems, followed by the formation of metallacycle **8**, where all of the key stereochemical features are established.^[9] The prediction that the proximal π -system of the allene will undergo coupling draws from our earlier observations made in the synthesis of kainic acid,^[4b] as well as related five-membered aldehyde-allene alkylative cyclization processes.^[7d]

For our initial examination of this approach, the requisite substrate **12** was first prepared (for ease of synthesis) as a mixture of allene epimers beginning from alkyne **10** (Scheme 3). Our

Correspondence to: John Montgomery.

Fax: (+) 734-763-1106 jmontg@mich.edu.

Supporting information for this article is available on the WWW under <http://www.chemeurj.org/> or from the author.

earlier studies had illustrated that unsaturated acyl oxazolidinones were the most effective enoate derivatives in nickel cyclization processes.^[4] Upon conversion of alcohol **11** to enoate **12**, one of the diastereomers of **12** was found to be unstable, based on facility of an undesired Diels Alder cycloaddition to afford product **13b** along with a single diastereomer of the desired enoate **12a**. The facility of this cycloaddition is notable given the comparatively harsh conditions required for other types of vinyl allene Diels Alder reactions.^[10] Based on the stereochemical outcome of the generation of product **13b**, allene **12b** can be assumed to be the unstable allene diastereomer that undergoes rapid cycloaddition to **13b**, and allene **12a** is therefore the diastereomer that is isolable.

As structure **12a** is the diastereomer anticipated to be required for the nickel-catalyzed cyclization, its participation in an alkylative cyclization reaction was examined (Scheme 4). However, we were unable to access the desired product **14**. Instead, Diels Alder cycloaddition predominated to afford structure **13a**, providing further proof of the structure of diastereomer **12a**. It is unclear if the cycloaddition is catalyzed by Ni(COD)₂ or ZnMe₂, and due to the limited material available, this question was not explored.^[11,12] Instead, we opted to redesign the strategy by removing the C3'-C4' unsaturation, thus removing the complexity of the competing Diels Alder cycloaddition.

In order to pursue this second-generation strategy, racemic allenyl mesylate **17** was prepared by a straightforward route from alkyne **15** and aldehyde **16** (Scheme 5). *N*-Alkylation of **18** with mesylate **17** followed by ester reduction afforded **19** as a mixture of allene epimers. Oxidation state adjustment and olefination of **19** provided diastereomers **20a** and **20b**, which were separated by preparative HPLC. This route was attractive in that both diastereomers of **20** were made available for investigation, and the stereochemistry of **20a** was later confirmed by independent asymmetric synthesis (*vide infra*).

With allene epimers **20a** and **20b** both in hand, nickel-catalyzed dimethyl zinc-promoted cyclization was investigated (Scheme 6). Gratifyingly, as anticipated from our strategic plan, cyclization of **20a** cleanly afforded desired compound **23a**, whereas cyclization of **20b** under identical conditions afforded diastereomer **23b** in a less-efficient and lower-yielding cyclization. The π -complex **21a**, followed by cyclization to metallacycle **22a**, likely explains the stereochemical outcome of this key cyclization process in the formation of **23a**.^[9] Alternatively, by utilizing the opposite allene chirality, modified orientation of substrate **20b** in π -complex **21b**, followed by cyclization to metallacycle **22b**, provides diastereomer **23b**. This outcome illustrates that the allene chirality is essential in establishing the desired C3 and C4 stereochemical arrangement. Oxazolidinones **23a** and **23b** were cleanly converted to methyl esters **24a** and **24b** in straightforward fashion.

Lastly, an improved route to compound **24a** involving asymmetric synthesis of the requisite allene fragment and an improved entry to a pyrrolidine precursor was developed (Scheme 7). Enantioselective addition of alkyne **15** to aldehyde **16** following the method of Carreira provided alcohol **25a** in 94% ee,^[13] and subjecting this structure to the allylic diazene method of Myers provided allene **26a** in 92% ee.^[14] Conversion to mesylate **17a** and *N*-alkylation of **27** provided access to diastereomerically pure **28a**. Conversion of **28a** to **19a** and then **20a** completed the stereoselective approach to **24a** and confirmed the structures of diastereomers **20a** and **20b**. In this route, precursor **27** was utilized rather than *N*-alkylation of **18** in order to avoid a problematic epimerization that had been observed in our parallel efforts to synthesize isodomoic acids G and H. This improved entry to compound **24a** avoids the tedious separation of **20a** and **20b** and provides a selective entry to the substructure that directly corresponds to that seen in domoic acid and a range of naturally occurring isodomoic acid structures.

In summary, the first examples of enoate-chiral allene alkylative cyclizations have been demonstrated. The process is effective in the synthesis of functionalized pyrrolidine derivatives that possess stereodefined alkenyl substituents. An advanced intermediate towards the synthesis of domoic acid was prepared using this method. The efficient synthesis of structure **24a** may be useful in future synthetic approaches to domoic acid and several naturally occurring isodomoic acids. A notable feature of this approach is that the C3-C4 bond, the C3 and C4 stereogenic centers, and the C1'-C2' trisubstituted alkene are introduced efficiently and with high selectivity in this new catalytic process.

Experimental Section

Preparation of compound 23a

To a 0.5 mL 0 °C THF solution of anhydrous ZnCl² (68 mg, 0.5 mmol), 0.67 mL of MeLi (0.66 mL, 1.0 mmol, 1.5 M in diethyl ether) was added dropwise. After stirring for 10 min at 0 °C, the mixture was transferred by cannula to a 5 mL 0 °C THF solution of Ni(COD)₂ (12 mg, 0.04 mmol). The mixture was cooled to -20 °C, then an 8 mL THF solution of **20a** (100 mg, 0.2 mmol) and titanium isopropoxide (60 mg, 60 µL, 0.2 mmol) was added dropwise. The mixture was slowly warmed to rt, and stirred for 2 h, quenched with sat. NH₄Cl, extracted with ethyl acetate, and the combined extracts were washed with brine and dried over MgSO₄. After filtration, the solvent was removed under vacuum, and the crude product was purified by flash column chromatography (hexane/ethyl acetate, 2:1 to 1:2) to yield 78 mg (77%, 94:6 *dr*) of **23a** as a thick oil.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to acknowledge receipt of NIH grant GM57014 in support of this work. The authors thank colleagues at Wayne State University, where early portions of this work were conducted. JM thanks the faculty and students of the Institute of Chemical Research of Catalonia (ICIQ) for hospitality during a sabbatical stay.

References

1. a Monaghan DT, Bridges RJ, Cotman CW. *Annu. Rev. Pharmacol. Toxicol.* 1989; 29:365–402. [PubMed: 2543272] b Moloney MG. *Nat. Prod. Rep.* 1998; 15:205–219. [PubMed: 9586226] c Chamberlin, R.; Bridges, R. *Drug Design For Neuroscience*. Kozikowski, AP., editor. Raven Press, Ltd.; New York: 1993. p. 231-259. d McGeer, EG.; Olney, JW.; McGeer, PL. *Kainic Acid as a Tool in Neurobiology*. Raven Press; New York: 1978.
2. a Lefebvre KA, Robertson A. *Toxicon.* 2010; 56:218–230. [PubMed: 19505488] b Wright JLC, et al. *Can. J. Chem. Rev. Can. Chim.* 1989; 67:481–490.
3. a Parsons AF. *Tetrahedron.* 1996; 52:4149–4174. b Moloney MG. *Nat. Prod. Rep.* 1999; 16:485–498. [PubMed: 10467739] c Moloney MG. *Nat. Prod. Rep.* 2002; 19:597–616. [PubMed: 12430725] d Takemoto T, Daigo K. *Chem. Pharm. Bull.* 1958; 6:578.
4. a Chevliakov MV, Montgomery J. *Angew. Chem. Int. Ed.* 1998; 37:3144–3146. b Chevliakov MV, Montgomery J. *J. Am. Chem. Soc.* 1999; 121:11139–11143. c Ni Y, Amarasinghe KKD, Ksebati B, Montgomery J. *Org. Lett.* 2003; 5:3771–3773. [PubMed: 14507227] d Ni Y, Kassab RM, Chevliakov MV, Montgomery J. *J. Am. Chem. Soc.* 2009; 131:17714–17718. [PubMed: 19899794]
5. For other approaches to kainic and allo-kainic acid, see reference 3 and the following representative examples: a Oppolzer W, Thirring K. *J. Am. Chem. Soc.* 1982; 104:4978–4979. b Sakaguchi H, Tokuyama H, Fukuyama T. *Org. Lett.* 2008; 10:1711–1714. [PubMed: 18402455] c Xia Q, Ganem B. *Org. Lett.* 2001; 3:485–487. [PubMed: 11434316] d Martinez MM, Hoppe D. *Org. Lett.* 2004;

- 6:3743–3746. [PubMed: 15469338] e Yoo SE, Lee SH. *J. Org. Chem.* 1994; 59:6968–6972. f Farwick A, Helmchen G. *Org. Lett.* 2010; 12:1108–1111. [PubMed: 20148523]
6. For other approaches to domoic acid and naturally occurring isodomoic acids: a Ohfuné Y, Tomita M. *J. Am. Chem. Soc.* 1982; 104:3511–3513. b Clayden J, Knowles FE, Baldwin IR. *J. Am. Chem. Soc.* 2005; 127:2412–2413. [PubMed: 15724994] c Denmark SE, Liu JHC, Muhuhi JM. *J. Am. Chem. Soc.* 2009; 131:14188–14189. [PubMed: 19807173]
7. For development of related cyclization methods: a Montgomery J, Savchenko AV. *J. Am. Chem. Soc.* 1996; 118:2099–2100. b Montgomery J, Chevliakov MV, Brielmann HL. *Tetrahedron.* 1997; 53:16449–16462. c Montgomery J, Oblinger E, Savchenko AV. *J. Am. Chem. Soc.* 1997; 119:4911–4920. d Song M, Montgomery J. *Tetrahedron.* 2005; 61:11440–11448. e Ikeda S, Sato Y. *J. Am. Chem. Soc.* 1994; 116:5975–5976. f Ikeda S, Yamamoto H, Kondo K, Sato Y. *Organometallics.* 1995; 14:5015–5016. g Kang SK, Yoon SK. *Chem. Commun.* 2002:2634–2635.
8. For representative transition metal-catalyzed couplings involving chiral allenes: a Brummond KM, Kerekes AD, Wan HH. *J. Org. Chem.* 2002; 67:5156–5163. [PubMed: 12126400] b Ng SS, Jamison TF. *J. Am. Chem. Soc.* 2005; 127:7320–7321. [PubMed: 15898774] c Urabe H, Takeda T, Hideura D, Sato F. *J. Am. Chem. Soc.* 1997; 119:11295–11305. d Wender PA, Glorius F, Husfeld CO, Langkopf E, Love JA. *J. Am. Chem. Soc.* 1999; 121:5348–5349.
9. While this simplified model serves as a useful stereochemical predictor, computational studies suggested a more complex role of the organozinc in promoting the cyclization. See: Hratchian HP, Chowdhury SK, Gutierrez-Garcia VM, Amarasinghe KKD, Heeg MJ, Schlegel HB, Montgomery J. *Organometallics.* 2004; 23:4636–4646.
10. a Deutsch EA, Snider BB. *J. Org. Chem.* 1982; 47:2682–2684. b Keck GE, Kachensky DF. *J. Org. Chem.* 1986; 51:2487–2493. For facile ene reaction – Diels Alder cascades involving vinyl allenes: c Robinson JM, Sakai T, Okano K, Kitawaki T, Danheiser RL. *J. Am. Chem. Soc.* 2010; 132:11039–11041. [PubMed: 20698669] d Sakai T, Danheiser RL. *J. Am. Chem. Soc.* 2010; 132:13203–13205. [PubMed: 20815385] For a review on vinyl allene cycloadditions: e Murakami M, Matsuda T, Krause N, Hashmi ASK. *Cycloadditions of Allenes. Modern Allene Chemistry.* 2004; 2:727–815. Wiley-VCH Weinheim For an overview of cyclizations and cycloadditions of allenes: f Wei LL, Xiong H, Hsung RP. *Acc. Chem. Res.* 2003; 36:773–782. [PubMed: 14567711]
11. For nickel-catalyzed [4+2] reactions: a Wender PA, Smith TE. *J. Org. Chem.* 1995; 60:2962–2963. b Wender PA, Smith TE. *J. Org. Chem.* 1996; 61:824–825.
12. For other metal-catalyzed processes of vinyl allenes: a Murakami M, Itami K, Ito Y. *J. Am. Chem. Soc.* 1999; 121:4130–4135. b Ma SM. *Chem. Rev.* 2005; 105:2829–2871. [PubMed: 16011326] c Hashmi ASK. *Angew. Chem. Int. Ed.* 2000; 39:3590–3593.
13. El-Sayed E, Anand NK, Carreira EM. *Org. Lett.* 2001; 3:3017–3020. [PubMed: 11554832]
14. Myers AG, Zheng B. *J. Am. Chem. Soc.* 1996; 118:4492–4493.

Scheme 1.
Representative kainoid amino acids.

Scheme 2.
First-generation strategy.

Scheme 3.

Examination of vinyl allene cyclizations. (a) *i.* **10**, *n*BuLi, then crotonaldehyde, THF, -50 °C, 65 %, *ii.* Bu₃SnH, AIBN, 90 °C, then MsCl, Et₃N, CH₂Cl₂, then (*n*Bu)₄NF, THF, 45 %; (b) oxalyl chloride, DMSO, Et₃N, CH₂Cl₂, -78 to -20 °C, then ((2-oxooxazolidin-3-yl)methyl)triphenylphosphonium bromide, DMAP, -20 °C to rt, 14 % of **12a**, 31 % of **13b**.

Scheme 4.
Attempted vinyl allene alkylative cyclization.

Scheme 5.

Preparation of cyclization substrates. (a) *i.* **15**, *n*BuLi, THF, -78 °C, then **16**, 65 %, *ii.* PPh₃, DEAD, *o*-nitrobenzene sulfonyl hydrazine, THF, 15 °C to rt, 61 %, *iii.* PPTS, EtOH, 55 °C, 85 %, *iv.* MsCl, Et₃N, CH₂Cl₂, 0 °C, 98 %; (b) *i.* KHMDS, **17**, THF, 0 °C, 41 %, *ii.* NaBH₄, EtOH, 0 °C, 80 %; (c) oxalyl chloride, DMSO, Et₃N, CH₂Cl₂, -78 to -20 °C, then ((2-oxooxazolidin-3-yl)methyl)triphenylphosphonium bromide, DMAP, -20 °C to rt, 69 % combined of **20a** and **20b** (1:1, separated by HPLC).

Scheme 6.

Alkylative cyclization of an enoate / chiral allene. (a) ZnCl_2 , MeLi , $\text{Ti}(\text{O}-i\text{Pr})_4$, $\text{Ni}(\text{COD})_2$ (20 mol %), -20°C to rt, 77 % (in the case of **23a**), 24 % (in the case of **23b**); (b) MeOMgBr , toluene/THF, 0°C to rt, 75 % (in the case of **24a**), 79 % (in the case of **24b**).

Scheme 7.

Enantioselective preparation of a cyclization precursor. (a) zinc trifluoromethane sulfonate, . (1*R*, 2*S*)-(-)-*N*-methylephedrine, **15**, then **16**, toluene, rt, 58 % (94 % ee); (b) PPh₃, DEAD, *o*-nitrobenzene sulfonyl hydrazine, THF, -15 °C to rt, 61 % (92 % ee); (c) *i*. PPTS, EtOH, 55 °C, 85 %, *ii*. MsCl, Et₃N, CH₂Cl₂, 0 °C, 98 %; (d) **17a**, **27**, benzyltriethylammonium bromide, 50 % NaOH, CH₂Cl₂, 50 %, (e) PPTS, EtOH, 55 °C, 76 %. (f) oxalyl chloride, DMSO, Et₃N, CH₂Cl₂, -78 to -20 °C, then ((2-oxooxazolidin-3-yl)methyl)triphenylphosphonium bromide, DMAP, -20 °C to rt, 74 %.