Rare disease

Ovarian metastasis in a transposed ovary 10 years after primary cervical cancer: the importance of histologic examination and review of literature

Julienne A Janse,¹ Daisy MDS Sie-Go,² Henk WR Schreuder¹

¹Division of Woman and Baby, Department of Gynaecologic Oncology, University Medical Centre, Utrecht, The Netherlands;

²Department of Pathology, University Medical Centre Utrecht, Utrecht, Netherlands

Correspondence to Dr Henk WR Schreuder, h.w.r.schreuder@umcutrecht.nl

Summary

Cases of cervical carcinoma metastasising to the transposed ovary are rarely reported in the literature. In this report, the authors present the case of a 53-year-old woman with a persisting, unsuspected cyst in the right transposed ovary, 10 years after treatment for adenosquamous carcinoma of the cervix. It is the first report describing a secondary ovarian malignancy originating from a cervical adenosquamous carcinoma in a transposed ovary. In addition, this is the first account of an ovarian metastasis 10 years after primary treatment for cervical cancer. Furthermore, pathologic examination with immunohistochemistry and human papillomavirus genotyping played a key role in the diagnostic process, as the case did not raise suspicion by ultrasound findings neither by cytological examination after cytological aspiration or by appearance during surgery.

BACKGROUND

Secondary malignancies can be hard to trace back to their original site, and histopathologic examination often leads to the final diagnosis. Ovarian metastases of cervical cancers are uncommon. The incidence is reported to vary between 0.9% and 2.2%.^{1–3} Approximately, 5% of all secondary ovarian malignancies originate from the cervix.⁴ In literature most cases described are diagnosed during primary radical surgery which included bilateral oophorectomy. To prevent the loss of ovarian function by radical surgery and postoperative radiotherapy in premenopausal women, the possibility exists to perform ovarian transposition. Until now, only four cases have been described of ovarian metastasis in a transposed ovary.^{5–7} This is the first case reported to authors' knowledge, of an ovarian metastasis 10 years after primary cervical cancer.

CASE PRESENTATION

A 43-year-old Caucasian female was treated in 1997 for a FIGO Stage Ib1 adenosquamous carcinoma of the cervix. Diagnosis was based on histopathological analysis after a vaginal hysterectomy for excessive bleeding in another institution. The surgical margins were free of tumour, but angio-invasion was present. The patient was sent to our university institution for completion of treatment. Surgical removal of parametrium, vaginal cuff and a bilateral pelvic lymphadenectomy was performed. The ovaries did not look suspicious of tumour involvement and the right ovary was transposed above the pelvic brim along the paracolic gutter. Histopathological analysis of the surgical specimen revealed no tumour. Thirty-six lymph nodes were removed and examined, no metastatic disease was found. No additional treatment was given. She was followed biannually for 5 years and after that yearly.

Ten years after primary treatment, she presented with a continuous pain in the right upper abdomen which was present since several months.

INVESTIGATIONS

On abdominal ultrasound a 6 cm large, unilocular, thin-walled and clear cyst, was seen in the transposed right ovary without free fluid. Ca-125 was measured (31 U/ml). There was no suspicion of malignancy (figure 1). Despite the use of medroxyprogesteron, the cyst persisted. An ultrasound-guided aspiration of the cyst was carried out and cytological analysis showed only scanty cell material and no malignant cells. Two weeks after aspiration the cyst recurred.

DIFFERENTIAL DIAGNOSIS

- ▶ Simple cyst
- ▶ Follicular cyst
- ▶ Serous cystadenoma
- ▶ Paraovarian cyst
- ▶ Serous borderline cyst.

TREATMENT

Because of recurrence of the cyst after aspiration, a laparoscopic removal of the transposed right ovary was performed. During an uncomplicated procedure the right ovary was removed, it contained a smooth and benign looking cyst which was 7 cm large. The small left ovary was lying retroperitoneal and was not visualised or removed during the operation.

HISTOLOGY

Macroscopically, the right ovary measuring 7.5 cm in greatest diameter, demonstrated a partly cystic tumour.

Figure 1 Ultrasound ovarian cyst in transposed ovary. (A) Measurements and aspect, (B) doppler.

Microscopically, the tumour showed mostly solid areas with squamous differentiation (figure 2). The atypical cells showed hyperchromatic polymorphic nuclei and a small amount of cytoplasm, which focally displayed intracytoplasmic mucin. There was a transition into gland-like structures also showing hyperchromatic nuclei and some intracytoplasmic mucin (figure 3). Focally there was a very small area where the atypical epithelial cells displayed a more sex cord-like arrangement. Immunohistochemical investigation of this ovarian tumour showed positivity with the following markers: CK AE1/3, CK7, CEA, CK19, partially with CK 5/6 and p63. The MIB1 showed a high proliferation index. The following immunohistochemical markers were negative on the tumour: CK14, CK20 and p53, confirming the diagnosis of an adenosquamous carcinoma.

The original tumour of the uterine cervix of 1997 was re-examined additionally and displayed the same histopathological results (figure 4). Microscopically, the original tumour stromal invasion of the uterine cervix was at least 7.5 mm with focal angio-invasive growth but without perineural invasion. There was a 5 mm stromal tumour-free margin.

Molecular biological examination was performed with the Linear Array human papillomavirus (HPV) Genotyping Test (Linear Array HPV Genotyping Kit: 03378179 190, Linear Array HPV Detection Kit: 208693; Roche, Basel Switzerland) which revealed HR-HPV type 18 in both tumours.

In summary, the histopathological investigation surprisingly revealed as a coincidental finding a small

Figure 2 Ovarian tumour showing a transition into gland-like structures (H&E, 250 $\times$).

Figure 3 Ovarian tumour showing a transition into gland-like structures. Hyperchromatic nuclei and some intracytoplasmic mucin are also present (arrow) (H&E, 400 $\times$).

adenosquamous carcinoma in the unsuspected right ovary. To reveal the origin of this uncommon finding molecular biological examination was performed on the ovary and the original cervical cancer from 1997. This displayed high-risk HPV type 18 in both the ovarian carcinoma and cervical primary tumour, confirming the diagnosis of a metastasis of cervical origin.

FOLLOW-UP

A PET-CT scan and a bone scintigraphy were performed, without evidence of a malignancy or further metastasis. After multidisciplinary discussion and consultation of the patient, the decision was made to carry out a PET-CT scan 6, 12, 24 and 36 months after surgery. Until now (36 months

Figure 4 Primary cervical tumour of 1997 with similar transition into gland-like structures and presence of hyperchromatic nuclei and intracytoplasmic mucin (arrows) (H&E, 250 $\times$).

Table 1 Summary of reported cases and current case of ovarian metastasis in a transposed ovary after primary cervical cancer

Authors	Age	Stage (FIGO)	Histologic type	Initial treatment	Time to recurrence	Treatment of recurrence	Follow-up
Nguyen <i>et al</i> ⁵	35 years	IB1	Squamous cell carcinoma	Vaginal hysterectomy, radical parametrectomy, pelvic lymphadenectomy, ovarian transposition	8 years	Exploratory laparotomy, BSO, radiotherapy	Recurrence after 4 months, palliative chemotherapy, lost to follow-up
Morice <i>et al</i> ⁶	Patient 1 34 years	IB1	Squamous cell carcinoma	Radical hysterectomy, pelvic and para-aortic lymphadenectomy, ovarian transposition, brachytherapy	3 years	Staging laparotomy as for ovarian cancer, BSO, right colectomy, chemotherapy	Died 15 months after recurrence
	Patient 2 34 years	IB1	Squamous cell carcinoma	Radical hysterectomy, pelvic lymphadenectomy, ovarian transposition, brachytherapy	3 years	Laparotomy, BSO, para-aortic lymphadenectomy, chemoradiation	Chemoradiation during publication
Delotte <i>et al</i> ⁷	36 years	IB1	Adenocarcinoma	Laparoscopic staging, ovarian transposition, pelvic radiation, brachytherapy, laparoscopic type II radical hysterectomy	2 years	Laparoscopic staging as for ovarian cancer, BSO.	5 years disease free
Janse <i>et al</i> (present case)	43 years	IB1	Adenosquamous carcinoma	Vaginal hysterectomy, radical parametrectomy, vaginal cuff, pelvic lymphadenectomy, ovarian transposition	10 years	Laparoscopic adnex extirpation	3 years disease free

BSO, bilateral salpingo-oophorectomy.

after removal of the ovary) the patient has demonstrated no evidence of recurrent disease.

DISCUSSION

Cases of cervical carcinoma metastasizing to the ovary are rarely reported in literature. The overall metastatic rate is estimated to vary between 0.9% and 2.2%, but different rates are seen when one looks at the histological

types specifically.^{1–3} Metastatic incidence for squamous cell carcinoma and adenocarcinoma is ranging between 0.4% and 1.3% and 1.7% and 8.2% respectively.^{1–3 8 9} Adenosquamous carcinoma was not taken into account separately but pooled with adenocarcinoma, as only very few cases were described. However, in the literature a controversy remains regarding a poorer prognosis of adenosquamous carcinoma in comparison with adenocarcinoma of

the uterine cervix.^{10–12} The variability between these studies might be due to poor diagnostic reproducibility.¹³

Independent risk factors for ovarian metastases in uterine cervical carcinoma were found to be the histologic type (adenocarcinoma),^{1–3} blood vessel invasion,⁹ uterine corpus involvement,³ unaffected peripheral stroma thickness (<3 mm)², age (>45 years)² and FIGO stage.² In the present case, only histologic type and blood vessel invasion can be determined as risk factors.

The four cases of ovarian metastasis on a transposed ovary described until now, all originated from FIGO stage IB1 cervical carcinoma (table 1).^{5–7} Morice *et al* reported two patients with squamous cell carcinoma who both presented 3 years after primary treatment with bilateral abdominal masses during clinical examination of 7 cm and 10 cm respectively.⁶ Delotte *et al* outlined one patient with adenocarcinoma who presented 2 years after primary treatment with abdominal pain and a 4.5 × 3 cm solid mass at MRI.⁷ Lastly, Nguyen *et al* described one patient with squamous cell carcinoma who presented 8 years after primary treatment with abdominal pain and a 6–7 cm solid mass during clinical examination and CT.⁵

The current case report is a unique case of an isolated ovarian metastasis. It is the first case describing a secondary ovarian malignancy originating from a cervical adenosquamous carcinoma in a transposed ovary. In addition, this is the first report of an ovarian metastasis 10 years after primary treatment for cervical cancer. Furthermore, immunohistologic and HPV genotyping played a key role in the diagnostic process, as the case did not raise suspicion by ultrasound findings neither by cytological examination after cytological aspiration nor by appearance during surgery.

Learning points

- ▶ Metastasis from primary cervical cancer in a transposed ovary is exceptionally rare.
- ▶ Metastasis from cervical cancer can occur in a transposed ovary up to 10 year after treatment of the primary tumour.
- ▶ Immunohistologic examination and HPV genotyping can be essential in the diagnostic process.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. Shimada M, Kigawa J, Nishimura R, *et al*. Ovarian metastasis in carcinoma of the uterine cervix. *Gynecol Oncol* 2006;**101**:234–7.
2. Landoni F, Zanagnolo V, Lovato-Diaz L, *et al*. Ovarian metastases in early-stage cervical cancer (IA2-IIA): a multicenter retrospective study of 1965 patients (a Cooperative Task Force study). *Int J Gynecol Cancer* 2007;**17**:623–8.
3. Kim MJ, Chung HH, Kim JW, *et al*. Uterine corpus involvement as well as histologic type is an independent predictor of ovarian metastasis in uterine cervical cancer. *J Gynecol Oncol* 2008;**19**:181–4.
4. de Waal YR, Thomas CM, Oei AL, *et al*. Secondary ovarian malignancies: frequency, origin, and characteristics. *Int J Gynecol Cancer* 2009;**19**:1160–5.
5. Nguyen L, Brewer CA, DiSaia PJ. Ovarian metastasis of stage IB1 squamous cell cancer of the cervix after radical parametrectomy and oophorectomy. *Gynecol Oncol* 1998;**68**:198–200.
6. Morice P, Haie-Meder C, Pautier P, *et al*. Ovarian metastasis on transposed ovary in patients treated for squamous cell carcinoma of the uterine cervix: report of two cases and surgical implications. *Gynecol Oncol* 2001;**83**:605–7.
7. Delotte J, Ferron G, Kuei TL, *et al*. Laparoscopic management of an isolated ovarian metastasis on a transposed ovary in a patient treated for stage IB1 adenocarcinoma of the cervix. *J Minim Invasive Gynecol* 2009;**16**:106–8.
8. Nakanishi T, Wakai K, Ishikawa H, *et al*. A comparison of ovarian metastasis between squamous cell carcinoma and adenocarcinoma of the uterine cervix. *Gynecol Oncol* 2001;**82**:504–9.
9. Yamamoto R, Okamoto K, Yukiharu T, *et al*. A study of risk factors for ovarian metastases in stage Ib–IIb cervical carcinoma and analysis of ovarian function after a transposition. *Gynecol Oncol* 2001;**82**:312–16.
10. Farley JH, Hickey KW, Carlson JW, *et al*. Adenosquamous histology predicts a poor outcome for patients with advanced-stage, but not early-stage, cervical carcinoma. *Cancer* 2003;**97**:2196–202.
11. Tiltman AJ. The pathology of cervical tumours. *Best Pract Res Clin Obstet Gynaecol* 2005;**19**:485–500.
12. Vinh-Hung V, Bourgain C, Vlastos G, *et al*. Prognostic value of histopathology and trends in cervical cancer: a SEER population study. *BMC Cancer* 2007;**7**:164.
13. Alfsen GC, Reed W, Abeler VM. Reproducibility of classification in non-squamous cell carcinomas of the uterine cervix. *Gynecol Oncol* 2003;**90**:282–9.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Janse JA, Sie-Go DMDS, Schreuder HWR. Ovarian metastasis in a transposed ovary 10 years after primary cervical cancer: the importance of histologic examination and review of literature. *BMJ Case Reports* 2011;**10**:1136/bcr.04.2011.4155, date of publication

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow