Tubular network formation protects mitochondria from autophagosomal degradation during nutrient starvation

Angelika S. Rambold, Brenda Kostecky, Natalie Elia, and Jennifer Lippincott-Schwartz¹

The Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD 20892

Contributed by Jennifer Lippincott-Schwartz, May 10, 2011 (sent for review April 19, 2011)

Mitochondria are highly dynamic organelles that mediate essential cell functions such as apoptosis and cell-cycle control in addition to their role as efficient ATP generators. Mitochondrial morphology changes are tightly regulated, and their shape can shift between small, fragmented units and larger networks of elongated mitochondria. We demonstrate that mitochondrial elements become significantly elongated and interconnected shortly after nutrient depletion. This mitochondrial morphological shift depends on the type of starvation, with an additive effect observed when multiple nutrients are depleted simultaneously. We further show that starvation-induced mitochondrial elongation is mediated by down-regulation of dynamin-related protein 1 (Drp1) through modulation of two Drp1 phosphorylation sites, leading to unopposed mitochondrial fusion. Finally, we establish that mitochondrial tubulation upon nutrient deprivation protects mitochondria from autophagosomal degradation, which could permit mitochondria to maximize energy production and supply autophagosomal membranes during starvation.

autophagy | mitofusin

Mitochondria are dynamic organelles that mediate many essential cell functions. Depending on the cellular context, mitochondria shift between fragmented and tubular network-like morphologies by means of coordinated fission and fusion (1, 2). Proteins responsible for mitochondrial fusion include the outer mitochondrial membrane proteins mitofusin 1 (Mfn1) and mitofusin 2 (Mfn2) (3–5) and the inner membrane protein optic atrophy 1 (Opa1) (6). Fission is mediated by dynamin-related protein 1 (Drp1) (7, 8) and its interaction with binding partners Fis1 and/or mitochondrial fission factor (Mff) (9, 10). Multiple mechanisms, including phosphorylation, sumoylation, and ubiquitination, coordinate Drp1 fission capacity (11–14). Phosphorylation of Drp1 at S616 by Cdk1/cyclin B results in increased Drp1 fission activity (13). Conversely, phosphorylation of Drp1 at S637 by PKA decreases fission by causing Drp1 retention in the cytosol, whereas dephosphorylation of S637 by calcineurin causes Drp1 translocation to the mitochondria and increased mitochondrial fission (11, 15).

Mitochondrial morphological dynamics are linked to regulation of many specific cell functions. Changes in mitochondrial cristae and mitochondrial fragmentation, for example, play a vital role in apoptosis (16, 17). Ca^{2+} transfer (18), cell-cycle regulation (13, 19), and mitochondrial quality control (20, 21) are all closely tied to changes in mitochondrial morphology. Furthermore, stress conditions and changes in energy source can induce significant mitochondrial morphological changes (22, 23). Very recently, nutrient starvation was shown to induce mitochondrial elongation and to protect mitochondria from autophagic degradation (24). In addition to the above functions, mitochondria have recently been linked to autophagosome biogenesis during starvation conditions (25), and it is possible that mitochondrial morphological changes play a role in this process.

Autophagy is a catabolic process that proceeds via formation of a double-membrane vesicle, the autophagosome. Both cyto-

solic and organellar components can be engulfed by autophagosomes, which then fuse with lysosomes. Once delivered to the lysosome, autophagosomal contents are recycled, allowing prolonged cell survival when nutrients are scarce (26). Recent research shows that the autophagy machinery can target specific substrates for degradation depending on the cell's needs in specific contexts such as mitochondrial stress (27, 28). Autophagy induced by starvation, however, has traditionally been viewed as a nonspecific process in which cytosolic and organellar components are turned over en masse. Cells become highly dependent on autophagy when nutrients are severely limited because autophagy allows for replenishment of vital macromolecular precursors, including amino acids, sugars, and fatty acids. These breakdown products provide substrates for biosynthesis of core cell components and metabolic substrates for the ATP production required to drive macromolecule synthesis during starvation. Despite a presumed need under starvation conditions for efficient ATP production provided by mitochondria, it has been unclear whether changes in mitochondrial morphology occur to facilitate promotion of cell survival and how those dynamic changes in mitochondrial structure might be mediated.

Here, we address these questions by examining mitochondrial morphology under nutrient deprivation. Our results demonstrate that mitochondrial elongation is induced shortly after starvation. This mitochondrial tubulation is reversible and depends on the specific nutrients depleted. Mitochondrial elongation does not occur when autophagy is induced by serum depletion or glucose elimination. Rather, mitochondrial elongation is a specific response to deprivation of glutamine and/or amino acids, with faster, more extensive mitochondrial tubulation upon additional depletion of glucose and serum. We demonstrate that starvation-induced mitochondrial elongation is mediated by down-regulation of Drp1, leading to unopposed mitochondrial fusion. The mitochondrial elongation response is further shown to protect mitochondria from autophagic turnover under starvation conditions. This ability of autophagosomes to exclude mitochondria from autophagic sequestration during starvation may serve to provide efficient aerobic ATP production when energy sources are limited and also highlights the previously unappreciated substrate specificity of the autophagosomal pathway during nutrient deprivation.

Results

Mitochondria Elongate During Cell Starvation. To determine whether mitochondria undergo morphological changes upon starvation, we imaged mouse embryonic fibroblasts (MEFs) ex-

Author contributions: A.S.R. and J.L.-S. designed research; A.S.R., B.K., and N.E. performed research; A.S.R., B.K., and J.L.-S. analyzed data, and A.S.R., B.K., and J.L.-S. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: lippincj@mail.nih.gov.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1107402108/-DCSupplemental.

Fig. 2. Mitochondrial tubulation is more efficiently induced by additive nutrient deprivation. MEFs were transfected with mito-YFP and starved for 2 h (DPBS, D-GSG, and no Ser) or 6 h (no glc, low glc, no aa, no gln, and D-GG). Media compositions are given in the table. Mitochondrial morphology was scored as described in Fig. 1 ($n \geq 3$). Representative images of cells are shown in the first column. Images in the second column are magnified views of the boxed areas in the first column. glc, glucose; gln, glutamine; aa, amino acids. (Scale bar: 15 μ m.)

Because Drp1 is regulated by phosphorylation, we sought to assess the potential involvement of Drp1 phosphorylation sites in starvation-induced mitochondrial elongation. Drp1-S616 phosphorylation decreased upon starvation (Fig. 3C) as determined by immunoblotting with a phospho-S616-specific antibody. This decreased Drp1-S616 phosphorylation is consistent with an increase in mitochondrial elongation attributable to inhibited Drp1 fission activity. A phosphomimetic mutant (Drp1-S616E) over-

expressed in MEFs led to less mitochondrial elongation than observed in cells transfected with WT Drp1 (wtDrp1) during starvation (Fig. 3B and Fig. S3B). Overexpression of a non-phosphorylatable Drp1-S616A mutant, on the other hand, did not affect mitochondrial elongation after starvation (Fig. S3C).

We next investigated the effect of a second posttranslational modification on Drp1-mediated mitochondrial fission during starvation. Overexpression of Drp1-S637A in MEFs resulted in sig-

Fig. 3. Unopposed fusion mediates mitochondrial tubulation during starvation. (A, C, and D) MEFs were treated with the media indicated for 2.5 h (A) or 5 h (C and D) and lysed in ice-cold lysis buffer A, and samples were subjected to immunoblotting against Mfn1, Mfn2, Opa1, Drp1, and pS616-Drp1 as indicated (whole-cell lysates were probed in A and C; isolated mitochondria were probed in D). (B) At 16 h after transfection with the indicated YFP-tagged Drp1 constructs, cells were treated with full medium or starved in D-GSG for 2 h. Mitochondrial morphology was scored as described in Fig. 1 ($n = 3$).

nificantly fewer cells with elongated mitochondria compared with wtDrp1-transfected cells after starvation (Fig. 3*B* and Fig. S3*B*). This finding is consistent with a report showing that phosphorylation of Drp1-S637 results in cytosolic retention of Drp1 and a decrease in mitochondrial fission (15). Overexpression of the double mutant Drp1-S616E/S637A during starvation resulted in a statistically significant decrease in mitochondrial tubulation compared with either Drp1-S616E- or Drp1-S637A-transfected cells, indicating that a synergistic effect of the two posttranslational modifications is possible (Fig. 3*B* and Fig. S3*D*).

A comparison of Drp1 levels associated with mitochondria revealed a substantial decrease in mitochondrial localization of Drp1 upon starvation, whereas total Drp1 levels remained constant (Fig. 3*A* and *D*), suggesting a reduction in the ability of Drp1 to effect fission after nutrient depletion. Together, these data suggest that starvation inhibits mitochondrial fission both by decreasing Drp1 fission activity and by preventing Drp1 localization to mitochondria, leading to unopposed Opa1/Mfn1-dependent mitochondrial fusion.

Elongation Protects Mitochondria from Mitophagy. Mitochondrial fragmentation and dysfunction have been described as preconditions for autophagy of mitochondria (mitophagy) under different conditions (21, 27, 36), which led us to investigate whether starvation-induced mitochondrial fusion can prevent mitochondria from being targeted for autophagy. We did not observe a decrease in mitochondrial mass during a 5-h starvation in WT cells, consistent with our previously reported data (25). However, mitochondrial fusion-deficient Opa1KO cells showed a significant decrease in mitochondrial proteins Tom20, Hsp60, and Complex V (subunit α) under starvation conditions (Fig. 4*A*). Furthermore, autophagosome-engulfed mitochondria were observed in nutrient-deprived Opa1KO cells containing fragmented mitochondria but not in WT cells with elongated mitochondria (Fig. S4).

To more clearly establish whether elongation can protect mitochondria from autophagy, we investigated whether high levels of mitophagy could be reversed by starvation-induced mitochondrial fusion. Here, we took advantage of the fact that serum starvation induces autophagy without inducing mitochondrial fusion (Fig. 2). First, we quantified mitophagy levels in WT and Mfn2KO MEFs transfected with GFP-LC3 and mitochondrially targeted red fluorescent protein (mito-RFP) using live-cell imaging. For better identification of mitochondrial material engulfed in autophagosomes, we blocked lysosomal turnover with bafilomycin A1. A lower incidence of mitophagy was observed in serum-starved WT cells compared with Mfn2KO cells (Fig. 4*B*). Additionally, there was a large subpopulation (29% of cells) in the starved Mfn2KO cells with >7 mitochondria-containing autophagosomes (approximately 5% in WT cells). The increased incidence of mitophagy could suggest that mitochondria from Mfn2-deficient cells are more prone to dysfunction because of the lack of endoplasmic reticulum-mitochondria tethering and altered calcium handling (34) and is consistent with evidence that serum starvation is associated with enhanced turnover of dysfunctional mitochondria (37). Next, we investigated whether the high levels of mitophagy in serum-starved Mfn2KO cells could be reduced if mitochondrial fusion was induced through glutamine depletion and glucose reduction. Indeed, although 29% of serum-starved Mfn2KO cells had >7 mitochondria in autophagosomes, only 1% of the D-GSG-starved cells contained >7 engulfed mitochondria (Fig. 4*B*). These findings support the hypothesis that mitochondrial elongation has a strong capacity to prevent mitochondrial turnover during starvation-induced autophagy.

Discussion

Mitochondrial morphological changes have been associated with many vital cellular functions. Mitochondrial fragmentation is important for progression of apoptosis (38, 39) and M phase (40)

Fig. 4. Mitochondrial tubulation ameliorates autophagosomal degradation of mitochondria during starvation. (A) WT and Opa1KO MEFs were treated with the indicated media for 5 h, and equal numbers of cells were lysed. Whole-cell lysates were subjected to immunoblotting against the proteins indicated. (B) Mfn2KO cells show increased mitophagy during serum starvation, which can be rescued through mitochondrial tubulation. WT and Mfn2KO MEF cells were transfected with GFP-LC3 and mito-RFP and starved for serum or D-GSG for 4 h in the presence of bafilomycin A1, and live images were acquired. Cells were grouped into categories according to the number of mitochondria in autophagosomes and are presented as percentage of the total number of transfected cells counted ($n = 3$). (Scale bar: 15 μ m.)

and for degradation of defective mitochondria (21). The number of processes involving mitochondrial fusion is equally diverse, underscoring the importance of understanding how changes in mitochondrial morphology are related to specific outcomes in the cell. Several lines of evidence, including the data presented here, indicate that mitochondrial fusion can serve a protective function, leading to the exchange of mitochondrial DNA, reorganization of mitochondrial cristae, and delay in apoptosis (22, 38, 39, 41). It is therefore reasonable that such mitochondrial morphological changes could also be beneficial for cells under conditions of low nutrient supply. Precedent for such energy source-induced mitochondrial morphological change was established on the observation that cells grown with galactose/glutamine instead of glucose resulted in more interconnected mitochondria with increased cristae (23). Here we demonstrate that mitochondria undergo rapid Mfn1- and Opa1-dependent elongation, mediated by two Drp1 posttranslational modifications after nutrient depletion (Fig. 5). The extent and rate of mitochondrial elongation increases upon additive nutrient starvation, with elongation protecting mitochondria from sequestration and subsequent degradation by autophagosomes.

Starvation-induced mitochondrial tubulation requires both Opa1 and Mfn1. Although Mfn2 appears capable of providing slight redundancy for Mfn1 fusion activity, the lack of a strong mitochondrial fusion phenotype in Mfn1KO cells indicates Mfn2 alone is

Protein concentrations were measured with bicinchoninic acid (Pierce) according to the manufacturer's instructions, and 10 μ g of protein was loaded on 4–12% NUPAGE Gels (Invitrogen). Primary antibodies were added for 16 h at 4 °C and secondary antibodies for 1 h at 24 °C. Antibody binding was detected with SuperSignal West Pico Chemiluminescent Substrate (Pierce).

Imaging and Statistics. Cells were stained as described previously (47). Cells were seeded either in LabTek chambers (Thermo Scientific) or on coverslips for live-cell imaging and immunofluorescence, respectively. Live and fixed cells were imaged on an LSM 510 META Zeiss microscope using a 63 $\times$ oil

objective at 37 °C. Images were analyzed with Karl Zeiss MicroImaging Software. Brightness and contrast were adjusted in Adobe Photoshop CS. Data were expressed as means $\pm$ SD. Statistical analysis among groups was performed using Student's *t* test.

ACKNOWLEDGMENTS. We thank R. Youle for contributing the Bax/BakDKO cells and the YFP-Drp1 construct and N. Mizushima and D. C. Chan for contributing the ATG5KO and Mfn-deficient cell lines, respectively. We thank K. Mitra for helpful discussions. A.R. was supported by a postdoctoral fellowship from the Deutsche Forschungsgemeinschaft.

- Bereiter-Hahn J (1990) Behavior of mitochondria in the living cell. *Int Rev Cytol* 122: 1–63.
- Nunnari J, et al. (1997) Mitochondrial transmission during mating in *Saccharomyces cerevisiae* is determined by mitochondrial fusion and fission and the intramitochondrial segregation of mitochondrial DNA. *Mol Biol Cell* 8:1233–1242.
- Santel A, et al. (2003) Mitofusin-1 protein is a generally expressed mediator of mitochondrial fusion in mammalian cells. *J Cell Sci* 116:2763–2774.
- Santel A, Fuller MT (2001) Control of mitochondrial morphology by a human mitofusin. *J Cell Sci* 114:867–874.
- Chen H, et al. (2003) Mitofusins Mfn1 and Mfn2 coordinately regulate mitochondrial fusion and are essential for embryonic development. *J Cell Biol* 160:189–200.
- Meeusen S, et al. (2006) Mitochondrial inner-membrane fusion and crista maintenance requires the dynamin-related GTPase Mgm1. *Cell* 127:383–395.
- Otsuga D, et al. (1998) The dynamin-related GTPase, Dnm1p, controls mitochondrial morphology in yeast. *J Cell Biol* 143:333–349.
- Bleazard W, et al. (1999) The dynamin-related GTPase Dnm1 regulates mitochondrial fission in yeast. *Nat Cell Biol* 1:298–304.
- Yoon Y, Krueger EW, Oswald BJ, McNiven MA (2003) The mitochondrial protein hFis1 regulates mitochondrial fission in mammalian cells through an interaction with the dynamin-like protein DLP1. *Mol Cell Biol* 23:5409–5420.
- Otera H, et al. (2010) Mff is an essential factor for mitochondrial recruitment of Drp1 during mitochondrial fission in mammalian cells. *J Cell Biol* 191:1141–1158.
- Cribbs JT, Strack S (2007) Reversible phosphorylation of Drp1 by cyclic AMP-dependent protein kinase and calcineurin regulates mitochondrial fission and cell death. *EMBO Rep* 8:939–944.
- Harder Z, Zunino R, McBride H (2004) Sumo1 conjugates mitochondrial substrates and participates in mitochondrial fission. *Curr Biol* 14:340–345.
- Taguchi N, Ishihara N, Jofuku A, Oka T, Mihara K (2007) Mitotic phosphorylation of dynamin-related GTPase Drp1 participates in mitochondrial fission. *J Biol Chem* 282: 11521–11529.
- Yonashiro R, et al. (2006) A novel mitochondrial ubiquitin ligase plays a critical role in mitochondrial dynamics. *EMBO J* 25:3618–3626.
- Cereghetti GM, et al. (2008) Dephosphorylation by calcineurin regulates translocation of Drp1 to mitochondria. *Proc Natl Acad Sci USA* 105:15803–15808.
- Scorrano L, et al. (2002) A distinct pathway remodels mitochondrial cristae and mobilizes cytochrome c during apoptosis. *Dev Cell* 2:55–67.
- Youle RJ, Karbowski M (2005) Mitochondrial fission in apoptosis. *Nat Rev Mol Cell Biol* 6:657–663.
- Park MK, Ashby MC, Erdemli G, Petersen OH, Tepikin AV (2001) Perinuclear, perigranular and sub-plasmalemmal mitochondria have distinct functions in the regulation of cellular calcium transport. *EMBO J* 20:1863–1874.
- Mitra K, Wunder C, Roysam B, Lin G, Lippincott-Schwartz J (2009) A hyperfused mitochondrial state achieved at G1-S regulates cyclin E buildup and entry into S phase. *Proc Natl Acad Sci USA* 106:11960–11965.
- Chen H, Chomyn A, Chan DC (2005) Disruption of fusion results in mitochondrial heterogeneity and dysfunction. *J Biol Chem* 280:26185–26192.
- Twig G, et al. (2008) Fission and selective fusion govern mitochondrial segregation and elimination by autophagy. *EMBO J* 27:433–446.
- Tondera D, et al. (2009) SLP-2 is required for stress-induced mitochondrial hyperfusion. *EMBO J* 28:1589–1600.
- Rosignol R, et al. (2004) Energy substrate modulates mitochondrial structure and oxidative capacity in cancer cells. *Cancer Res* 64:985–993.
- Gomes LC, Benedetto GD, Scorrano L (2011) During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. *Nat Cell Biol* 13: 589–598.
- Hailey DW, et al. (2010) Mitochondria supply membranes for autophagosome biogenesis during starvation. *Cell* 141:656–667.
- Rabinowitz JD, White E (2010) Autophagy and metabolism. *Science* 330:1344–1348.
- Narendra D, Tanaka A, Suen DF, Youle RJ (2008) Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol* 183:795–803.
- Schweers RL, et al. (2007) NIX is required for programmed mitochondrial clearance during reticulocyte maturation. *Proc Natl Acad Sci USA* 104:19500–19505.
- Mizushima N, et al. (2001) Dissection of autophagosome formation using Apg5-deficient mouse embryonic stem cells. *J Cell Biol* 152:657–668.
- Song Z, Chen H, Fiket M, Alexander C, Chan DC (2007) OPA1 processing controls mitochondrial fusion and is regulated by mRNA splicing, membrane potential, and Yme1L. *J Cell Biol* 178:749–755.
- Wei MC, et al. (2001) Proapoptotic BAX and BAK: A requisite gateway to mitochondrial dysfunction and death. *Science* 292:727–730.
- Hoppins S, et al. (2011) The soluble form of Bax regulates mitochondrial fusion via MFN2 homotypic complexes. *Mol Cell* 41:150–160.
- Karbowski M, Norris KL, Cleland MM, Jeong SY, Youle RJ (2006) Role of Bax and Bak in mitochondrial morphogenesis. *Nature* 443:658–662.
- de Brito OM, Scorrano L (2008) Mitofusin 2 tethers endoplasmic reticulum to mitochondria. *Nature* 456:605–610.
- Ishihara N, Eura Y, Mihara K (2004) Mitofusin 1 and 2 play distinct roles in mitochondrial fusion reactions via GTPase activity. *J Cell Sci* 117:6535–6546.
- Gomes LC, Scorrano L (2008) High levels of Fis1, a pro-fission mitochondrial protein, trigger autophagy. *Biochim Biophys Acta* 1777:860–866.
- Elmore SP, Qian T, Grissom SF, Lemasters JJ (2001) The mitochondrial permeability transition initiates autophagy in rat hepatocytes. *FASEB J* 15:2286–2287.
- Frank S, et al. (2001) The role of dynamin-related protein 1, a mediator of mitochondrial fission, in apoptosis. *Dev Cell* 1:515–525.
- Lee YJ, Jeong SY, Karbowski M, Smith CL, Youle RJ (2004) Roles of the mammalian mitochondrial fission and fusion mediators Fis1, Drp1, and Opa1 in apoptosis. *Mol Biol Cell* 15:5001–5011.
- Yaffe MP (1999) The machinery of mitochondrial inheritance and behavior. *Science* 283:1493–1497.
- Chen H, et al. (2010) Mitochondrial fusion is required for mtDNA stability in skeletal muscle and tolerance of mtDNA mutations. *Cell* 141:280–289.
- Karbowski M, Neutznar A, Youle RJ (2007) The mitochondrial E3 ubiquitin ligase MARCH5 is required for Drp1 dependent mitochondrial division. *J Cell Biol* 178:71–84.
- Nakamura N, Kimura Y, Tokuda M, Honda S, Hirose S (2006) MARCH-V is a novel mitofusin 2- and Drp1-binding protein able to change mitochondrial morphology. *EMBO Rep* 7:1019–1022.
- Kristensen AR, et al. (2008) Ordered organelle degradation during starvation-induced autophagy. *Mol Cell Proteomics* 7:2419–2428.
- Iwata J, et al. (2006) Excess peroxisomes are degraded by autophagic machinery in mammals. *J Biol Chem* 281:4035–4041.
- Collins CA, et al. (2009) Atg5-independent sequestration of ubiquitinated mycobacteria. *PLoS Pathog* 5:e1000430.
- Rambold AS, et al. (2006) Association of Bcl-2 with misfolded prion protein is linked to the toxic potential of cytosolic PrP. *Mol Biol Cell* 17:3356–3368.