Dinucleotide repeat polymorphism at the human dopamine β -hydroxylase (DBH) locus

C.J.Porter, J.Nahmias¹, J.Wolfe¹ and I.W.Craig
Genetics Laboratory, Department of Biochemistry, South Parks Road, Oxford, OX1 3QU and ¹The Galton Laboratory, University College London, 4 Stephenson Way, London NW1 2HE, UK

Source/Description: Cosmid clone cDBH14 was isolated by screening a human cosmid library (1) with an 800bp *Bam*HI/*Eco*RI fragment derived from a DBH cDNA clone (2). A 1kb *Pst*I subclone of cosmid cDBH14 was shown to hybridize to the probe (GT)₁₅, and was sequenced outwards from the repeat using the primers (GT)₇-A/C/T-N and (AC)₇-C/G/T-N (3). In addition, a phage clone, λ D8.1, was selected from a human genomic library by PCR and hybridization to a DBH cDNA probe (2). A 1kb *Hind*III subclone of phage λ D8.1 was shown to hybridize to poly(CA) and was fully sequenced. The phage sequence has been submitted to the EMBL data library with the accession number X63418.

Primer Sequences:

GCAGTCACGCATCCTTATGG (GT strand).
CAGCTCTGGGCTCATGCTC (AC strand).

Frequency: Estimated from 224 chromosomes from unrelated individuals. PIC = 0.47, observed heterozygosity = 0.51.

Allele F1 (180bp)	0.013
Allele F2 (178bp)	0.093
Allele F3 (176bp)	0.265
Allele F4 (174bp)	0.625
Allele F5 (172bp)	0.004

Chromosomal Localization: DBH has been localized to 9q34 by linkage analysis (4) and *in situ* hybridization (5).

Mendelian Inheritance: Consistent with Mendelian inheritance in two 3-generation and two 2-generation families.

PCR Conditions/Comments: Amplification was carried out using the following conditions: 30 cycles of 1 minute at 94°C, 45 seconds at 59°C and 25 seconds at 70°C. Alleles were separated by electrophoresis on 8% polyacrylamide. The (GT) strand corresponds to the +strand of the DBH gene.

Acknowledgements: We are grateful to Jacques Mallet who provided the DBH cDNA clone, to Derek Blake for the phage library and to Graeme Black for some of the family DNAs. This work was carried out with support from the MRC as part of the UK human genome mapping project (MRC grant 8916330), and from the Wellcome Trust.

References: 1) Cachon-Gonzales, M.B. (1991) PhD Thesis. University of London. 2) Lamouroux, A. *et al.* (1987) *EMBO J.* **6**, 3931–3937. 3) Browne, D.L. (1991) *Cytogenet. Cell Genet.* **58**, 2147. 4) Wilson, A.F. *et al.* (1988) *Am. J. Hum. Genet.* **42**, 160–166. 5) Craig, S.P. *et al.* (1988) *Cytogenet. Cell Genet.* **48**, 48–51.

RFLP for Bgl II at the human neurofilament medium chain (NEF3) gene locus

Yuan Ding, Danielle R.Reed, Mary C.Baltazar and R.Arlen Price*

Behavioral Genetics Laboratory, Department of Psychiatry, University of Pennsylvania, 422 Curie Blvd, CRB-118, Philadelphia, PA 19104-6141, USA

Source/Description: pHNFM contains a 8.5 kb *Bam*HI-*Hind* III fragment that encodes the entire human NEF3 gene (1), the vector is EMBL3 and a 726 bp *Bam*HI fragment, corresponding to a part of exon 3, was used as a probe.

Polymorphism: Bgl II identifies a two-allele polymorphism with fragment sizes of either 6.9 kb or 6.4 kb.

Frequency: Studied in 30 unrelated individuals.

A1	6.9 kb allele	0.52
A2	6.4 kb allele	0.48

Observed heterozygosity = 0.56.

Not Polymorphic For: *Bam*HI, *Hind* III, *Eco*RI, *Pvu* II, *Hae* III, *Hha* I, *Taq* I, *Bcl*, *Msp* I.

Chromosomal Localization: Localized to 8p21 (D.Nelson, personal communication).

Mendelian Inheritance: Co-dominant segregation shown in 5 informative families, with a total of 32 individuals.

Probe Availability: Provided by Dr.V.M.-Y.Lee (1).

Other Comments: Low background under usual stringency conditions.

Acknowledgements: This work was supported in part by NIH grants MH43880, DK44073 and MH44210.

References: 1) Myers *et al.* (1987) *EMBO J.* **6**, 1617–1626.

* To whom correspondence should be addressed